Progress reports

Sea Levels: Abrupt events and mechanisms of change

Robin Edwards

School of Natural Sciences, Trinity College Dublin, Ireland.

[**Citation:** Edwards, R.J. (2005) Sea levels: abrupt events and mechanisms of change *Progress in Physical Geography* **29(4)** 599-608. DOI: 10.1191/0309133305pp465pr]

International Geoscience Programme (IGCP) Project 495 'Quaternary Land-Ocean Interactions: Driving Mechanisms and Coastal Responses' commenced in 2004, three decades after the first project specifically concerned with sea levels began (Project 61: 1974-82). Over thirty years of co-ordinated international research have produced a detailed picture of the spatial variability inherent in sea level change. Project 495 will undoubtedly refine this picture but in addition, it seeks to develop a more detailed understanding of the mechanisms responsible for these patterns. This report reviews examples of recent research concerned with identifying underlying mechanisms, and concentrates on the role of studies examining abrupt events and rapid changes.

I. Tsunamis

The last couple of years have seen a number of papers and special issues concerned with tsunamis (e.g. Tappin, 2004; Pelinovsky & Tinti, 2005). This subject has particular resonance in light of the tragic loss of life associated with the Indian Ocean tsunami of December 2004. The nature and extent of a tsunami's impact is

related to the mechanism responsible for its generation (e.g. seismic event, mass movement etc.). Consequently, this research naturally progresses from identification of deposits to an examination of the events responsible for them. This requires the linkage of terrestrial and marine records, and involves modern observations, palaeoenvironmental reconstructions and computer model simulations.

1. Modern Observations

Tsunamis leave deposits that are 'out of place', such as marine sand layers within a terrestrial peat or freshwater lake (e.g. Dawson et al., 1991; Bondevik et al., 1997a, b.). In themselves, these attributes are not unique, since storm surges can also deposit marine sediments many metres above normal tidal levels (Dawson & Shi, 2000). Ongoing research is investigating the processes and impacts of modern tsunamis to develop a clearer picture of the distinctive anatomy of these events (e.g. Maramai et al., 2005a, b). Comparative studies examining both storm and tsunami deposits are particularly valuable in this respect (e.g. Nanayama et al., 2000). A recent example is the study by Goff et al. (2004) in New Zealand, who examine sediments from a 15th century tsunami and a large storm that occurred in 2002. The results show clear differences in their sedimentology, continuity and extent. This type of information enables the development of a set of diagnostic criteria for the identification of palaeotsunami deposits (e.g. Goff et al., 2001; Smith et al., 2004; Bondevik et al., 2005; Williams et al., 2005).

In contrast to their sandy counterparts, rocky shorelines currently lack detailed sedimentary facies models describing the processes responsible for the formation of coarse deposits (Felton & Crook, 2003). Consequently, interpretations from these environments are more equivocal, relying on supporting evidence and argument (e.g. McMurtry et al., 2004; Scheffers, 2004). This situation is complicated by new data

indicating that the influence of storm waves may be greater than previously thought. Whilst attempts have been made to model the size of wave required to lift blocks of a given mass (e.g. Nott, 2003), recent studies of modern deposits suggest these significantly overestimate the size of the waves involved (e.g. Mastronuzzi & Sanso, 2004). For example, Williams & Hall (2004) graphically illustrate the power of storm waves in their description of megaclasts from cliff-tops along the Atlantic Irish coast. Here, boulders weighing several tonnes are recorded up to 50 m above sea level. It is likely that as more data are collected, the power of storm waves will be reassessed, and this may prompt the reinterpretation of some purportedly tsunamigenic deposits.

2. Palaeoenvironmental reconstructions

The number, distribution and height of waves associated with a tsunami reflect the mechanism responsible for its formation. For example, in the case of a fault-related tsunami, runup height rarely exceeds twice the fault slip (Okal & Synolakis, 2004). Consequently, given maximum water levels of 25-30 m recorded in Sumatra, Stein & Okal (2005) suggest the recent Indian Ocean event was triggered by a fault displacement of between 12 and 15 m. Mass movements have the potential to produce even bigger waves, but at present these remain difficult to quantify, particularly when the slide or slump is submarine in nature (Pelinovsky & Tinti, 2005). This picture is further complicated when mass movements are triggered by seismic events. Investigations of past tsunamis can be used to develop and test models of tsunami generation, and identify regions at greatest risk of future events.

Smith et al. (2004) combine data from 32 sites in northern Britain, to provide a comprehensive review of evidence for the Storegga Slide tsunami. This evidence predominantly takes the form of a widespread 'out of place' sand layer containing marine microfossils and radiocarbon dated to c. 8000 cal. yr BP. Sedimentological analysis of these deposits provides information on the anatomy of this intensively studied palaeotsunami. These results confirm that the slide was actually associated with more than one wave, supporting earlier evidence of multiple waves inferred from deposits on the Shetland Isles (Bondevik et al., 2003). Tooley & Smith (2005) show that this signal is robust even within a high-energy setting, describing two fining-up sequences in coarse deposits of sand and gravel from Scotland.

Reconstructing runup height from a palaeotsunami requires information on the maximum altitude attained by the waves, and the altitude of sea level at the time of the event. Runup heights based on the inland limit of sand layers are treated as minimum estimates since water levels commonly exceed the height of sediment deposition (Dawson, 1999; Dawson & Shi, 2000). Many studies rely on estimates of former relative sea-level derived from geophysical models of the glacial-isostatic adjustment process (e.g. Bondevik et al., 2003). In an interesting development, Smith et al. (2004) use the tsunami deposits as a time horizon. By locating the inland limit of inter-tidal sediments capped by this marker, the shoreline position at the time of the tsunami can be reconstructed and subsequent patterns of land movement identified. A variable pattern of runup height is revealed which Smith et al. (2004) attribute to wave modification by the coastline, and differences in the state of the tide at the time of the event.

Accurate dating is central to all reconstructions and is particularly important where the timing of the event is considered a diagnostic feature. In common with many sea

level investigations, assigning a precise year to a tsunami deposit is complicated due to the difficulties associated with constructing precise chronologies from sedimentary sequences (Edwards, 2003). In the case of the Storegga Slide deposits, whilst there is no shortage of radiocarbon dates, the results show typical ranges of several hundred years due to a combination of problems such as erosion and the uncertainties inherent in the radiocarbon technique. Age estimates also vary depending on the dating method used, and Bondevik et al. (2003, in press) note that conventional 'bulk' radiocarbon dates are commonly too young due to contamination. Whilst the precise year is uncertain, fish skeletons and plant macrofossils preserved within Storegga slide deposits suggest that the tsunami occurred in late autumn (Bondevik et al., 1997; Dawson & Smith, 2000).

Whilst dating sediments can constrain the timing of an undated event, it may also be used to identify a tsunami deposit by association with a known occurrence. Williams et al. (2005) attempt to fingerprint the sources of nine muddy sand layers preserved within a tidal marsh in Washington State, USA. They compare the ages of the layers with historical events and other dated tsunami deposits in the region. The authors conclude that four to six of the layers are probably the result of tsunamis, but that these are likely to have been generated by a variety of sources, involving both mass movements and seismic events from near and far-field locations. Williams et al. (2005) note that this process is complicated by the fact that the uncertainties associated with their age data are of similar magnitude to the recurrence interval of events in this region.

3. Model simulation and marine data

As well as providing estimates of past relative sea-levels, computer models are used to simulate the actual slides responsible for generating tsunamis and the patterns of wave propagation expected to result from different mechanisms of generation (McMurty et al., 2004; Fryer et al., 2004; Løvholt et al., 2005). For example, Bondevik et al. (2005) compare the extensive set of runup data from Norway, Scotland and the Shetland Islands, with numerical simulations of the Storegga slide. Their best-fit model suggests that sea levels along the Norwegian coast fell by around 20 m during the first 30 minutes following the slide. The simulation also predicts the generation of multiple waves, matching the reconstructions based on sedimentological data. In an alternative application, Okal (2005) uses a model to fingerprint the source of the 1906 Pacific-wide tsunami. Whilst the influence of the two candidate earthquakes is indistinguishable in terms of time (they occurred within 30 minutes of each other), the simulated tsunamis have distinctly different far-field characteristics, indicating that the Chilean event was in fact the most likely cause.

Additional information on tsunami source is provided by marine seismic data. The location, extent and architecture of the Storegga slide is now well-constrained by geophysical surveys (Haflidason et al., 2004). The triggering mechanism is still under investigation, and Bryn et al. (2005) suggest a strong earthquake is a likely cause. This may have been facilitated by excess porewater pressure in the sediments brought about by high rates of deposition. Solheim et al. (2005) suggest that immediately following deglaciation, rapid sedimentation coupled with glacio-isostatic seismicity could produce conditions favourable for slope failure. They report seven large pre-Holocene slides in the area that appear to form a complex of failures related to the glacial-interglacial cycle. The recent discovery on the Shetland Islands of two tsunami-like deposits post-dating the Storegga Slide suggests that this instability persists throughout the Holocene (Bondevik et al., in press). Hutton &

Syvitskli (2004) model sediment failures under changing sea levels, and note that whilst most occur during sea level falls or lowstands, the largest volume failures are associated with rising sea levels and highstands.

In fact, the nature of coastal sedimentation is increasingly being tied with tsunami risk. Using a catalogue of historic tsunamis in the Pacific Ocean, Gusiakov (2005) shows that sedimentation has a strong control on the likelihood that an earthquake will generate a tsunami. In this paper, written almost a year before the Indian Ocean tsunami of 2004, the extremely high efficiency of earthquakes in the western part of Indonesia is highlighted. In a recent paper, Syvitski et al. (2005) use a model to examine how the flux of sediment into the global coastal ocean has changed due to human activities. One of their results is that, in contrast to many other parts of the globe, Indonesian rivers now deliver much more sediment to their coastal waters than before. This research highlights the need for a greater understanding of land-ocean sediment fluxes and processes operating on the shelf (Long, 2003).

II. Sea levels and seismic events

In addition to tsunamis, earthquakes are associated with relative sea-level changes driven by vertical land movement. Evidence for cycles of land uplift and subsidence have been known in coastal deposits from the Pacific coast of North America for some years (e.g. Atwater, 1987; Atwater & Hemphill-Haley, 1997). Of current interest is the observation that in some areas a period of pre-seismic subsidence occurs which could potentially be used as an indicator of forthcoming large earthquakes (Shennan et al., 1999).

Distinguishing pre-seismic subsidence from co-seismic change requires detailed investigations employing precise indicators of relative sea-level. Quantitative paleoenvironmental reconstructions employing microfossils have the potential to provide these records (e.g. Sawai et al. 2004a). Zong et al. (2003) used diatoms and pollen to examine relative sea-level changes at two sites in Cook Inlet, Alaska, resulting from the 1964 earthquake. Their data, with a maximum precision of ± 0.06 m, indicate pre-seismic subsidence of around 0.15 m at both locations. In two related papers, Hamilton & Shennan (2005; in press) also use a diatom-based transfer function for tide level to reconstruct changes associated with a series of seismic events in the area. Their results show evidence for pre-seismic subsidence of similar magnitude to that associated with the 1964 earthquake, although these subtle changes are at the upper limit of their transfer function's resolution.

Elsewhere Hayward et al. (2004) use a combination of foraminifera and diatoms to identify sudden elevation changes in three Holocene sedimentary sequences from Ohiwa Harbour, New Zealand. Their records indicate potential seismic related subsidence of around 2m, although the reconstructions are of lower precision than the transfer functions from North America. In Japan, Sawai et al. (2004b) use a diatom-based transfer function to reconstruct relative sea-level changes in eastern Hokkaido. Instrumental data from this area record recent land subsidence, whilst geological evidence, such as elevated marine terraces, indicates long-term net uplift. Sawai et al. (2004b) focus their attention on a large earthquake and tsunami that occurred in the 17th century. Detailed analysis above and below the tsunami sand layer indicates gradual pre-seismic subsidence followed by post-seismic uplift. Intriguingly, this post-seismic uplift appears to have persisted for several decades rather than occurring abruptly. The authors conclude that a more detailed picture of

earthquake deformation cycles in this area is necessary before the long-term balance between inter-seismic subsidence and post-seismic uplift can be determined.

III. Meltwater Pulses

Abrupt changes in sea level have occurred during the late Glacial and Holocene in response to injections of meltwater from terrestrial ice sheets. Work is ongoing to constrain the timing, magnitude and source of these meltwater pulses (MWP). In Ireland, Clark et al. (2005) present five AMS radiocarbon dates from foraminifera within stratified glaciomarine sediments which they use to support the idea of a rapid sea level rise associated with a MWP around 19 000 cal. years BP. The existence of this pulse has been the subject of some debate since its earlier identification in marine sediments from Bonaparte Gulf, Australia (Yokohama et al., 2000, 2001, 2003; Clark & Mix, 2002; Lambeck et al., 2002a, b; Peltier et al., 2002a, b; Milne et al., 2002; Shennan & Milne, 2003). In fact controversy also surrounds interpretations based on the reported glaciomarine sediments from the Irish Sea basin (see McCarroll et al., 2001; Hiemstra et al., in press). McCabe et al. (in press) suggest that a series of late Glacial relative sea-level changes are recorded by raised marine muds and boulder pavements in Ireland, reflecting the combination of eustatic sea level rise and isostatic adjustments to a dynamic, fluctuating ice-sheet. These interpretations are incompatible with modelled relative sea levels (Lambeck & Purcell, 2001) and work in the Irish Sea basin and neighbouring areas continues to try and account for these important discrepancies.

Whilst the existence of MWP 1a is less controversial, work continues to constrain its precise age (Shennan et al., 2005) and the source of the water responsible for it. Peltier (in press) questions the suggestion by Clark et al. (2002) that the spatial

pattern (or 'fingerprint') of reconstructed sea level rise is indicative of an Antarctic source. He suggests that vertical uncertainties associated with critical sea level data are too large to conclusively rule out a northern Hemisphere source. Baroni & Hall (2004) present data from Terra Nova Bay, Antarctica, indicating deglaciation of the Ross Sea took place comparatively late, with rapid retreat occurring sometime after 8000 cal. year BP. These authors suggest that the presence of large amounts of grounded ice in the Ross Sea around 8000 years ago makes a significant Antarctic contribution to the earlier MWP 1a less likely.

South America has recently received attention from sea level modellers since the distinctive, differential pattern of relative sea-levels change along its eastern coast provides a useful testing ground for their models. Peltier (in press) uses South American sea level data compiled by Rostami et al. (2000), to test the accuracy of rotational effects as modelled by ICE-4G (VM2), concluding that these are 'rather accurate'. Milne et al. (in press) also analyse data from this area, but with the aim of constraining the Holocene highstand, and separating the contribution of eustatic and non-eustatic components. Interestingly, they choose not to use the data of Rostami et al (2000), suggesting it possesses some large observational uncertainties. Instead, they prefer a sub-set of data comprising the most reliable sea level indicators, such as mangrove peats, vermitids and marine molluscs. Milne et al. (in press) suggest their best-fit model indicates a relatively fast eustatic rise in sea level of 7-8 mm/yr in the early Holocene, declining rapidly around 7000 BP.

IV. Recent Sea Level Changes

Separating mass-related eustatic sea level changes from those due to steric effects or the redistribution of water around the globe becomes increasingly difficult after the

early Holocene addition of meltwater diminishes. A classic example of this is the 'attribution problem': direct estimates of 'global' sea level rise based on tide gauge data indicate a rate of 1.5 to 2.0 mm/yr for the last century, but indirect estimates, based on modelling of mass and volume change, predict values much lower than this. The 'fingerprinting' approach can be employed to assist in deciphering the contrasting records provided by satellite altimetry, tide gauges and longer term geological indicators (e.g. Clarke et al., 2003). Nakada & Inoue (in press) analyse seven long duration tide gauge records to estimate sea level change over the last 140 to 200 years. After correction for land level movements, their results reveal a spatial fingerprint consistent with a meltwater contribution from the Greenland ice sheet equivalent to 1 mm/yr, plus an additional 0.5 mm/yr from another source (e.g. thermal expansion and/or melting from Antarctica/mountain glaciers). The authors caution that their results are preliminary given the small number of stations and the fact that the use of data from a tectonically active area may violate an underlying assumption in their analysis.

Miller & Douglas (2004) evaluate the suggestion that tide gauge estimates of sea level rise are perhaps 2-3 times too high due to being located in areas of anomalous ocean warming (Cabanès et al. 2001). They analyse tide gauge data from the Atlantic and Pacific Oceans and compare them with average dynamic heights determined from oceanographic observations. They conclude that migration of the Gulf Stream introduced an error into the analysis of Cabanès et al. (2001) and that the new analysis indicates a discrepancy between tide gauges and steric effects of about 1.5 mm/yr. On this basis they suggest that more mass than previously thought may be being added to the oceans. Some support for this comes from Meehl et al. (2005) who publish the results of two global coupled climate models indicating only 3-5 cm of the observed 15-20 cm rise in 20th century sea levels is due to steric change.

They conclude that either the observations are wrong or melting from glaciers and ice sheets provides a more significant contribution to recent sea level rise than currently thought. Intriguing new data from Antarctica, comprising over 2000 aerial photographs and 100 satellite images, charts changes in 244 marine glaciers from 1940 onwards, and shows 87% are in overall retreat (Cook et al. 2005). Whilst these have no direct influence on sea level (since they are floating) the authors speculate that melting may cause accelerations of the glaciers that feed them, resulting in an increased 'draining' of Antarctica.

Miller & Douglas (2004) do not rule out an alternative explanation for their results involving substantial mass redistribution, but they note that this would require sea level falls during the 20th century over large areas of the central oceans (away from most of the tide gauges). Interestingly, Möner et al. (2004) infer a recent fall in eustatic sea level in the central Indian Ocean based on observations from the Maldives.

A couple of recent papers point toward the possible detection of an acceleration in the long-term rate of sea level rise. Donnelly et al. (2004) construct a high precision relative sea-level curve from Connecticut, USA, covering the last 700 years. The curve is a composite derived from compaction-free AMS radiocarbon dated plant macrofossils and tide gauge data. The authors suggest it indicates a nearly three-fold increase in the rate of rise in the latter half of the 19th century. Gehrels et al. (2005) use a precise foraminiferal transfer function in combination with a detailed chronology comprising AMS radiocarbon dates, pollen data and a suite of short-lived radionuclides, to produce a high-resolution record of sea-level change for the past 1000 years from Nova Scotia. Their data indicate a doubling in the mean rate of relative sea-level rise to around 3.2 mm/yr between about AD 1900 and AD 1920.

This paper highlights the importance of chronology construction, noting that erroneous fluctuations in relative sea-level records can be generated during the calibration of radiocarbon dates.

V. Concluding Remarks

Recent studies examining tsunamis and earthquake hazard provide an excellent example of how the combination of terrestrial and marine data, coupled with the integration of modern observations, palaeoenvironmental reconstructions and computer model simulations, can produce a detailed picture of the mechanisms responsible for an event. Well-dated abrupt events have played an important role in the hunt for the driving mechanisms of climate change, especially where these produce widespread signals that can be extracted from proxy records (e.g. the '8.2 kyr event', Schmidt & LeGrande, 2005). Similarly, abrupt sea level changes, such as those related to meltwater pulses, are beginning to cast light on the relationships among climate, ice sheets and ocean levels.

As geological records are probed at higher levels of resolution, the pattern of variability exhibited alters. In some instances, changes are greater than previously thought, such as the surprising amount of interglacial sea level variability emerging from coral records (Thompson and Goldstein, 2005). In others, an apparently widespread abrupt event may be resolved into separate anomalies or more gradual changes, complicating the relationship between cause and effect (e.g. Rohling & Pälike, 2005). Sea level research faces the particular challenge of linking satellite, tide gauge and geological records together, each of which possess contrasting precisions, resolutions and apparent variability.

Whilst satellite missions such as GRACE will provide new pictures of the global ocean, and feedback into the development of glacio-isostatic models (Peltier, 2004), a long-term perspective can only be provided by geological data. Consequently, attempts to link detailed, high-precision sea-level reconstructions with instrumental records are of critical importance (Gehrels et al., 2001; 2005; Donnelly et al., 2004). These will require continued improvement in the methods used to fix the age and altitude of past relative sea-level. Age control is a particular challenge, and will require better methods of age interpolation (e.g. Gehrels et al., 2005), a fuller understanding of radiocarbon ages derived from different organic components (e.g. van de Plassche et al., 2005), and the adoption of alternative tools, such as luminescence dating (Duller, 2004). It also requires clear communication between disciplines of the errors and limitations associated with data. As Woodroffe & Horton (2005) observe in their recent review of sea level data from the Indo-Pacific, correlations are compromised by the failure of many studies to adequately account for the vertical uncertainties associated with the sea level indicators used. Hence, as Project 495 commences, it is evident that the guidelines developed during IGCP Project 61 are as relevant today as they were 30 years ago.

References

- Atwater, B.F.** 1987: Evidence for great Holocene earthquakes along the outer coast of Washington State. *Science* 236, 942-944.
- Atwater, B.F., and Hemphill-Haley, E.** 1997: *Recurrence intervals for great earthquakes of the past 3,500 years at northeastern Willapa Bay, Washington*. US Geological Survey Professional Paper No. 1576, 1-108.

Baroni, C., and Hall, B.L. 2004: A new Holocene relative sea-level curve for Terra Nova Bay, Victoria Land, Antarctica. *Journal of Quaternary Science* 19, 377-396.

Bondevik, S., Svendsen, J.I., Johnsen, G., Mangerud, J., and Kaland, P.E. 1997a: The Storegga tsunami along the Norwegian coast, its age and runup. *Boreas* 26, 29-53.

Bondevik, S., Svendsen, J.I., and Mangerud, J. 1997b: Tsunami sedimentary facies deposited by the Storegga tsunami in shallow marine basins and coastal lakes, Western Norway. *Sedimentology* 44, 1115-1131.

Bondevik, S., Mangerud, J., Dawson, S., Dawson, A., and Lohne, Ø. 2003: Record-breaking height for 8000-year-old tsunami in the North Atlantic. *Eos* 84 (31), 289-300.

Bondevik, S., Løvholt, F., Harbitz, C., Mangerud, J., Dawson, A., and Svendsen, J.I. 2005: The Storegga Slide tsunami – comparing field observations with numerical simulations. *Marine and Petroleum Geology* 22, 195-208.

Bondevik, S., Mangerud, J., Dawson, S., Dawson, A., and Lohne, Ø. in press: Evidence for three North Sea tsunamis at the Shetland Islands between 8000 and 1500 years ago. *Quaternary Science Reviews*.

Bryn, P., Berg, K., Forsberg, C.F., Solheim, A., and Kvalstad, T.J. 2005: Explaining the Storegga Slide. *Marine and Petroleum Geology* 22, 11-19.

Cabanes, C., Cazenave, A., and Le Provost, C., 2001: Sea level rise during past 40 years determined from satellite and in situ observations. *Science* 294, 840-842.

Clark, J.A., Haidle, P.E. and Cunningham, L.N. 2003: Comparison of satellite altimetry to tide gauge measurement of sea level: predictions of glacio-isostatic adjustment. *Journal of Climate* 15, 3291-3300.

Clark, P.U., and Mix, A.C. 2002: Ice sheets and sea level of the Last Glacial Maximum. *Quaternary Science Reviews* 21, 1-7.

Clark, P.U., Mitrovica, J.X., Milne, G.A., and Tamisiea, M.E. 2002: Sea level fingerprinting as a direct test for the source of global meltwater pulse 1a. *Science* 295, 2438-2441.

Clark, P.U., McCabe, A.M., Mix, A.C., and Weaver, A.J. 2004: Rapid rise of sea level 19,000 years ago and its global implications. *Science* 304, 1141-1144.

Cook, A.J., Fox, A.J., Vaughan, D.G., and Ferrigno, J.G. 2005: Retreating glacier fronts on the Antarctic Peninsula over the past half-century. *Science* 308, 541-544.

Dawson, A.G. 1999: Linking tsunami deposits, submarine landslides and offshore earthquakes. *Quaternary International* 60, 119-126.

Dawson, A.G., and Shi, S. 2000: Tsunami deposits. *Pure and Applied Geophysics* 157, 875-897.

Dawson, A.G., and Smith, D.E. 2000: The sedimentology of a middle Holocene tsunami facies in northern Sutherland, Scotland, UK. *Marine Geology* 170, 69-79.

Dawson, A.G., Foster, I.D.L., Shi, S., Smith, D.E., and Long, D. 1991: The identification of tsunami deposits in coastal sediment sequences. *Science of Tsunami Hazards* 9, 73-82.

Donnelly, J.P., Cleary, P., Newby, P., and Ettinger, R. 2004: Coupling instrumental and geological records of sea-level change: Evidence from southern New England of an increase in the rate of sea-level rise in the late 19th century. *Geophysical Research Letters* 31, L05203.

Duller, G.A.T. 2004: Luminescence dating of Quaternary sediments: recent advances. *Journal of Quaternary Science* 19, 183-192.

Edwards, R.J. 2003: Constructing chronologies of sea-level change from salt-marsh sediments. In: **Buck, C.E.**, and **Millard, A.R.** (eds.), *Tools for Constructing Chronologies: Crossing Disciplinary Boundaries*: Springer Verlag (London) Ltd, 191-213.

Felton, E.A., and **Crook, K.A.W.** 2003: Evaluating the impacts of huge waves on rocky shorelines: as essay review of the book 'Tsunami – The Underrated Hazard'. *Marine Geology* 197, 1-12.

Fryer, G.J., **Watts, P.**, and **Pratson, L.F.** 2004: Source of the great tsunami of 1 April 1946: a landslide in the upper Aleutian forearc. *Marine Geology* 203, 201-218.

Gehrels, W.R., **Belknap, D.F.**, **Black, S.**, and **Newnham, R.M.** 2002: Rapid sea-level rise in the Gulf of Maine, USA, since AD 1800. *The Holocene* 12, 383-389.

Gehrels, W.R., **Kirby, J.R.**, **Prokoph, A.**, **Newnham, R.M.**, **Achterberg, E.P.**, **Evans, H.**, **Black, S.**, and **Scott, D.B.** 2005: Onset of recent rapid sea-level rise in the western Atlantic Ocean. *Quaternary Science Reviews* 24, 2083-2100.

Goff, J., **Chagué-Goff, C.**, and **Nichol, S.** 2001: Palaeotsunami deposits: A New Zealand perspective. *Sedimentary Geology* 143, 1-6.

Goff, J., **McFadgen, B.G.**, and **Chagué-Goff, C.** 2004: Sedimentary differences between the 2002 Easter storm and the 15th-century Okoropunga tsunami, southeastern North Island, New Zealand. *Marine Geology* 204, 235-250.

Gusiakov, V.K. 2005: Tsunami generation potential of different tsunamigenic regions in the Pacific. *Marine Geology* 215, 3-9.

Haflidason, H., Sejrup, H.P., Nygård, A., Mienert, J., Bryn, P., Lien, R., Forsberg, C.F., Berg, K., and Masson, D. 2004: The Storegga Slide: architecture, geometry, and slide development. *Marine Geology* 213, 201-234.

Hamilton, S., and Shennan, I. 2005: Late Holocene great earthquakes and relative sea-level change at Kenai, southern Alaska. *Journal of Quaternary Science* 20, 95-111.

Hamilton, S., and Shennan, I. in press: Late Holocene relative sea-level changes and the earthquake deformation cycle around upper Cook Inlet, Alaska. *Quaternary Science Reviews*.

Hayward, B.W., Cochran, U., Southall, K., Wiggins, E., Grenfell, H.R., Sabaa, A., Shane, P.R., and Gehrels, W.R. 2004: Micropalaeontological evidence for the Holocene earthquake history of the eastern Bay of Plenty, New Zealand, and a new index for determining the land elevation record. *Quaternary Science Reviews* 23, 1651-1667.

Hiemstra, J.F., Evans, D.J.A., Scourse, J.D., McCarroll, D., Furze, M.F.A., and Rhodes, E. in press: New evidence for a grounded Irish Sea glaciation of the Isles of Scilly, UK. *Quaternary Science Reviews*.

Hutton, E.W.H., and Syvitski, J.P.M. 2004: Advances in the numerical modeling of sediment failure during the development of a continental margin. *Marine Geology* 203, 367-380.

Lambeck, K., and Purcell, A. 2001: Sea-level change in the Irish Sea since the last glacial maximum: constraints from isostatic modelling. *Journal of Quaternary Science* 16, 497-505.

Lambeck, K., Yokoyama, Y., and Purcell, A. 2002a: Into and out of the Last Glacial Maximum: sea-level change during oxygen isotope stages 3 and 2. *Quaternary Science Reviews* 21, 343-360.

Lambeck, K., Yokoyama, Y., Purcell, A. and Johnston, P. 2002b: Reply to the comment by W.R. Peltier. *Quaternary Science Reviews* 21, 415-418.

Long, A.J. 2003: The coastal strip: sea-level change, coastal evolution and land-ocean correlation. *Progress in Physical Geography* 27, 423-434.

Løvholt, F., Harbitz, C.B., and Haugen, K.B. 2005: A parametric study of tsunamis generated by submarine slides in the Ormen Lange/Storegga area off western Norway. *Marine and Petroleum Geology* 22, 219-231.

Maramai, A., Graziani, L., and Tinti, S. 2005a: Tsunamis in the Aeolian Islands (Southern Italy): A review. *Marine Geology* 215, 11-21.

Maramai, A., Graziani, L., Alessio, G., Burrato, P., Colini, L., Cucci, L., Nappi, R., Nardi, A., and Vilardo, G. 2005b: Near- and far-field survey report of the 30 December 2002 Stromboli (Southern Italy) tsunami. *Marine Geology* 215, 93-106.

Mastronuzzi, G., and Sansò, P. 2004: Large boulder accumulations by extreme waves along the Adriatic coast of southern Apulia (Italy). *Quaternary International* 120, 173-184.

McCabe, A.M., Clark, P.U., and Clark, J. in press: AMS ^{14}C dating of deglacial events in the Irish Sea Basin and other sectors of the British-Irish ice sheet. *Quaternary Science Reviews*.

McCarroll, D., Knight, J., and Rijdsdijk, K. 2001: Introduction: The glaciation of the Irish Sea basin. *Journal of Quaternary Science* 16, 391-392.

McMurty, G.M., Watts, P., Fryer, G.J., Smith, J.R., and Imamura, F. 2004: Giant landslides, mega-tsunamis, and paleo-sea level in the Hawaiian Islands. *Marine Geology* 203, 219-233.

Meehl, G.A., Washington, W.M., Collins, W.D., Arblaster, J.M., Hu, A., Buja, L.E., Strand, W.G., and Teng, H. 2005: How much more global warming and sea level rise. *Science* 307, 1769-1772.

Miller, L., and Douglas, B.C. 2004: Mass and volume contributions to twentieth-century global sea level rise. *Nature* 428, 406-409.

Milne, G.A., Mitrovica, J.X., and Schrag, D.P. 2002: Estimating past continental ice volume from sea-level data. *Quaternary Science Reviews* 21, 361-376.

Milne, G.A., Long, A.J., and Bassett, S.E. in press: Modelling Holocene relative sea-level observations from the Caribbean and South America. *Quaternary Science Reviews*.

Mörner, N.-A., Tooley, M., and Possnert, G. 2004: New perspectives for the future of the Maldives. *Global and Planetary Change* 40, 177-182.

Nakada, M., and Inoue, H. in press: Rates and causes of recent global sea-level rise inferred from long tide gauge records. *Quaternary Science Reviews*.

Nanayama, F., Shigeno, K., Satake, K., Shimokawa, K., Koitabashi, S., Miyasaka, S., and Ishii, M. 2000: Sedimentary differences between the 1993 Hokkaido-nanasei-oki tsunami and the 1959 Miyakojima typhoon at Taisei, southwestern Hokkaido, northern Japan. *Sedimentary Geology* 135, 255-264.

Nott, J. 2003: Waves, coastal boulder deposits and the importance of the pre-transport setting. *Earth and Planetary Science Letters* 210, 269-276.

Okal, E.A. 2005: A re-evaluation of the great Aleutian and Chilean earthquakes of 1906 August 17. *Geophysical Journal International* 161, 268-282.

Okal, E.A., and Synolakis, C.E. 2004: Source discriminants for near-field tsunamis. *Geophysical Journal International* 158, 899-912.

Pelinovsky, E., and Tinti, S. 2005: Editorial for 'Tsunamis in the Pacific, in the Atlantic and in Europe: relevant cases and hazard evaluation'. *Marine Geology* 215, 1-2.

Peltier, W.R. 2002a: Comments on the paper of Yokoyama et al. (2000), entitled "Timing of the Last Glacial Maximum from observed sea level minima". *Quaternary Science Reviews* 21, 409-414.

Peltier, W.R. 2002b: On eustatic sea level history: Last Glacial Maximum to Holocene. *Quaternary Science Reviews* 21, 377-396.

Peltier, W.R. 2004: Global glacial isostasy and the surface of the ice age Earth: the ICE-5G(VM2) model and GRACE. *Annual Reviews of Earth and Planetary Science* 32, 111-149.

Peltier, W.R. in press: On the hemispheric origins of meltwater pulse 1a. *Quaternary Science Reviews*.

Rohling, E.J., and Pälike, H. 2005: Centennial-scale climate cooling with a sudden cold event around 8,200 years ago. *Nature* 434, 975-979.

Rostami, K., Peltier, W.R., and Mangini, A. 2000: Quaternary marine terraces, sea level changes and uplift history of Patagonia, Argentina: comparisons with predictions of the ICE-4G(VM2) model of the global process of glacial isostatic adjustment. *Quaternary Science Reviews* 19, 1495-1525.

Sawai, Y., Horton, B.P., and Nagumo, T. 2004a: The development of a diatom-based transfer function along the Pacific coast of eastern Hokkaido, northern Japan - an aid in paleoseismic studies of the Kuril subduction zone. *Quaternary Science Reviews* 23, 2467-2483.

Sawai, Y., Satake, K., Kamataki, T., Nasu, H., Shishikura, M., Atwater, B.F., Horton, B.P., Kelsey, H.M., Nagumo, T., and Yamaguchi, M. 2004b: Transient uplift after a 17th-century earthquake along the Kuril subduction zone. *Science* 306, 1918-1920.

Scheffers, A. 2004: Tsunami imprints on the Leeward Netherlands Antilles (Aruba, Curaçao, Bonaire) and their relation to other coastal problems. *Quaternary International* 120, 163-172.

Schmidt, G.A., and LeGrande, A.N. 2005: The Goldilocks abrupt climate change event. *Quaternary Science Reviews* 24, 1109-1110.

Shennan, I., and Milne, G. 2003: Sea-level observations around the Last Glacial Maximum from the Bonaparte Gulf, NW Australia. *Quaternary Science Reviews* 22, 1543-1547.

Shennan, I., Scott, D., Rutherford, M., and Zong, Y. 1999: Microfossil analysis of sediments representing the 1964 earthquake, exposed at Girdwood Flats, Alaska, USA. *Quaternary International* 60, 55-73.

Shennan, I., Hamilton, S., Hillier, C., and Woodroffe, S. 2005: A 16 000-year record of near-field relative sea-level changes, northwest Scotland, United Kingdom. *Quaternary International* 133-134, 95-106.

Smith, D.E., Shi, S., Cullingford, R.A., Dawson, A.G., Dawson, S., Firth, C.R., Foster, I.D.L., Fretwell, P.T., Haggart, B.A., Holloway, L.K., and Long, D. 2004:

The Holocene Storegga Slide tsunami in the United Kingdom. *Quaternary Science Reviews* 23, 2291-2321.

Solheim, A., Berg, K., Forsberg, C.F., and Bryn, P. 2005: The Storegga Slide complex: repetitive large scale sliding with similar cause and development. *Marine and Petroleum Geology* 22, 97-107.

Stein, S., and Okal, E.A. 2005: Speed and size of the Sumatra earthquake. *Nature* 434, 581-582.

Syvitski, J.P.M., Vörösmarty, C.J., Kettner, A.J., and Green, P. 2005: Impact of humans on the flux of terrestrial sediment to the global coastal ocean. *Science* 308, 376-380.

Tappin, D.R. 2004: Preface for 'Submarine-slump-generated Tsunamis'. *Marine Geology* 203, 199-200.

Thompson, W.G., and Goldstein, S.L. 2005: Open-system coral ages reveal persistent suborbital sea-level cycles. *Science* 308, 401-404.

Tooley, M.J., and Smith, D.E. 2005: Relative sea-level change and evidence for the Holocene Storegga Slide tsunami from a high-energy coastal environment: Cocklemill Burn, Fife, Scotland, UK. *Quaternary International* 133-134, 107-119.

van de Plassche, O., Bohncke, S.J.P., Makaske, B., and van der Plicht, J. 2005: Water-level changes in the Flevo area, central Netherlands (5300-1500 BC): implications for relative mean sea-level rise in the Western Netherlands. *Quaternary International* 133-134, 77-93.

Williams, D.M., and Hall, A.M. 2004: Cliff-top megaclast deposits of Ireland, a record of extreme waves in the North Atlantic – storms or tsunamis? *Marine Geology* 206, 101-117.

Williams, H.F.L., Hutchinson, A., and Nelson, A.R. 2005: Multiple sources for late-Holocene tsunamis at Discovery Bay, Washington State, USA. *The Holocene* 15, 60-73.

Woodroffe, S.A., and Horton, B.P. 2005: Holocene sea-level changes in the Indo-Pacific. *Journal of Asian Earth Sciences* 25, 29-43.

Yokoyama, Y., Lambeck, K., De Deckker, P., Johnston, P., and Fifield, L.K. 2000: Timing of the Last Glacial Maximum from observed sea-level minima. *Nature* 406, 713-716.

Yokoyama, Y., Lambeck, K., De Deckker, P., Johnston, P., and Fifield, L.K. 2001: Sea-level at the Last Glacial Maximum: evidence from northwestern Australia to constrain ice volumes for Oxygen Isotope Stage 2. *Palaeogeography, Palaeoclimatology, Palaeoecology* 165, 281-297.

Yokoyama, Y., De Deckker, P., and Lambeck, K. 2003: Reply to “Sea-level observations around the Last Glacial Maximum from the Bonaparte Gulf, NW Australia” by I. Shennan and G. Milne. *Quaternary Science Reviews* 22, 1549-1550.

Zong, Y., Shennan, I., Combellick, R.A., Hamilton, S.L., and Rutherford, M.M. 2003: Microfossil evidence for land movements associated with the AD 1964 Alaska earthquake. *The Holocene* 13, 7-20.