Helpful and Hindering Events in Internet-Delivered Cognitive Behavioural Treatment for Generalized Anxiety

John Burke

E-Mental Health Research Group, School of Psychology, Trinity College, Dublin and Research and Innovation, SilverCloud Health, Dublin, Ireland

Derek Richards

E-Mental Health Research Group, School of Psychology, Trinity College, Dublin and Research and Innovation, SilverCloud Health, Dublin, Ireland

Ladislav Timulak

E-Mental Health Research Group School of Psychology, Trinity College, Dublin

Background: Anxiety disorders are a highly prevalent cause of impairment globally with generalized anxiety disorder (GAD) sharing many features with other anxiety disorders. **Aims:** The present study investigated the helpful and hindering events and impacts for individuals with generalized anxiety who engaged with a supported 6-week online intervention based on cognitive behavioural therapy (iCBT). **Method:** Participants ($n = 36$) completed the Helpful and Hindering Aspects of Therapy (HAT) for each session. A descriptive-interpretative framework was used to analyse the data. **Results:** Helpful events were identified by participants as CBT techniques including psychoeducation, monitoring, cognitive restructuring and relaxation, and found supporter interaction, mindfulness and reading personal stories helpful. The associated impacts were identified as support and validation; behavioural change/applying coping strategies; clarification, awareness, and insight; reassurance/relief; and self-efficacy/empowerment. Hindering events were identified as treatment content/form; and amount of work/technical issues, which led to impacts such as frustration/irritation; increased anxiety; and isolation. **Conclusion:** The implications of the results, potential future directions of research and limitations of the study are discussed.

Keywords: significant events in therapy, helpful and hindering events, internet-delivered interventions, generalized anxiety, descriptive-interpretative framework

Introduction

Generalized anxiety disorder (GAD) is common, persistent and characterized by excessive worry about several events, including minor day-to-day matters in the past, present or future. Often, GAD presents comorbidly with mood disorders (Tyrer and Baldwin, 2006) and can cause significant distress and impairment in daily life for individuals (Judd *et al.*, 1998). Lifetime prevalence rates are estimated at 6.2% (Somers *et al.*, 2006). It has been documented as one of the most difficult anxiety disorders to treat, yielding lower treatment response relative to other anxiety disorders (Jones, 2014).

Treatment of GAD

Reviews of GAD outcome research have indicated that pharmacotherapies and psychological treatments are effective (Tyrer and Baldwin, 2006). A comprehensive meta-analysis of psychological treatments for GAD concluding that CBT is the most empirically supported treatment available (Cuijpers *et al.*, 2014). However, many cases of GAD go undiagnosed, especially in university settings where services are often overburdened with long waiting lists and lack of resources (Mowbray *et al.*, 2006).

Internet-based psychological interventions for GAD

Internet-delivered interventions are accessible and offer greater access for those unlikely to seek traditional approaches to care due to stigmatization or preferences for self-reliance (Thomas *et al.*, 2015). Such interventions can complement university services that are not resourced to offer more traditional costly and time-intensive services (Andersson and Cuijpers, 2009). Richards and colleagues (2015) meta-analysis evaluating internet-delivered interventions for GAD found large effect sizes comparable to those of face-to-face therapy for GAD across several outcome measures. Despite these results, research examining how internet-delivered interventions for GAD can integrate into routine practice is unclear (Batelaan *et al.*, 2012). Furthermore, research has identified that there are largely differing responses to internet-delivered interventions across demographics, for certain personalities (Bendelin *et al.*, 2011), and those with differing attitudes (Rochlen *et al.*, 2004). Understanding the client experiences of iCBT is an integral aspect in supporting iCBT adoption into routine practice and evaluating significant events of therapy provides a means of doing this.

Significant events research

Significant events research (Elliott *et al.*, 1985) represents the study of the important moments in therapy that elicit change for the client, either of a helpful or unhelpful (hindering) nature (Timulak, 2010). Evaluating the client's experience of therapy is often used to ascertain the processes whereby therapeutic methods are effective and inform service provision. Most of the research has been conducted through qualitative methods, mainly semi-structured interviews (Elliott, 2008).

Significant events research in internet-delivered treatments is limited and the therapeutic processes involved are relatively unknown. There has, however, been some qualitative research examining the psychological experience of users in iCBT treatments for depression (Bendelin

et al., 2011; Gega *et al.*, 2013). The Helpful/Hindering Aspects of Therapy (HAT) form is one questionnaire that gathers written data revealing client-identified significantly helpful and hindering events (Elliott *et al.*, 2001). It has been used in previous studies that have assessed client-identified helpful and hindering events and their impacts (Gega *et al.*, 2013; Richards and Timulak, 2012). This provides the opportunity to gain a qualitative insight into client experiences of iCBT.

The current study intended to gain an understanding of the helpful and hindering events and associated impacts of iCBT for GAD symptoms, specifically within a student population. The present study intended to contribute to significant events research in iCBT and provide guidance for developing future iCBT programmes for GAD.

Methods

Design

The study employed a qualitative design using a descriptive and interpretative framework (Elliott and Timulak, 2005). The qualitative data were obtained from a larger randomized controlled trial (RCT) evaluating iCBT for GAD in a university population (Richards *et al.*, 2014, 2016).

Participants

Participants were recruited from an Irish University using an email advertisement for an internet-delivered course for anxiety called '*Calming Anxiety*'. Further information on the trial design and its main outcomes is available (Richards *et al.*, 2014, 2016). The primary outcome measure was the Generalized Anxiety Disorder 7-item scale (GAD-7; Spitzer *et al.*, 2006). Those obtaining a score of 10 or above on GAD-7, equating to moderate to severe levels of anxiety according to normative values were included in the overall analysis (mean = 12.8, $SD = 2.68$). These scores correlate to a formal diagnosis of GAD as outlined in DSM-IV (Spitzer *et al.*, 2006), but formal diagnostic assessment was not carried out on the present cohort.

Sample size

The original RCT recruited 137 students from an Irish-based university to engage with the '*Calming Anxiety*' iCBT intervention. Participants were randomized into the immediate treatment group ($n = 70$) or a waiting list control group ($n = 67$). This participant group consisted of 69% female participants ($n = 25$) and 31% male ($n = 11$). Age ranged from 18 to 45 years (mean = 24.69, $SD = 7.37$). A total number of 78 from a possible 216 HAT responses (36%) were obtained from the 36 participants. Figure 1 shows the breakdown of HAT responses per session.

Measure

The HAT form (Llewelyn, 1988) was utilized to assess the most helpful and hindering events in therapy. Participants were asked to describe, in their own words, what they felt were the most significant events in each module for them. This event is anything identified by the participant

Figure 1. Column chart depicting number of HAT forms complete for each session

that they, the course or supporter said/did. The form also asks for a description of what made the event helpful or hindering for them, to elucidate their associated impacts.

Procedure

Participants were randomly allocated into one of two conditions – immediate access to the internet-delivered treatment intervention and delayed access to treatment using a waiting list control. The treatment was an iCBT program entitled ‘*Calming Anxiety*’ composed of six modules of CBT content for GAD consisting of cognitive, emotional and behaviour components (see Richards *et al.*, 2014 for description of each module). In addition, participants received encouragement and specific feedback from their assigned supporter. Each supporter was a graduate psychology student qualified to Master’s level or higher. Participants completed HAT measures from their second login to the platform where they were asked to reflect on the previous session and complete the questions.

Data analysis

Descriptive and interpretative (DI) qualitative analysis (Elliott and Timulak, 2005) was used to analyse the data collected to help understand the clients’ perspectives on helpful and hindering events and impacts of the intervention. The first author received an Excel file containing the anonymized HAT data collected from participants of the main RCT. The process of analysis followed some distinct steps:

- (1) The first author studied the data thoroughly, noting certain themes appearing regularly across participant responses. This process was repeated several times until the first author had a sufficient grasp of themes within the data.
- (2) The data were divided into four domains: helpful events, helpful impacts, hindering events, hindering impacts. The clients’ descriptions were considered about the event they described as well as the associated impact of the event. Richards and Timulak (2012) recommend this structure to organize the data, to determine what events are helpful/hindering as well as the impact of these events.

Table 1. Instances for helpful and hindering events and helpful impacts for each category

Helpful events	Pts	Helpful impacts	Pts
Psychoeducation	16/36	Support and Validation	19/36
Supporter Interaction	15/36	Applying Coping strategies/Behavioural Change	19/36
Monitoring	10/36	Clarification, Awareness and Insight	17/36
Relaxation	6/36	Reassurance/Relief	12/36
Mindfulness	6/36	Self-Efficacy/Empowerment	7/36
Cognitive Restructuring	4/36		
Reading Personal Stories	3/36		
Other Skills	3/36		
Hindering events	Pts	Hindering impacts	Pts
Treatment Content/Form	11/36	Frustration/Irritation	7/36
Amount of Work/Technical Issues	7/36	Increased Anxiety	6/36
		Isolation	4/36

Pts, number of participants whose responses exhibited data associated with specified event or impact, e.g. 16/36 means 16 out of 36.

- (3) Meaning units were identified, i.e. pieces of text that when placed outside of their context held a specific meaning.
- (4) Through analysing these meaning units, reoccurring themes in the data were noted and general categories emerged. Each category was defined to reflect the content of data, ensuring each meaning unit was sufficiently represented by the category's definition to which it was assigned. This process was repeated several times to ensure accuracy.
- (5) Further review of each category, category definitions along with the assigned meaning units was carried out by the second author, who has experience in DI analysis, to ensure accurate labelling and categorization of an 'event' or 'impact', and revisions made. The third author, with significant experience in conducting qualitative research, then reviewed the steps of the analysis and the outline of the findings. The changes suggested by the second and third author were then discussed and incorporated to the final analysis.

Results

DI analysis revealed a total of eight helpful events (*Psychoeducation; Supporter Interaction; Monitoring; Relaxation; Mindfulness; Cognitive Restructuring; Reading Personal Stories; Other Skills*) and five associated impacts (*Support and Validation; Behavioural Change/Applying Coping Strategies; Clarification, Awareness and Insight; Reassurance/Relief; Self-Efficacy/Empowerment*). Table 1 details the number of participants who experienced each helpful and hindering event and impact. Two hindering events were identified (*Treatment Content/Form; Amount of Work/Technical Issues*), with three associated impacts (*Frustration/Irritation; Increased Anxiety; Isolation*). The majority of the data was collected after sessions 1 and 2 ($n = 51$). Responses from sessions 3 to 6 were lower ($n = 27$).

Therefore, the events and their associated impacts described pertain mainly to the initial sessions of the iCBT programme.

Each category was defined for helpful events and impacts (Table 2) and hindering events and impacts (Table 3). Each helpful and hindering event and impact is described in more detail below.

Helpful events

Supporter interaction ($n = 15$) encompasses any form of feedback, advice, encouragement or acknowledgement by the supporter in their personalized responses that were recognized as a helpful aspect of the course, e.g. *My supporter did a really good summary of how I was feeling, it was good to see it written down from an outside perspective.* *Mindfulness* ($n = 6$) techniques were another helpful event, e.g. *The mindfulness tools were helpful in terms of how to relax.* The *Reading Personal Stories* ($n = 3$) category relates to a specific component of the intervention where individuals describe their experience of anxiety making reference to common symptoms, thoughts, behaviours and feelings, e.g. *I liked reading and relating to the personal stories.* *Other Skills* ($n = 3$) were helpful events of learning unspecified skills to manage anxiety, e.g. *Learning about the exercises for reducing my anxiety.*

Participants reported learning four CBT techniques through their engagement with the programme. *Psychoeducation* ($n = 19$) involves learning a new piece of information relating to the prevalence of anxiety in the general population, what the potential causes of anxiety might be, how anxiety is perpetuated or how to reduce and manage anxiety, e.g. *Reading about how common anxiety is.* *Monitoring* ($n = 10$) refers to attending to one's actions, emotions, thoughts or physiological reactions, e.g. *I am trying to track thoughts or events that elicit anxiety, and then (sic) trying to find the way to fix or avoid them.* *Relaxation* ($n = 6$) involves learning methods for relaxation to reduce physiological tension, e.g. *Relaxation tools available online – very useful for calming down at the end of a day.* *Cognitive Restructuring* ($n = 4$) involves learning to refute cognitive distortions and replace them with healthier cognitions, e.g. *Identifying hot thoughts will be worthwhile when I can master it, in the moment.*

Helpful impacts

Five helpful impacts were identified. Nineteen participants described how they felt *Supported* or *Validated* through their engagement with the course. This feeling could occur because of supporter interaction, or clients' resonance with the course material, e.g. *I feel like having someone there look over my progress generally has encouraged me to keep going with the programme.* Nineteen participants reported that they *Applied a New Skill* or experienced *Behavioural Change* assisting them in managing their anxiety daily, e.g. *I took the suggestion of the mindfulness exercise and am working on integrating this type of approach into everyday life.* Seventeen participants reported developing an *Awareness* of the influences of anxiety, an *Insight* into recognizing their own anxiety or *Clarification* of aspects of anxiety that might previously have been ambiguous to them, e.g. *It made me focus on my own situation and think about everything I had just read and how it related to me.*

Twelve participants reported feeling *Reassured*, or gaining a sense of *Relief*, e.g. *It was reassuring to see some events/feelings outlined in the cycle of anxiety that I thought were exclusive to me.* Because of their engagement with the course, seven participants reported

Table 2. Definition for each category of helpful event and impact

Helpful event	Definition
Psychoeducation	Education on: how common anxiety is in the general population; the causal factors in anxiety (i.e. Thoughts, Feelings Behaviour (TFB) cycle; the mechanics whereby anxiety is perpetuated (i.e. avoidance; poor sleep hygiene) and the ways that anxiety can be reduced; learned through material in each session
Supporter Interaction	Receiving advice, encouragement, acknowledgement or general feedback from the supporter. Sharing/disclosing information with the supporter
Monitoring	Systematically attending to and evaluating one's current actions, thoughts, emotions, or physiological reactions
Relaxation	Learning new habits based on relaxation principles that encourage reduction of physiological tension or arousal
Mindfulness	Using mindfulness techniques stated in the course
Cognitive Restructuring	Learning to refute cognitive distortions, and to replace these cognitions with more helpful and healthier cognitions
Reading Personal Stories	Reading and relating to the personal stories described within the course
Other Skills	The learning of non-specific, unspecified skills that do not fit into any other category, that are helpful to manage anxiety
Helpful impact	Definition
Support and Validation	Feeling their thoughts, feelings, fears and anxieties are accepted, understood and validated through supporter interaction, or course material engagement (i.e. by means of relating to personal stories) Feeling that they are supported through their interaction with the course rather than feeling alone as they might have had previously
Applying Coping Strategies/Behavioural change	The application of new skills learned through participation in the course (i.e. CBT techniques; mindfulness) that they have continued to practise outside of the course requirements that have been helpful in managing or controlling their anxiety
Clarification, Awareness and Insight	Developing awareness of how certain life events and daily life experiences can influence anxiety; an insight into the recognition of their own anxiety, realizing the impact of their anxiety on their life and what they can do to control the anxiety they experience Clarifying certain aspects of the contributory and perpetuating factors of anxiety that participants may have previously held misconceptions about
Reassurance/Relief	Gaining reassurance from supporter or course material. Feeling relief through engagement with the course material and/or supporter feedback
Self-Efficacy/Empowerment	Feeling 'in control' of their anxiety, feeling that they have the ability to manage their own issues through engagement with the programme. Identifying their own role in tackling their anxiety and moving towards taking control of it themselves

Table 3. Definition for each category of hindering event and impact

Event	Definition
Treatment content/form	Comments made about the material encountered within the course (i.e. personal stories) or relating to the computer-based, short-term form of treatment. For example, comments made about the ineffectiveness or irrelevance of material
Technical issues	Any issue related to course structure, the layout and/or administration of the course, any technical difficulties encountered, or issues with time commitment required for engagement with the course
Impact	Definition
Frustration/irritation	Events experienced as a result of engagement with the course causing frustration and/or irritation for participants
Increased anxiety	Clients experiencing an increase in their anxiety as a result of engaging with the course
Isolation	Events making participants feel as though they are unsupported throughout their participation in the course, and feeling isolated as though they are experiencing symptoms that are not commonly experienced by others

instances of feeling ‘in control’ of their anxiety through *Self-Efficacy/Empowerment* and having the ability to manage the issues they are experiencing because of their anxiety, e.g. *I feel a little less out of control.*

Hindering events

Eleven participants in this category alluded to unhelpful course material (*Content/Form*) such as the personal stories not resonating with them, or to the online, short-term nature of the programme as unsuitable for them, e.g. *I just find it hard to start certain activities because the examples available, though still helpful, are not really like my problems or anxiety.* Seven participants noted issues with the course structure, administration or the time commitment required to engage (*Amount of Work/Technical Issues*), e.g. *Some of the content was a little bit long.*

Hindering impacts

Three hindering impacts were identified. This first category captured instances of *Frustration and Irritation* seven participants felt while completing the course, e.g. *I found much of the advice impossible to achieve with a student lifestyle, hours at college and a house full of students.* Six participants reported feeling an increased level of *Anxiety* because of their participation, for example due to monitoring and recording their anxiety or technical administration, e.g. *Anxiety triggered whenever I saw I had a [programme name] email.* Four participants reported feeling *Isolated* through their engagement with the programme, e.g. *The examples available, are not really like my problems or anxiety. It kind of makes me feel a bit more alone as no one seems to feel the same things I do.*

Discussion

The present study set out to gain an understanding of how students enrolled in a university course experience an iCBT intervention for GAD symptoms. Specifically, it aimed to identify the significantly helpful and hindering events of the course and their potential impacts. Participants found certain aspects of the course particularly helpful. CBT techniques such as *Psychoeducation*, *Monitoring*, *Relaxation* and *Cognitive Restructuring* were helpful for some participants. *Mindfulness* and *Reading Personal Stories* and *Other Skills* not specifically named were helpful, while the *Supporter Interaction* also played a significant role for some participants.

Several participants began to *Apply Coping Strategies* or experience *Behavioural Change* in their daily lives because of their engagement with the programme. Feeling *Supported and Validated* cultivated a feeling of *Self-Efficacy/Empowerment* for some participants. Developing *Clarification* on, and *Awareness* or *Insight* into how their own life is affected by anxiety, were helpful impacts. The internet-delivered course evoked a feeling of *Reassurance/Relief* among participants, who previously felt their anxiety was all consuming.

Some hindering aspects were also identified. The *Treatment Content/Form* left some people feeling *Frustrated* or *Irritated* by the *Amount of Work* required and *Technical Issues* relating to layout or administration. Some participants found the online platform too like their working environment. Those who did not feel supported and validated instead felt *Isolated* and some were left with an increased level of *Anxiety*.

The presence of *Applying Coping Strategies/Behavioural Change* as a prominent client-identified impact in the data indicates that ‘*Calming Anxiety*’ was successful in equipping some participants with strategies to manage their anxiety. This is an important facet of treatment, as these individuals may now have some resources to manage their anxiety and these may offset the need for more intensive treatment.

Some participants reported that the relational element developed through personalized feedback allowed their anxiety to be validated; evidence suggests that validation is important for improvement when treating any psychological disorder (Linehan, 1997). The responses of some participants suggest that the supporter was an integral component of the intervention for their outcomes. Supported internet-delivered interventions generally yield better outcomes (Andersson and Cuijpers, 2009; Richards and Richardson, 2012). However, over half of the participants did not cite the supporter in their responses as either a helpful or hindering aspect. The issues within research concerning the involvement of a supporter in iCBT were mentioned previously (Titov *et al.*, 2013) and the present results serve to pose more questions rather than provide answers in this debate.

It could be the case that individuals who did not highlight the supporter interaction in their responses were assigned a supporter less skilled in relating to participants through their feedback. Alternatively, it could be suggested that two pathways to treatment success exist – one relying on skills and strategies and the other on the interpersonal aspects of the programme and the supporter (Richards *et al.*, 2013; Richards *et al.*, 2017).

A sense of empowerment was experienced by seven participants and such an outcome is a welcome therapeutic benefit from the course. Some individuals, however, may have had a greater sense of self-efficacy prior to engagement, but all may have valued gaining knowledge on coping strategies that they could implement themselves and enhance self-management

(Dwyer *et al.*, 2011). This may be facilitative of students given the preference to foster a sense of autonomy (Gulliver *et al.*, 2012).

Awareness and insight are commonly reported categories in significant events research (Timulak, 2007). Gaining an insight into these detrimental effects of anxiety on one's behavioural and cognitive functioning, and how these effects could be managed revealed itself as a powerful impact within the present data. However, for both internet-delivered and traditional forms of therapy it is argued that the therapist (or supporter in this case) is required to facilitate the client in reaching the deepest potential of the significant insight resulting in the most significant change (Elliott, 1985). The issue of sufficient therapist presence in these moments is an area ripe for future research.

It may be the case that these significant insights are not navigated to their full potential. This is not necessarily a criticism that can only be levelled at iCBT, it can also be the case in other delivery mediums. To add to this dynamic, it may be the case that these differences are unrelated to the administration of the intervention. Instead, each client's personal and interpersonal characteristics colour their experiences of the supporter and report different types of events in keeping with these differences.

The delivery and presentation of the material in earlier iCBT interventions was identified as a hindering event in a significant number of client experience evaluation studies (Ritterband *et al.*, 2003). The present results indicate less prominent hindrances of engaging with this intervention, portraying the advancements in technology and expertise in internet-delivered interventions markedly improving the presentation and delivery. However, hindrances were identified relating to the content, general administration and technical features. The consideration of participants' hindering experiences and their impact can only advance our knowledge of how to further develop interventions to promote and optimize engagement.

Implications for clinical practice and future research

The present findings contribute to the growing list of studies evaluating client experiences of internet-delivered interventions. To the researchers' knowledge, it is the first qualitative study of participant-identified significant events, specifically iCBT for GAD symptoms.

Whether the involvement of a supporter is related to self-efficacy of an individual, and what characteristics are salient for participants placing a greater reliance on the supporter's role are two important avenues of research. Further research is welcomed on the function of the supporter: can support facilitate participants to gain knowledge, insight and develop strategies, or does it result in an over-reliance, leading to little or no learning of coping strategies? Given the importance of developing insight for therapeutic change, future research might be useful to understand the level of insight iCBT and internet-delivered interventions in general can bring and comprehend the impact of differing levels of support in this process. Furthermore, determining whether the supporter is capable of sufficiently navigating developments in insight for participants would refine the role of the supporter in internet-delivered interventions. There is also a case for evaluating supporter feedback to determine whether outcomes depend on the supporter they were assigned to determine if the intervention impacts client experiences above personal and interpersonal differences.

Delineating these complex interactions with internet-delivered interventions is important for continuous advancement in internet-delivered interventions research and bringing parity between the knowledge and research on internet-delivered interventions and traditional

methods of therapy. This challenge will require innovative methods to capture the experience of users and therapists.

Limitations

It is generally accepted that qualitative research can hold significance with lower numbers of participants and responses. However, the high rates of attrition between the total number of participants and the number of participants who completed HAT forms could be a potential limitation. Most events and impacts described were observed within the first two iCBT sessions. The participants had not yet navigated through modules describing graded exposure, challenging negative thoughts and summary modules. Therefore, there is the potential that some events occurred in the latter sessions that were not captured in the present analysis. In addition, participants may have described the experiences of the first two sessions differently having completed the entire programme. For example, technical issues and the associated frustration may not have been as prominent as participants increased their computer competence throughout their use of the programme. However, the present data do not provide the opportunity to examine the participant experiences of latter modules. Furthermore, it was not possible to capture the experiences of those who dropped out of the intervention, which would be useful to address potential shortcomings of iCBT in meeting user needs and expectations.

The data was gathered from a university student sample. This was a heterogeneous sample capturing a range of demographics. This sample is potentially more computer-literate and has a higher level of education in comparison with the general population.

The results of the primary study outcomes outlined how participants presented with GAD symptoms as opposed to GAD disorder (Richards *et al.*, 2016). This discrepancy may have been due to the use of the GAD-7 alone to establish GAD in the sample. This can be considered a limitation, as it cannot be assumed that a moderate/severe score on the GAD-7 equates directly to a formal GAD diagnosis. Despite this, client experiences were being evaluated rather than outcomes in the present paper, and GAD-7 scores represent a sufficient measure of participant's self-reported anxiety.

The authors used auditing to ensure the quality and accuracy of analysis but as is the case with all qualitative research, there is a subjective component to the analysis that may have influenced the interpretation of the data.

Conclusion

Analysing client experiences has demonstrated that iCBT (*Calming Anxiety*) contains helpful elements that can have relevant impacts. Many participants reported a greater insight into their anxiety and learned skills they could apply in their daily lives. Through the application of these skills, they gained a sense of empowerment and self-efficacy, motivating them to manage their anxiety. The programme left some participants feeling supported and validated, much of which was due to supporter interaction throughout the course. As previously noted, some aspects of the treatment content and form caused an increase in anxiety. The suitability, or unsuitability of the course material for a small percentage of participants presents an issue for developers, clinicians and researchers alike that needs to be addressed in the development of future iCBT interventions to reduce the instance of hindering, negative impacts because of engaging with the programme. Beginning to understand the more complex processes involved

in internet-delivered interventions is an important avenue for research and should be prioritized if internet-delivered interventions are to be widely implemented into routine care, namely in relation to therapeutic insight and the role of the supporter for participants.

The present study makes a much-needed contribution to the field of research evaluating client experiences within internet-delivered interventions. More specifically, it provides a qualitative perspective on the use of iCBT for GAD symptoms. Future research should continue to support the utilization of iCBT for GAD as a low-intensity intervention within routine practice of university psychological services to manage the increasing issue of untreated anxiety in the student population.

Acknowledgements

The authors would like to acknowledge the authors of the primary study for the provision of secondary data evaluated in this study (Richards *et al.*, 2014, 2016).

Thanks to all of the people who participated in the study. Thanks also to the student counselling service for hosting the study.

Ethical statements: The authors have abided by the Ethical Principles of Psychologists and Code of Conduct as set out by the APA. Ethical approval for the original study was obtained from the School of Psychology Research Ethics Committee, Trinity College Dublin, Ireland (22 November 2013). The authors adhere to both research ethics and Trinity College Dublin policy on good research practice. Approval for examination of the secondary data was provided by the same committee on 2 December 2015.

Conflicts of interest: Mr John Burke, Dr Derek Richards and Dr Ladislav Timulak declare no conflicts of interest.

Financial support: This trial was jointly funded from monies and resources provided by SilverCloud Health Ltd and Trinity College Dublin student counselling service.

References

- Andersson, G. and Cuijpers, P. (2009). Internet-based and other computerized psychological treatments for adult depression: a meta-analysis. *Cognitive Behaviour Therapy*, 38, 196–205. doi: [10.1080/16506070903318960](https://doi.org/10.1080/16506070903318960)
- Batelaan, N. M., Smit, J. H., Cuijpers, P., van Marwijk, H. W., Terluin, B. and van Balkom, A. J. (2012). Prevention of anxiety disorders in primary care: a feasibility study. *BMC Psychiatry*, 12, 1. doi: [10.1186/1471-244X-12-206](https://doi.org/10.1186/1471-244X-12-206)
- Bendelin, N., Hesser, H., Dahl, J., Carlbring, P., Nelson, K. Z. and Andersson, G. (2011). Experiences of guided internet-based cognitive-behavioural treatment for depression: a qualitative study. *BMC Psychiatry*, 11, 1. doi: [10.1186/1471-244X-11-107](https://doi.org/10.1186/1471-244X-11-107)
- Cuijpers, P., Sijbrandij, M., Koole, S., Huibers, M., Berking, M. and Andersson, G. (2014). Psychological treatment of generalized anxiety disorder: a meta-analysis. *Clinical Psychology Review*, 34, 130–140. doi: [10.1016/j.cpr.2014.01.002](https://doi.org/10.1016/j.cpr.2014.01.002)
- Dwyer, L. A., Hornsey, M. J., Smith, L. G., Oei, T. P. and Dingle, G. A. (2011). Participant autonomy in cognitive behavioral group therapy: an integration of self-determination and cognitive behavioral theories. *Journal of Social and Clinical Psychology*, 30, 24. doi: [10.1521/jscp.2011.30.1.24](https://doi.org/10.1521/jscp.2011.30.1.24)

- Elliott, R.** (1985). Helpful and nonhelpful events in brief counseling interviews: an empirical taxonomy. *Journal of Counseling Psychology*, 32, 307–322. doi: [10.1037/0022-0167.32.3.307](https://doi.org/10.1037/0022-0167.32.3.307)
- Elliott, R.** (2008). Research on client experiences of therapy: introduction to the special section. *Psychotherapy Research*, 18, 239–242. doi: [10.1080/10503300802074513](https://doi.org/10.1080/10503300802074513)
- Elliott, R., James, E., Reimschuessel, C., Cislo, D. and Sack, N.** (1985). Significant events and the analysis of immediate therapeutic impacts. *Psychotherapy: Theory, Research, Practice, Training*, 22, 620. doi: [10.1037/h0085548](https://doi.org/10.1037/h0085548)
- Elliott, R., Slatick, E. and Urman, M.** (2001). Qualitative change process research on psychotherapy: alternative strategies. *Psychological Test and Assessment Modeling*, 43, 69. doi: [10.1002/9781119973249.ch6](https://doi.org/10.1002/9781119973249.ch6)
- Elliott, R. and Timulak, L.** (2005). Descriptive and interpretative approaches to qualitative research. In J. Miles and P. Gilbert (eds), *A Handbook of Research Methods in Clinical and Health Psychology* (pp. 147–160). Oxford: Oxford University Press. Retrieved from: http://nideffer.net/classes/GCT_RPI_S14/readings/interpretive.pdf
- Gega, L., Smith, J. and Reynolds, S.** (2013). Cognitive behaviour therapy (CBT) for depression by computer vs. therapist: patient experiences and therapeutic processes. *Psychotherapy Research*, 23, 218–231. doi: [10.1080/10503307.2013.766941](https://doi.org/10.1080/10503307.2013.766941)
- Gulliver, A., Griffiths, K. M., Christensen, H. and Brewer, J. L.** (2012). A systematic review of help-seeking interventions for depression, anxiety and general psychological distress. *BMC Psychiatry*, 12, 1. doi: [10.1186/1471-244X-12-81](https://doi.org/10.1186/1471-244X-12-81)
- Jones, S. L.** (2014). *An efficacy trial of therapist-assisted internet-delivered cognitive-behaviour therapy for older adults with generalized anxiety*. Unpublished doctoral dissertation, University of Regina, Canada. Retrieved from: <http://ourspace.uregina.ca/handle/10294/5818>
- Judd, L., Kessler, R., Paulus, M., Zeller, P., Wittchen, H. and Kunovac, J.** (1998). Comorbidity as a fundamental feature of generalized anxiety disorders: results from the National Comorbidity Study (NCS). *Acta Psychiatrica Scandinavica Supplement*, 393, 6–11. doi: [10.1111/j.1600-0447.1998.tb05960.x](https://doi.org/10.1111/j.1600-0447.1998.tb05960.x)
- Linehan, M. M.** (1997). Validation and Psychotherapy. In A. C. Bohart and L. S. Greenberg (eds), *Empathy Reconsidered: New Directions in Psychotherapy* (pp. 353–392). doi: [10.1037/10226-016](https://doi.org/10.1037/10226-016)
- Llewellyn, S. P.** (1988). Psychological therapy as viewed by clients and therapists. *British Journal of Clinical Psychology*, 27, 223–237.
- Mowbray, C. T., Mandiberg, J. M., Stein, C. H., Kopels, S., Curlin, C., Megivern, D., Strauss, S., Collins, K. and Lett, R.** (2006). Campus mental health services: recommendations for change. *American Journal of Orthopsychiatry*, 76, 226–237. doi: [10.1037/0002-9432.76.2.226](https://doi.org/10.1037/0002-9432.76.2.226)
- Richards, D., Dowling, M., O'Brien, E., Viganò, N. and Timulak, L.** (2017). Significant events in an internet-delivered (Space from Depression) intervention for depression. *Counselling and Psychotherapy Research*. doi: [10.1002/capr.12142](https://doi.org/10.1002/capr.12142)
- Richards, D. and Richardson, T.** (2012). Computer-based psychological treatments for depression: a systematic review and meta-analysis. *Clinical Psychology Review*, 332, 329–342. doi: [10.1016/j.cpr.2012.02.004](https://doi.org/10.1016/j.cpr.2012.02.004)
- Richards, D., Richardson, T., Timulak, L. and McElvaney, J.** (2015). The efficacy of internet-delivered treatment for generalized anxiety disorder: a systematic review and meta-analysis. *Internet Interventions*, 2, 272–282. doi: [10.1016/j.invent.2015.07.003](https://doi.org/10.1016/j.invent.2015.07.003)
- Richards, D. and Timulak, L.** (2012). Satisfaction with therapist-delivered vs. self-administered online cognitive behavioural treatments for depression symptoms in college students. *British Journal of Guidance and Counselling*, 41, 193–207. doi: [10.1080/03069885.2012.726347](https://doi.org/10.1080/03069885.2012.726347)
- Richards, D., Timulak, L., Doherty, G., Sharry, J., McLoughlin, O., Rashleigh, C., Colla, A. and Joyce, C.** (2014). Low-intensity internet-delivered treatment for generalized anxiety symptoms in routine care: protocol for a randomized controlled trial. *Trials*, 15. doi: [10.1186/1745-6215-15-145](https://doi.org/10.1186/1745-6215-15-145)

- Richards, D., Timulak, L. and Hevey, D.** (2013). A comparison of two online cognitive-behavioural interventions for symptoms of depression in a student population: the role of therapist responsiveness. *Counselling and Psychotherapy Research*, 13, 184–193. doi: [10.1080/14733145.2012.733715](https://doi.org/10.1080/14733145.2012.733715)
- Richards, D., Timulak, L., Rashleigh, C., McLoughlin, O., Colla, A., Joyce, C., Doherty, G., Sharry, J., Duffy, D. and Anderson-Gibbons, M.** (2016). Effectiveness of an internet-delivered intervention for generalized anxiety disorder in routine care: a randomised controlled trial in a student population. *Internet Interventions*, 1–9. doi: [10.1016/j.invent.2016.10.003](https://doi.org/10.1016/j.invent.2016.10.003)
- Ritterband, L. M., Gonder-Frederick, L. A., Cox, D. J., Clifton, A. D., West, R. W. and Borowitz, S. M.** (2003). Internet interventions: in review, in use, and into the future. *Professional Psychology: Research and Practice*, 34, 527. doi: [10.1037/0735-7028.34.5.527](https://doi.org/10.1037/0735-7028.34.5.527)
- Roehlen, A. B., Zack, J. S. and Speyer, C.** (2004). Online therapy: review of relevant definitions, debates, and current empirical support. *Journal of Clinical Psychology*, 60, 269–283. doi: [10.1002/jclp.10263](https://doi.org/10.1002/jclp.10263)
- Spitzer, R. L., Kroenke, K., Williams, J. B. and Löwe, B.** (2006). A brief measure for assessing generalized anxiety disorder: the GAD-7. *Archives of Internal Medicine*, 166, 1092–1097. doi: [10.1001/archinte.166.10.1092](https://doi.org/10.1001/archinte.166.10.1092)
- Somers, J. M., Goldner, E. M., Waraich, P. and Hsu, L.** (2006). Prevalence and incidence studies of anxiety disorders: a systematic review of the literature. *Canadian Journal of Psychiatry*, 51, 100–113. Retrieved from: <http://www1.cpa-apc.org:8080/Publications/Archives/CJP/2006/february/sommers-RP.asp>
- Thomas, N., McLeod, B., Jones, N. and Abbott, J. A.** (2015). Developing internet interventions to target the individual impact of stigma in health conditions. *Internet Interventions*, 2, 351–358. doi: [10.1016/j.invent.2015.01.003](https://doi.org/10.1016/j.invent.2015.01.003)
- Timulak, L.** (2007). Identifying core categories of client-identified impact of helpful events in psychotherapy: a qualitative meta-analysis. *Psychotherapy Research*, 17, 305–314. doi: [10.1080/10503300600608116](https://doi.org/10.1080/10503300600608116)
- Timulak, L.** (2010). Significant events in psychotherapy: an update of research findings. *Psychology and Psychotherapy: Theory, Research and Practice*, 83, 421–447. doi: [10.1348/147608310X499404](https://doi.org/10.1348/147608310X499404)
- Titov, N., Dear, B. F., Johnston, L., Lorian, C., Zou, J., Wootton, B., Spence, J., McEvoy, P. M. and Rapee, R. M.** (2013). Improving adherence and clinical outcomes in self-guided internet treatment for anxiety and depression: randomised controlled trial. *PLoS One*, 8, e62873. doi: [10.1371/journal.pone.0062873](https://doi.org/10.1371/journal.pone.0062873)
- Tyrer, P. and Baldwin, D.** (2006). Generalised anxiety disorder. *The Lancet*, 368, 2156–2166. doi: [10.1016/S0140-6736\(06\)69865-6](https://doi.org/10.1016/S0140-6736(06)69865-6)