

## SHORT REPORT

# RNAi of *COL1A1* in mesenchymal progenitor cells

Sophia Millington-Ward<sup>\*1</sup>, Helena P McMahon<sup>1</sup>, Danny Allen<sup>1</sup>, Gearóid Tuohy<sup>1</sup>, Anna-Sophia Kiang<sup>1</sup>, Arpad Palfi<sup>1</sup>, Paul F Kenna<sup>1</sup>, Peter Humphries<sup>1</sup> and G Jane Farrar<sup>1</sup>

<sup>1</sup>Ocular Genetics Unit, Department of Genetics, Trinity College Dublin, Dublin 2, Ireland

Given that mutant *COL1A1* is known to cause Osteogenesis Imperfecta (OI), tools to modulate *COL1A1* expression are likely to be of significant therapeutic value. In this context, we have evaluated RNA interference (RNAi) as a means to downregulate *COL1A1* expression in Cos-7 cells and in human mesenchymal progenitor stem cells (MPCs), the latter cells giving rise to bone and therefore representing a target cell type for collagen-related disorders. In addition, allele-specificity, a key factor to the success of RNAi-based suppression, was explored with a view to developing a mutation-independent RNAi-based therapeutic for OI by targeting an intragenic SNP within transcripts derived from the *COL1A1* gene. Preferential suppression of individual polymorphic alleles that differed by a single nucleotide was observed.

*European Journal of Human Genetics* (2004) 12, 864–866. doi:10.1038/sj.ejhg.5201230

Published online 7 July 2004

**Keywords:** RNAi; polymorphism; *COL1A1*; gene therapy; stem cell; Osteogenesis Imperfecta

## Introduction

Over 150 *COL1A1* mutations give rise to the brittle bone disease Osteogenesis Imperfecta (OI) (<http://www.le.ac.uk/genetics/collagen/>). It would therefore be preferable to suppress mutant *COL1A1* alleles in a mutation-independent manner.<sup>1–4</sup> Clearly, efficiency and specificity are key factors in the efficacy of a suppression-based therapeutic and RNA interference (RNAi) represents a powerful new tool.<sup>5</sup> However, some studies show that RNAi may not be sequence specific at the level of a single nucleotide.<sup>5</sup> It is therefore imperative, when contemplating using RNAi-based therapies, to determine whether allele-specificity really can be achieved.

Mesenchymal progenitor stem cells (MPCs) are stem cells, which can give rise to bone and which represent target cells for OI therapies. Engraftment of donor MPCs has been achieved in children with OI.<sup>6</sup> However, the use of a patient's own MPCs transduced with a therapeutic gene may aid in the cells' long-term survival. To date, RNAi

has not been shown in MPCs. In this study, short interfering RNAs (siRNAs<sup>5</sup>) targeting *COL1A1* have been evaluated in MPCs. siRNAs were also assessed for their ability to distinguish between two *COL1A1* targets that differ by a single nucleotide (2pq=0.41, SNP ID: rs1061237<sup>7,1,2,4</sup>). Such sequence specificity would enable generation of two siRNAs targeting each of the polymorphic variants of *COL1A1*, which together could be used to treat 40% of OI patients with dominant-negative *COL1A1* mutations – those patients who are heterozygous for the polymorphism.

## Materials and methods

### siRNAs

siRNA target sequences (below) differed by at least four nucleotides from any known gene (BLASTN2.2.6<sup>8</sup>). siRNAs were synthesised<sup>5</sup> (Qiagen-Xeragon, Germantown, MD 20874, USA). The C/T polymorphism is highlighted in bold print. siRNA7a1, targeting human *COL7A1*, was used as a negative control.

siRNAT1 target: 5' AA CTGAACCCCTCAAAGCC 3'  
(4592–4610 of sense strand Z74615)

\*Correspondence: Dr S Millington-Ward, Ocular Genetics Unit, Department of Genetics, Trinity College Dublin, Dublin 2, Ireland. Tel: +353 1 6082482; Fax: +353 1 6083848; E-mail: sophia@maths.tcd.ie  
Received 2 December 2003; revised 16 April 2004; accepted 22 April 2004

siRNAT1 sense: 5' CUGAACCCCCUCAAAGCC TT 3'  
 siRNAT1 antisense:  
 3' TT GACUUGGGGAGUUUUCGG 5'  
 siRNAT2 target: 5' CC CCTCAAAGCCAAAAAATG  
 3'(4600–4618 of sense strand Z74615)  
 siRNAT2 sense: 5' CCUCAAAGCCAAAAAUG GG 3'  
 siRNAT2 antisense:  
 3' GG GGAGUUUUCGGUUUUUAC 5'  
 siRNAC target: 5' CC CCCCAAAGCCAAAAAATG  
 3'(4600–4618 of sense strand Z74615)  
 siRNAC sense: 5' CCCCAAAGCCAAAAAUG GG 3'  
 siRNAC antisense:  
 3' GG GGGUUUUCGGUUUUUAC 5'  
 siRNA7a1 target: 5' AA GGGGCAGGGGTCAAGCTA  
 3'(588–516 of sense strand NM000094)  
 siRNA7a1 sense: 5' GGGGCAGGGGUCAAGCUA TT 3'  
 siRNA7a1 antisense:  
 3' TT CCCCGUCCCCAGUUCGAU 5'

#### Target *COL1A1* constructs

siRNA targets (15940–16620 Acc. No. AF017178) with either the T- or C-allele of human *COL1A1* were cloned into pIRES2-EGFP (Clontech Laboratories Inc., Palo Alto, USA Cat. No. 6029-1).

#### Transient transfections of Cos-7 cells and MPCs

Cos-7 cells were transfected in quadruplicate in 2 cm<sup>2</sup> wells with 0.8 µg of target and 50 nmol of siRNA using Lipofectamine 2000. RNA was isolated<sup>4</sup> after 48 h. MPCs<sup>9</sup> were transfected with 50 nmol of siRNA using Oligofectamine (Invitrogen Cat. No. 12252-011). RNA was harvested after 7 h. When two siRNAs were cotransfected, 25 nmol of each was used.

#### Real time RT-PCR

Quadruplicate transfection samples were pooled and *COL1A1* RNA levels analysed by real time RT-PCR, standardised to *β-actin* or *GAPDH*.<sup>4</sup> *β-Actin* expression profile was checked using *GAPDH* and *COL1A2*.<sup>4</sup>

#### ELISA assays

MPC-T/T (below) cells were transfected in quadruplicate with siRNAs. Quadruplicate samples were pooled and ELISAs were performed, in duplicate, 28 h post-transfection.<sup>4</sup>

#### Statistical analysis

ANOVAs and least significant difference tests (LSDs) were performed using Data Desk 6.0 (Data Descriptions Inc., New York, USA). Differences were significant at  $P \leq 0.05$ . Tables include standard errors.

#### Results

##### siRNA studies for OI therapeutics

Three mutation-independent siRNAs targeting a polymorphic region in the *COL1A1* gene were evaluated in human MPCs. siRNAT1 and siRNAT2 were designed to target the T-allele variant whereas siRNAC targets the C-allele variant of *COL1A1*.<sup>7,1,2,4</sup> MPCs were homozygous (MPC-T/T) or heterozygous (MPC-C/T) for the polymorphism.

All siRNAs suppressed *COL1A1* (Table 1). siRNAT2 downregulates the T-allele of *COL1A1* in MPC-T/T cells more than either siRNAT1 or siRNAC (84.62, 65.23, 67.66 respectively) ( $P=0.004$  and  $0.01$ ). siRNAT1 and siRNAC are equally active in these cells even though siRNAC has a single-base mismatch with respect to its target ( $P=0.7$ ). When both siRNAT1 and siRNAC or siRNAT2 and siRNAC are transfected together, lower levels of suppression are achieved than with either siRNAT2 or siRNAC alone, suggesting that the two siRNAs may compete/interfere with one another.

In MPC-C/T cells, siRNAT2 and siRNAC are better than siRNAT1 at suppressing *COL1A1* ( $P<0.0001$ ). Unlike the results found in MPC-T/T cells, siRNAC is better than siRNAT2 at downregulating the combined C- and T-allele transcripts found in the MPC-C/T cells ( $P<0.0001$ ). This suggests that siRNAC may be better at downregulating the

**Table 1** *COL1A1* expression levels in MPC-T/T, MPC-C/T, and Cos-7 cells after siRNA transfections

siRNA	<i>COL1A1</i> levels in MPC-T/T (%)	n, p	<i>COL1A1</i> levels in MPC-C/T (%)	n, p
siRNA 7a1	100 ± 3.28	8	100 ± 2.67	16
siRNA T1	34.77 ± 6.66	8, <0.0001	73.44 ± 3.40	10, <0.0001
siRNA T2	15.38 ± 2.50	8, <0.0001	51.97 ± 2.46	10, <0.0001
siRNA C	32.34 ± 3.52	8, <0.0001	17.62 ± 1.24	8, <0.0001
siRNA T1+siRNA C	45.95 ± 5.21	8, <0.0001	51.92 ± 2.64	10, <0.0001
siRNA T2+siRNA C	46.15 ± 4.50	8, <0.0001	39.88 ± 1.14	8, <0.0001
siRNA	T-allele levels in Cos-7 (%)	n, p	C-allele levels in Cos-7 (%)	n, p
siRNA 7a1	100 ± 1.46	9	100 ± 0.95	9
siRNA T1	51.64 ± 3.36	9, <0.0001	65.35 ± 1.48	9, <0.0001
siRNA T2	35.54 ± 1.97	8, <0.0001	38.61 ± 2.00	9, <0.0001
siRNA C	54.22 ± 2.55	9, <0.0001	47.98 ± 2.93	9, <0.0001

**Table 2** Type I collagen protein expression levels in MPCs after siRNA transfections

siRNA or shRNA	Type I collagen (ng/ml)	Type I collagen levels in MPCs (%)
siRNA 7a1	83.04	100
siRNA T1	51.48	62
siRNA T2	23.04	28
siRNA C	43.20	52
siRNA T1+siRNA C	34.08	41
siRNA T2+siRNA C	29.76	36

C-allele of *COL1A1* than siRNAT2 is at downregulating the T-allele. This is corroborated by the fact that lower levels of suppression are achieved with both siRNAT2 and siRNAC than with siRNAC singly ( $P < 0.0001$ ). When siRNAT1 and siRNAC or siRNAT2 and siRNAC were evaluated in MPCs together, an intermediate level of suppression to that obtained with each siRNA singly was found.

Results quantifying *de novo* formed Type I collagen protein, excreted from MPC-T/T cells transfected with siRNAs, provide evidence that suppression of RNA is mirrored at the protein level (Table 2). Again, siRNAT2 was the most efficient siRNA, followed by siRNAC and siRNAT1. Protein levels were suppressed by up to 72%.

Since MPC cells with the rare homozygous C-allele genotype were not available, siRNAs were evaluated in Cos-7 cells expressing a partial C- or T-allele *COL1A1* gene. Downregulation of up to 65% was observed. As found in MPCs, siRNAT2 was more active than siRNAT1. In addition, all three siRNAs targeted their own correct target allele better than their incorrect allele, showing allele preference rather than allele-specificity (Table 1).

## Discussion

The use of mutation-independent siRNAs targeting a common and therefore potentially therapeutically significant polymorphism in the *COL1A1* gene has been investigated. While levels of *COL1A1* suppression of up to 85% were observed, none of the siRNAs evaluated were found to be completely allele-specific. Specificity of siRNA at the single-nucleotide resolution has been achieved in a number of studies.<sup>5,10–12</sup> One suggests that to achieve sequence-specificity, small amounts of siRNA should be used.<sup>5</sup> Another study does not confirm this.<sup>5</sup> The positioning of a single-nucleotide mismatch within the siRNA molecule at position 9, 10 or 11 of the target (from the 5' end of the target and not including 3' siRNA overhang sequences) may convey specificity.<sup>5,10,11</sup> In our study the single mismatch was situated at position 3 (siRNAT2 and siRNAC) or 11 (siRNAT1). Constraints due to *COL1A1*

target sequence disallowed the study of additional siRNAs targeting this polymorphism.

In summary, RNAi-based suppression of *COL1A1* has been demonstrated in MPCs and allele-specificity evaluated. Clearly, in many cases, the lack of allele-specificity will be an enormous barrier when using RNAi to suppress mutant alleles. In addition, constraints incurred by both the target sequence and the necessity to design molecules that will not target multiple genes will be challenging when designing RNAi therapeutics. The use of mutation-independent RNAi targets, such as polymorphic sites, is therefore appealing.

## Acknowledgements

We thank C Allers and P Conget for providing MPCs, and the CBBF, the Health Research Board Ireland and Enterprise Ireland for their support.

## References

- 1 Millington-Ward S, O'Neill B, Tuohy G *et al*: Strategems *in vitro* for gene therapies directed to dominant mutations. *Hum. Mol. Genet.* 1997; **6**: 1415–1426.
- 2 Millington-Ward S, O'Neill B, Kiang AS, Humphries P, Kenna PF, Farrar GJ: A mutation-independent therapeutic strategem for osteogenesis imperfecta. *Antisense Nucleic Acid Drug Dev.* 1999; **9**: 537–542.
- 3 O'Neill B, Millington-Ward S, O'Reilly M *et al*: Ribozyme-based therapeutic approaches for autosomal dominant retinitis pigmentosa. *Invest. Ophthalmol. Vis. Sci.* 2000; **41**: 2863–2869.
- 4 Millington-Ward S, Allers C, Tuohy G *et al*: Validation in mesenchymal progenitor cells of a mutation-independent *ex vivo* approach to gene therapy for osteogenesis imperfecta. *Hum. Mol. Genet.* 2002; **11**: 2201–2206.
- 5 Dykxhoorn DM, Novina CD, Sharp PA: Killing the messenger: short RNAs that silence gene expression. *Nat. Rev. Mol. Cell Biol.* 2003; **4**: 457–467.
- 6 Horwitz EM, Prockop DJ, Fitzpatrick LA *et al*: Transplantability and therapeutic effects of bone marrow-derived mesenchymal cells in children with osteogenesis imperfecta. *Nat. Med.* 1999; **5**: 309–313.
- 7 Westerhausen AI, Constantinou CD, Prockop DJ: A sequence polymorphism in the 3'-nontranslated region of the pro alpha 1 chain of type I procollagen. *Nucleic Acids Res.* 1990; **18**: 4968.
- 8 Altschul SF, Madden TL, Schaffer *et al*: Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* 1997; **25**: 3389–3402.
- 9 Conget PA, Minguell JJ: Phenotypical and functional properties of human bone marrow mesenchymal progenitor cells. *J. Cell Physiol.* 1999; **181**: 67–73.
- 10 Ding H, Schwarz DS, Keene A *et al*: Selective silencing by RNAi of a dominant allele that causes amyotrophic lateral sclerosis. *Aging Cell* 2003; **2**: 209–217.
- 11 Miller VM, Xia H, Marrs GL *et al*: Allele-specific silencing of dominant disease genes. *Proc. Natl. Acad. Sci. USA* 2003; **100**: 7195–7200.
- 12 Abdelgany A, Wood M, Beeson D: Allele-specific silencing of a pathogenic mutant acetylcholine receptor subunit by RNA interference. *Hum. Mol. Genet.* 2003; **12**: 2637–2644.