
BRIEF COMMUNICATION

Detection of prospective memory deficits in mild cognitive impairment of suspected Alzheimer's disease etiology using a novel event-based prospective memory task

ALBERTO BLANCO-CAMPAL,^{1,2} ROBERT F. COEN,³ BRIAN A. LAWLOR,³ JOSEPH B. WALSH,³
AND TERESA E. BURKE¹

¹School of Psychology, University College Dublin, Dublin, Ireland

²North Eastern Area of the Health Service Executive, Dundalk, Ireland

³Mercer's Institute for Research on Ageing, St James's Hospital, Dublin, Ireland

(RECEIVED April 21, 2008; FINAL REVISION September 20, 2008; ACCEPTED September 29, 2008)

Abstract

We investigated the relative discriminatory efficacy of an event-based prospective memory (PM) task, in which specificity of the instructions and perceptual salience of the PM cue were manipulated, compared with two widely used retrospective memory (RM) tests (Rivermead Paragraph Recall Test and CERAD-Word List Test), when detecting mild cognitive impairment of suspected Alzheimer's disease etiology (MCI-AD) ($N = 19$) from normal controls (NC) ($N = 21$). Statistical analyses showed high discriminatory capacity of the PM task for detecting MCI-AD. The Non-Specific-Non-Salient condition proved particularly useful in detecting MCI-AD, possibly reflecting the difficulty of the task, requiring more strategic attentional resources to monitor for the PM cue. With a cutoff score of $<4/10$, the Non-Specific-Non-Salient condition achieved a sensitivity = 84%, and a specificity = 95%, superior to the most discriminative RM test used (CERAD-Total Learning: sensitivity = 83%; specificity = 76%). Results suggest that PM is an early sign of memory failure in MCI-AD and may be a more pronounced deficit than retrospective failure, probably reflecting the greater self-initiated retrieval demands involved in the PM task used. Limitations include the relatively small sample size, and the use of a convenience sample (i.e. memory clinic attenders and healthy active volunteers), reducing the generalizability of the results, which should be regarded as preliminary. (*JINS*, 2009, 15, 154–159.)

Keywords: Prospective memory, Mild cognitive impairment, Memory disorders, Neuropsychological tests, Dementia, Early detection

INTRODUCTION

Prospective memory (PM) is remembering to carry out intended actions (such as remembering to take medication) at an appropriate point in the future (McDaniel & Einstein, 2007). A critical difference between PM and retrospective memory (RM) tasks (e.g., list-learning or story recall), is that there is no external agent requesting a memory search when the target event occurs (McDaniel & Einstein, 2000). It is generally accepted that PM requires a greater degree of self-initiated retrieval operations, compared with RM tasks,

requiring inhibition of an ongoing task, and switching to another action (Einstein & McDaniel, 1996).

Einstein and McDaniel (1996) distinguished between *time-based tasks*, requiring performance of actions at a certain time and *event-based tasks*, requiring performance of actions in response to external events. For event-based tasks, encoding factors such as specificity of instructions and retrieval factors such as perceptual salience of PM cues are associated with superior prospective remembering. The multiprocess model of PM argues that both factors reduce the self-initiated retrieval demands of PM tasks (McDaniel & Einstein, 2000). These factors have been investigated in older healthy adults (Henry et al., 2004), but not in individuals with Mild Cognitive Impairment (MCI).

MCI is a broad syndrome characterized by impaired cognitive performance that lies on the continuum between normal

Correspondence and reprint requests to: Alberto Blanco-Campal, School of Psychology, University College Dublin, Belfield, Dublin 4, Ireland.
E-mail: Alberto.Blanco-Campal@hse.ie

ageing and a dementing process, and is associated with increased risk of developing dementia (Kluger et al., 2000). MCI represents a heterogeneous group presenting with different cognitive profiles and underlying etiologies (Petersen et al., 2001). We focus on Mild Cognitive Impairment-Alzheimer's Disease (MCI-AD), a term describing individuals with MCI whose underlying disease is clinically suspected to be AD (Portet et al., 2006).

Only two studies have previously investigated PM in MCI. Kazui et al. (2005) found that while single-trial event-based PM tasks proved sensitive to MCI, the degree of specificity was not as robust, concluding that PM tasks were not useful for detecting MCI. More recently, Troyer and Murphy (2007) found that a time-based PM task was particularly sensitive to MCI, but no data regarding the discriminatory capacity of this task were presented. Interestingly, Driscoll et al. (2005) found significant PM deficits in healthy elderly APOE $\epsilon 4$ gene carriers, known to be associated with increased risk and decreased age of onset for AD relative to noncarriers.

The present study intends to fill current gaps in the literature by focusing on the relative vulnerability of PM and RM to the cognitive impairment in MCI-AD. Our primary hypotheses were that (1) An event-based PM task would show higher discriminatory efficacy to detect MCI-AD from NC relative to RM tests due to its higher self-initiated retrieval demands. (2) Conditions of non-specificity of instructions and perceptual non-salience of the PM cue would be particularly sensitive to MCI-AD because they require greater strategic attentional resources to monitor the PM cue.

METHOD

Participants

Participants were 21 normal controls (NC), and 19 individuals with a diagnosis of MCI-AD, in accordance with the Report of the MCI Working Group of the European Consortium on Alzheimer's disease (Portet et al., 2006) (see Table 1).

NC were recruited from two local Active Retirement Groups (community-based social groups). Participants qualified as NC if they scored $\geq 27/30$ on the Mini-Mental State Examination (MMSE; Folstein et al., 1975), >8 on Clock Drawing (Manos, 1999), ≥ 7 on the Geriatric Depression Scale-15 item Short Form (Sheikh & Yesavage, 1986), and were functioning normally in the community as documented by self-report on the Lawton-Brody Instrumental Activities of Daily Living (Lawton & Brody, 1969). MCI-ADs were identified through a Hospital Memory Clinic (Swanwick et al., 1996). Methods were approved by the Hospital's ethics board. Before enrolment in the study, patients underwent a neuropsychological evaluation using a range of standardized cognitive tests including Cambridge Cognitive Examination (CAMCOG; Huppert et al., 1995); Delayed Word Recall (DWR) test (Coen et al., 1997; Knopman & Ryberg, 1989), and executive functions, using both phonemic (FAS; Benton & Hamsher, 1978), and semantic (animals, fruits, vegetables; Monsch et al., 1992) verbal fluency tasks.

Table 1. Demographic and clinical details of the two groups of participants

	NC ($n = 21$)	MCI-AD ($n = 19$)
	Mean (SD)	Mean (SD)
Age at testing (years)		
Mean (SD)	72.5 (5.6)	71.05 (5.6)
Range	(63–79)	(61–80)
Gender		
Male (%)	6 (28.6%)	9 (47%)
Female (%)	15 (71.4%)	10 (53%)
MMSE		
Mean (SD)	29.4 (0.7)	25.72 (1.97)
Education ^a		
≤ 13 Years	13	16
>13 Years	8	3
NART-R		
Mean (SD)	110 (6.6)	106 (11.4)
Range	(100–122)	(85–124)
GDS-15	1.14 (1.52)	2.58 (2.78)

Note. MCI-AD = mild cognitive impairment-Alzheimer's disease; NC = normal controls; SD = standard deviation; MMSE = Mini-Mental State Exam; NART-R = National Adult Reading Test-Revised; GDS-15 = Geriatric Depression Scale.

^a In Ireland, completion of Secondary Level (High-School equivalent), entails 13 years of formal education.

Exclusion criteria for both groups included a diagnosis of dementia (DSM-IV-TR; American Psychiatric Association, 2000), primary psychiatric disorder, history of alcohol or substance dependence, cognitive impairment caused by concomitant medications, or the presence of a major neurological illness.

The two groups were matched on age [$t(38) = 0.83$; $p = .41$], gender [$\chi^2 = 1.50$; $df = 1$; $p = .22$], years of education [$\chi^2 = 2.48$; $df = 1$; $p = .115$], estimated premorbid intellect (NART-R; Nelson, 1991) [$t(38) = 1.33$; $p = .19$], and affect (GDS-15) [$u = 140$; $p = .174$]. As expected, the NC group had higher scores on MMSE [$t(38) = 8.01$; $p \leq .0001$] (see Table 1).

Measures

Retrospective memory tests. Both groups were assessed using two RM tests, not part of the diagnostic process: The Rivermead Paragraph Recall Test from the Rivermead Behavioural Memory Test (RBMT-PRT; Wilson et al., 1985), and the Word List Memory subtest from the Consortium to Establish a Registry for Alzheimer's disease (CERAD-Word List; Welsh et al., 1994).

The RBMT-PRT is a paragraph with 21 separate ideas, read to the subject who must recall it immediately and after a 10- to 15-min delay. The CERAD-Word List has 10 words presented over three trials. Participants read out each word in turn, followed by immediate recall. Delayed recall is assessed after a 5-min delay.

The Silly Sentences PM Task. This was a novel event-based PM task using a dual-task paradigm where a PM intention was embedded in an ongoing task.

Ongoing Task. This consisted of a lexical decision task, preventing continuous rehearsal of the PM intention. Participants were shown two series of 108 short sentences, in which two different conditions of the Silly Sentences PM task were embedded.

Participants were told that we were interested in determining their ability to verify a series of sentences, individually presented in the middle of the computer screen with the alternatives true/false written underneath, and that sentences would automatically appear and disappear. Participants were asked to read each sentence and to say “true” (e.g., Africa is home to tigers), or “false” (e.g., lions live in hotels) as appropriate, and told that no immediate performance feedback would be provided. Sentences were inspired by the Silly Sentences Test, an instrument measuring speed of language comprehension (Baddeley et al., 1992).

To control for differences in information processing-speed, and based on piloting results, rate of presentation was 6 s per sentence for NC and 8 s for MCI-AD. This was done to avoid possible processing-speed deficits precluding the noticing of the PM cue in MCI-AD. While it could be argued that this introduces variations in the retention delay between critical sentences, existing research has failed to find increased forgetting with delay variations of just several minutes or more (Brandimonte & Passolunghi, 1994). Sentences were projected on a 15.1-inch screen laptop, displayed in lower case, (Times Roman, 44-point), on a white background. Participants were initially presented with six practice sentences (three true, three false) to familiarize them with the procedure.

Prospective memory task. Following the practice trials, participants were told that we were also interested in studying their ability to remember to do something in the future.

With the aim of creating a task that placed a differential degree on strategic attentional resources, two variables were manipulated (i.e., specificity of the instructions and perceptual salience of the PM cue) (see Table 2).

In the *Specific condition*, participants were given specific instructions concerning the PM cue (e.g., when you see the word “lions” written in any of the sentences, remember to say “animals”). Ten critical sentences containing the PM cue lions were embedded throughout the first 108 sentences. Within the specific condition, the perceptual salience of the recall cue (RC) was varied so that 5 of 10 critical sentences were presented with the word “lions” in italics (i.e., Specific-Salient condition), and the other 5 in regular format (i.e., Specific-Non-Salient). Presentation of salient and non-salient sentences was intermixed.

In the *Non-Specific condition*, participants were given general instructions (i.e., when you see the name of any type of animal remember to say “animals”). Ten critical sentences containing the PM cue of an exemplar of the category “animals” were embedded throughout the second 108 verification sentences’ trial. Each one of the 10 PM cues was only shown once, and exemplars were equated against the exemplar “lion” in terms of their typicality using Irish-based norms (Brown & Semrau, 1986). Within the Non-Specific condition, the perceptual salience of the RC was also varied so that 5 of 10 critical sentences were presented with the name of the animal in italics (e.g., *horses*) (i.e., Non-Specific-Salient condition), and the other 5 in regular format (e.g., elephants) (i.e., Non-Specific-Non-Salient). Presentation of these sentences was also intermixed.

Procedure. Participants were tested individually under all testing conditions. The two PM tasks (i.e., Specific and Non-Specific) were introduced separately, and order of administration

Table 2. Parameters of the prospective memory task and performance of NC and MCI-AD

Experimental condition	PM Intention	Examples	Ongoing task	PM task type (Processing)	PM trials	Correct PM responses	
						MCI-AD	NC
Specific							
Salient	When you see the word	Pens are smaller than <i>lions</i> <i>Lions</i> have long stripes	Lexical Decision Task (Silly Sentences)	^a Focal +	5	67.4%	100%
Non-Salient	lions remember to say animals	Airplanes travel faster than lions Lions live in hotels		^a Focal -	5	62.2%	99%
Total 10						64.7%	99.5%
Non-Specific							
Salient	When you see the name of any	Africa is home to <i>tigers</i> <i>Horses</i> have stripes	Lexical Decision Task (Silly Sentences)	^b Non-focal ↓	5	37.8%	95%
Non-Salient	type of animal remember to say animals	Tusks form part of elephants Cats are manufactured goods		^b Non-focal ↑	5	28.4%	94%
Total 10						33.2%	95%

Note. ^aHigh in focal processing; ⁻Low in focal processing; [↓]Low in non-focal processing; [↑]High in non-focal processing. NC = normal controls; MCI-AD = mild cognitive impairment-Alzheimer’s disease.

^a The PM cue overlaps with the information constellation relevant to performing the ongoing task (McDaniel et al., 2008).

^b The PM cue is present in the environment but not part of the information being considered by the person (McDaniel et al., 2008).

was counterbalanced. In both conditions, participants were told to keep the instructions in mind because they would not be repeated and were asked to repeat back the instructions to ensure their learning. Participants were not told how often the PM cue would appear. Critical sentences appeared relatively equally throughout the verification sentences (on average every 10 sentences, at a rate of 1.5 min approximately). Cues were counterbalanced for location (i.e., beginning or end of sentences), and verification form (i.e., true or false). In both conditions, participants were told that making the PM response precluded the need to do the sentence verification task. To minimize the RM demands of the task, the intended action (i.e., say animals) was held constant and highly associated with the PM cue in both conditions. Following the PM task, it was verified that 100% of individuals remembered the instructions. Total time of administration for each condition (Specific/Non-Specific) was 14 min.

Statistical Analyses

Data distribution in the PM task and several conditions of the RM tests was skewed. For this reason, either nonparametric tests (Mann-Whitney *U*), or their parametric equivalents (*t* test), were used, as appropriate, to analyze group differences.

Analysis of group differences. A compromise approach (setting *alpha* (α) level for statistical significance (*p*) at $\alpha = 0.01$) to balance the risk of Type I and Type II errors was adopted (Aron et al., 2006).

Discriminatory efficacy of prospective and retrospective memory. A series of Receiver Operating Characteristic Curve (ROC) analyses was produced to examine the relative discriminative efficacy of PM tasks compared with RM tests. The cut-off scores providing the highest sensitivity balanced against optimum specificity were selected. Positive predictive values

(PPV = true positives / [true positives + false positives]), negative predictive values (NPV = true negatives / [true negatives + false negatives]), and overall accuracy (i.e., percentage of MCI cases accurately categorized as MCI + percentage of NCs correctly classified as NC) were also calculated.

RESULTS

Prospective and Retrospective Memory: Analysis of Group Differences

There were significant differences between the NC and MCI-AD groups on all the RM tests and all PM conditions, reaching our criterion ($p < .01$) on all but RBMT (PRT)-Savings (see Table 3). These differences remained even when education was accounted for (i.e., analysis of covariance and case summaries).

Discriminatory Capacity of PM tasks and RM Tests: Detecting MCI-AD From NC

ROC analysis was significant for all measures of memory administered except for RBMT-PRT-Savings. Overall accuracy (OA) ranged from 69% to 90%. With the exception of the Silly Sentences-Specific-Salient condition, which showed poor sensitivity to MCI-AD (<55%), all other measures demonstrated reasonable sensitivity and specificity. Notably, the Silly Sentences-Non-Specific-Total score (sensitivity = 84%; specificity = 95%) and its individual conditions, the Silly Sentences Non-Specific-Non-Salient (sensitivity = 84%; specificity = 95%), and Silly Sentences-Non-Specific-Salient (sensitivity = 73%; specificity = 95%) demonstrated the highest discriminatory power to detect MCI-AD from NC, and was superior to all RM tests, certainly in terms of specificity and overall accuracy (see Table 4).

Table 3. Analysis of group differences on memory functioning

Test	NC (<i>n</i> = 21) <i>M</i> (<i>SD</i>)	MCI-AD (<i>n</i> = 19) <i>M</i> (<i>SD</i>)	NC vs. MCI-AD <i>U/t</i>	% Difference MCI-AD vs. NC
Retrospective Memory				
CERAD-Word List-Total Learning	21.7 (4.0)	14.3 (4.9)	<i>t</i> = 5.20***	66%
CERAD-Word List-Delay Recall	7.4 (2.2)	4.1 (2.4)	<i>t</i> = 4.37***	55%
CERAD- Word List-Savings	85.0 (21.3)	60.0 (31)	<i>U</i> = 86.00*	71%
RBMT(PRT)-Immediate Recall	9.0 (2.5)	5.28 (3)	<i>t</i> = 4.20***	59%
RBMT (PRT)-Delay Recall	7.3 (2.6)	3.1 (3.9)	<i>t</i> = 3.9***	43%
RBMT (PRT)-Savings	80.1 (18.3)	44.1 (40.5)	<i>U</i> = 86.00	55%
Prospective Memory				
Silly Sentences-Non-Specific-Total	9.4 (0.9)	3.3 (3.3)	<i>U</i> = 21.50***	35%
Silly Sentences-Non-Specific Salient	4.7 (0.5)	1.9 (1.8)	<i>U</i> = 43.00***	40%
Silly Sentences-Non-Specific-Non-Salient	4.7(0.6)	1.4 (1.7)	<i>U</i> = 20.50***	30%
Silly Sentences-Specific-Total	9.9 (0.2)	6.5 (3.8)	<i>U</i> = 57.50***	66%
Silly Sentences-Specific-Salient	5.0 (0)	3.4 (1.9)	<i>U</i> = 94.50*	68%
Silly Sentences-Specific-Non-Salient	4.9 (0.2)	3.1 (1.9)	<i>U</i> = 68.50***	63%

Note: NC = normal controls; MCI-AD = mild cognitive impairment-Alzheimer’s disease; CERAD = Consortium to Establish a Registry for Alzheimer’s Disease; RBMT (PRT) = Rivermead Behavioural Memory Test-Paragraph Recall Test; *M* = mean; *SD* = standard deviation. * Significant value ($p < .01$); **Significant value ($p < .001$); ***Significant value ($p < .0001$).

Table 4. Summary of the ROC analyses with cutoff scores for NC ($n = 21$) vs. MCI-AD ($n = 19$)

	Max. AUC (<i>SE</i>)	<i>p</i>	Cutoff score	Sensitivity	Specificity	OA	PPV	NPV
Retrospective Memory								
CERAD-Word List-Total Learning	0.87(0.04)	***	<19	83%	76%	80%	75%	84%
CERAD-Word List-Delay Recall	0.86(0.06)	***	<6	72%	86%	80%	81%	86%
CERAD- Word List-Savings	0.77(0.07)	*	<70%	56%	81%	69%	71%	68%
RBMT(PRT)-Immediate Recall	0.82(0.07)	**	<9	82%	71%	76%	70%	83%
RBMT (PRT)-Delay Recall	0.83(0.07)	**	<6	77%	81%	79%	77%	81%
RBMT (PRT)-Savings	0.75(0.08)	*	<69	65%	90%	79%	85%	76%
Prospective Memory								
Silly Sentences-Non-Specific-Total	0.96(0.03)	***	<8	84%	95%	90%	94%	87%
Silly Sentences-Non-Specific Salient	0.89(0.05)	***	<4	74%	95%	85%	93%	80%
Silly Sentences-Non-Specific-Non-Salient	0.94(0.03)	***	<4	84%	95%	90%	94%	87%
Silly Sentences-Specific-Total	0.85(0.06)	***	<10	74%	95%	85%	93%	80%
Silly Sentences-Specific-Salient	0.76(0.08)	*	<5	53%	100%	78%	100%	70%
Silly Sentences-Specific-Non-Salient	0.82(0.07)	***	<5	68%	95%	83%	93%	77%

Note. ROC = receiver operating characteristics curve; AUC = area under the curve; *SE* = standard error; OA = overall accuracy; PPV = positive predictive value; NPV = negative predictive value; NC = normal controls; MCI = mild cognitive impairment; CERAD = Consortium to Establish a Registry for Alzheimer's Disease; RBMT (PRT) = Rivermead Behavioural Memory Test-Paragraph Recall Test.

* Significant value ($p < .01$); ** Significant value ($p < .001$); ***Significant value ($p < .0001$).

DISCUSSION

The first aim of this study was to investigate the relative discriminatory efficacy of a newly developed PM task compared with two traditionally-used RM tests. The sensitivity, specificity, positive and negative predictive indices derived from ROC analyses showed that an event-based PM task (i.e., Silly Sentences-Non-Specific), was superior to traditionally used RM tests (RBMT-PRT and CERAD-Word List) in discriminating MCI-AD from NC. This is consistent with the idea that PM tasks demand a greater degree of self-initiative retrieval operations relative to RM tasks (Einstein & McDaniel, 1996), requiring the monitoring of the environment to identify the cue that signals the initiation of the action. This finding is most noteworthy since the CERAD-Delay Recall (Welsh et al., 1994), and RBMT have been found to discriminate very mild AD from NC, and MCI-AD from NC (Kazui et al., 2005).

The second aim of the study was to investigate whether varying parameters of the event-based PM task (i.e., specificity of the instructions and perceptual salience of the PM cue) would result in differential levels of discrimination between the groups. According to the multiprocess view of prospective remembering, the cognitive processes recruited in PM tasks, and the likelihood of success, are determined by the nature and demands of the ongoing task, and parameters of the PM cue (McDaniel & Einstein, 2000). This theory posits that ongoing tasks that encourage focal processing of the PM cue, and cues that are salient relative to the existing context in which they are presented, are more likely to produce an involuntary orienting response to the PM cue, and stimulate a relatively spontaneous retrieval of the intended action. In contrast, non-focal tasks, using non-salient PM cues, are likely to require more strategic attentional resources to monitor for the PM cue signaling the appropriateness of

performing the intended action (McDaniel & Einstein, 2000; McDaniel et al., 2008; McDaniel & Einstein, 2007). Consistent with the multiprocess theory, we found that the Silly-Sentences-Non-Specific-Non-Salient condition was superior to the Silly-Sentences-Specific-Salient in discriminating MCI-AD from NC. This differential pattern is consistent with the early presence of defective executive functioning in MCI-AD (Grober et al., 2008).

This study contributes to the limited literature on PM in MCI. Our findings are consistent with the two previous studies (Kazui et al., 2005; Troyer & Murphy, 2007), indicating that event-based PM tasks are sensitive to the cognitive effects of early MCI-AD. In addition, we have shown that a certain type of event-based PM task, the Silly Sentences-Non-Specific-Non-Salient, achieved a greater discriminatory power relative to traditional RM tests. This finding has research and clinical implications. First, paraphrasing McDaniel's & Einstein (2000) idea applied to the study of PM in MCI, we should move from investigating whether MCI affects prospective remembering to studying the types of PM tasks that provide a robust cognitive marker in MCI, aiding early detection. Second, future research will be needed to develop shorter PM tests appropriate for use in clinical settings.

Limitations of this study include, small sample size, use of convenience samples (MCI-AD were a selected group attending a memory clinic, and the NC were healthy Active Retirement Group participants), and, albeit not statistically significant, differences on educational profiles, all reducing the generalizability of our results. Present findings are regarded as preliminary results awaiting replication in larger unselected samples. Despite these limitations, the findings indicate that PM deficits are relatively more pronounced compared with RM failure in MCI-AD and that the Silly Sentences PM task may prove useful in the early detection of this group.

ACKNOWLEDGMENTS

The authors thank the participants and their families who cooperated in this study. The manuscript is new and original, is not currently under review by any other publication, and has not been previously published either electronically or in print. The project did not receive any financial support, and there are no conflicts of interest affecting this manuscript.

REFERENCES

- American Psychiatric Association. (2000). *Diagnostic and Statistical Manual of Mental Disorders* (4th ed. text revision). Washington, DC: American Psychiatric Association Press.
- Aron, A., Aron, E.N., & Coups, E. (2006). *Statistics for Psychology*. London: Prentice Hall.
- Baddeley, A., Emslie, H., & Nimmo-Smith, I. (1992). *The Speed and Capacity of Language Processing (SCOLP) Test*. Bury St Edmunds: Thames Valley Test Company.
- Benton, A.L. & Hamsler, K. (1978). *Multilingual Aphasia Examination manual*. Iowa City: University of Iowa.
- Brandimonte, M.A. & Pasolunghi, M.C. (1994). The Effect of cue-familiarity, cue-distinctiveness, and retention interval on prospective remembering. *Quarterly Journal of Experimental Psychology: Human Experimental Psychology*, *47*, 565–588.
- Brown, W.P. & Semrau, G. (1986). The rated typicality of items in semantic categories: Some Irish norms. *Irish Journal of Psychology*, *2*, 123–146.
- Coen, R.F., Kirby, M., Swanwick, G.R., Maguire, C.P., Walsh, J.B., Coakley, D., O'Neill, D., & Lawlor, B.A. (1997). Distinguishing between patients with depression or very mild Alzheimer's disease using the Delayed-Word-Recall test. *Dementia and Geriatric Cognitive Disorders*, *8*, 244–247.
- Driscoll, I., McDaniel, M.A., & Guynn, M.J. (2005). Apolipoprotein E and prospective memory in normally aging adults. *Neuropsychology*, *19*, 28–34.
- Einstein, G.O. & McDaniel, M.A. (1996). Retrieval processes in prospective memory: Theoretical approaches and some new empirical findings. In M. Brandimonte, G.O. Einstein, & M.A. McDaniel (Eds.), *Prospective Memory: Theory and applications* (pp. 115–142). Mahwah, NJ: Lawrence Erlbaum Associates.
- Folstein, M.F., Folstein, S.E., & McHugh, P.R. (1975). "Mini-mental State": A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, *12*, 189–198.
- Grober, E., Hall, C.B., Lipton, R.B., Zonderman, A.B., Resnick, S.M., & Kawas, C. (2008). Memory impairment, executive dysfunction, and intellectual decline in preclinical Alzheimer's disease. *Journal of the International Neuropsychological Society: JINS*, *14*, 266–278.
- Henry, J.D., MacLeod, M.S., Phillips, L.H., & Crawford, J.R. (2004). A meta-analysis review of prospective memory in aging. *Psychology and Aging*, *19*, 27–39.
- Huppert, F.A., Brayne, C., Gill, C., Paykel, E.S., & Beardsall, L. (1995). CAMCOG- a concise neuropsychological test to assist dementia diagnosis: Sociodemographic determinants in an elderly population sample. *British Journal of Clinical Psychology*, *34*, 529–541.
- Kazui, H., Matsuda, A., Hirono, N., Mori, E., Miyoshi, N., Ogino, A., Tokunaga, H., Ikejiri, Y., & Takeda, M. (2005). Everyday memory impairment of patients with Mild Cognitive Impairment. *Dementia and Geriatric Cognitive Disorders*, *19*, 331–337.
- Kluger, A., Golomb, J., & Ferris, S.H. (2000). Mild cognitive impairment. In N. Qizilbash (Ed.), *Evidence-based Dementia Practice* (pp. 341–354). Oxford: Oxford University Press.
- Knopman, D.S. & Ryberg, S. (1989). A verbal memory test with high predictive accuracy for dementia of the Alzheimer type. *Archives of Neurology*, *46*, 1097–1105.
- Lawton, P.M. & Brody, E.M. (1969). Assessment of older people: Self-maintaining and instrumental activities of daily living. *Gerontologist*, *9*, 179–186.
- Manos, P.J. (1999). Ten-point clock test sensitivity for Alzheimer's disease in patients MMSE scores greater than 23. *International Journal of Geriatric Psychiatry*, *14*, 454–458.
- Monsch, A.U., Bondi, M.W., Butters, N., Salmon, D.P., Katzman, R., & Thal, L.J. (1992). Comparison of verbal fluency tasks in the detection of dementia of the Alzheimer's type. *Archives of Neurology*, *49*, 1253–1258.
- McDaniel, M.A. & Einstein, G.O. (2000). Strategic and automatic processes in prospective memory retrieval: A multiprocess framework. *Applied Cognitive Psychology*, *14*, S127–S144.
- McDaniel, M.A. & Einstein, G.O. (2007). Prospective memory: A new research enterprise. In M.A. McDaniel & G.O. Einstein (Eds.), *Prospective Memory: An Overview and Synthesis of an Emerging Field* (pp. 1–11). Thousand Oaks, CA: Sage Publications.
- McDaniel, M.A., Einstein, G.O., & Rendell, P.G. (2008). The puzzle of inconsistent age-related declines in prospective memory: A multiprocess explanation. In M. Kligel, M.A. McDaniel & G.O. Einstein (Eds.), *Prospective Memory: Cognitive, Neuroscience, Developmental, and Applied Perspectives* (pp. 141–158). New York: Lawrence Erlbaum Associates.
- Nelson, H.E. (1991). *National Adult Reading Test (NART). Test Manual (revised)*. Windsor: NFER-Nelson.
- Petersen, R.C., Doody, R., & Kuz, A. (2001). Current concepts in mild cognitive impairment. *Archives of Neurology*, *58*, 1985–1992.
- Portet, F., Ousset, P.J., Visser, P.J., Frisoni, G.B., Nobili, F., Scheltens, P., Vellas, B., & Touchon, J. (2006). Mild cognitive impairment (MCI) in medical practice: A critical review of the concept and new diagnostic procedure. Report of the MCI Working Group of the European Consortium on Alzheimer's Disease (EADC). *Journal of Neurology Neurosurgery and Psychiatry*, *77*, 714–718.
- Sheikh, J.A. & Yesavage, J.A. (1986). Geriatric Depression Scale (GDS): Recent findings and development of a shorter version. In T.L. Brink (Ed.), *Clinical Gerontology: A Guide to Assessment and Intervention* (pp. 165–173). New York, NY: Haworth Press.
- Swanwick, G.J., Coen, R.F., O'Mahony, D., Tully, M., Bruce, I., Buggy, F., Lawlor, B.A., Walsh, J.B., & Coakley, D. (1996). A memory clinic for the assessment of mild dementia. *Irish Medical Journal*, *89*, 104–105.
- Troyer, A.K. & Murphy, K.J. (2007). Memory for intentions in amnesic mild cognitive impairment: Time- and event-based prospective memory. *Journal of the International Neuropsychological Society: JINS*, *13*, 365–369.
- Welsh, K.A., Butters, N., Mohs, R.C., Beekly, D., Edland, S., Fillenbaum, G., & Heyman, G. (1994). The consortium to establish a registry for Alzheimer's disease (CERAD). Part V. A normative study of the neuropsychological battery. *Neurology*, *44*, 609–614.
- Wilson, B.A., Cockburn, J., & Baddeley, A.D. (1985). *The Rivermead Behavioural Memory Test*. Bury St Edmunds, UK: Thames Valley Test.