

Accepted Manuscript

Lanthanide luminescence sensing of copper and mercury ions using an imino-diacetate based Tb(III)-cyclen chemosensor

Brian K. McMahon, Thorfinnur Gunnlaugsson

PII: S0040-4039(10)01392-4
DOI: [10.1016/j.tetlet.2010.07.174](https://doi.org/10.1016/j.tetlet.2010.07.174)
Reference: TETL 38212

To appear in: *Tetrahedron Letters*

Received Date: 10 June 2010
Revised Date: 16 July 2010
Accepted Date: 30 July 2010



Please cite this article as: McMahon, B.K., Gunnlaugsson, T., Lanthanide luminescence sensing of copper and mercury ions using an iminodiacetate based Tb(III)-cyclen chemosensor, *Tetrahedron Letters* (2010), doi: [10.1016/j.tetlet.2010.07.174](https://doi.org/10.1016/j.tetlet.2010.07.174)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.


Graphical Abstract

To create your abstract, type over the instructions in the template box below.
Fonts or abstract dimensions should not be changed or altered.

Lanthanide luminescence sensing of copper and mercury ions using an iminodiacetate based Tb(III)-cyclen chemosensor

Brian K. McMahon, Thorfinnur Gunnlaugsson*

Leave this area blank for abstract info.





Pergamon

TETRAHEDRON
LETTERS

Lanthanide luminescence sensing of copper and mercury ions using an iminodiacetate based Tb(III)-cyclen chemosensor


Brian K. McMahon, Thorfinnur Gunnlaugsson*

School of Chemistry, Centre for Synthesis and Chemical Biology, Trinity College Dublin, Dublin 2, Ireland.

Abstract— The design, synthesis and photophysical evaluation of **1.Tb.Na**, a Tb(III)-cyclen based sensor, possessing a phenyl iminodiacetate based receptor, for the selective detection of Cu(II) and Hg(II) ions in water is demonstrated. Sensitisation of the Tb(III) 5D_4 excited state was achieved by excitation of the phenyl receptor, which in water gave rise to characteristic time-delayed and line-like Tb(III) emission. The Tb(III) emission was shown to be pH independent over the physiological pH window. The changes in the Tb(III) emission were monitored by carrying out metal titrations using various group I, II and transition metal ions. Of these, only the titrations of Cu(II) and Hg(II) gave rise to modulations in the Tb(III) emission; resulting in quenching in the Tb(III) emission by *ca.* 65% and 40%, respectively. © 2010 Elsevier Science. All rights reserved


One of the main research areas within the field of supramolecular chemistry is the development of sensors for use in competitive solution.¹⁻⁴ In particular, the detection and monitoring of biologically relevant ions *in vitro* and *in vivo* has become of critical importance using luminescent sensors.^{3,4} Due to shorter lived background emission (autofluorescence) and light scattering from such biological environments, the development of sensors possessing long-lived emission lifetimes has become an active area of research.⁵⁻⁷ The unique photophysical properties of the lanthanides,⁸⁻¹⁰ such as long wavelength emissions, sharp line-like bands and relatively long-lived excited state lifetimes [μ s for Yb(III) and Nd(III) to ms for Eu(III) and Tb(III)], makes them very attractive candidates for such application in *in vivo* sensing as well as for biological imaging, another topical area of research.^{5,10-12} Careful functionalisation of the macrocyclic framework, 1,4,7,10-tetraazacyclododecane (cyclen) has allowed for the formation of many examples of thermodynamically stable and kinetically inert complexes with Ln^{III} ions, which have been employed in luminescent sensing of metal ions.¹²⁻¹⁷ These possess a suitable antennae/receptor unit which can populate the lanthanide excited states *via* an energy sensitization process (*e.g.* the antenna effect); which are perturbed upon binding to ions, giving rise to changes in both the lanthanide emission intensities and lifetimes.^{15,16}

The design of sensors for detecting ions such as Cu(II) and Hg(II) in aqueous solution at low concentration levels is of current interest in both environmental and biomedical monitoring.^{13,18-20} Bound Cu(II) plays a vital role in many enzymatic processes, while free Cu(II) can disrupt and damage various biological processes, and is the cause of several conditions such as Wilson and Menkes diseases.^{21,22} Similarly, Hg(II) is toxic in humans and can accumulate in the blood-brain barrier which can result in severe neurological disorders.¹⁹ Hence, the design and development of novel sensors which show a strong and selective response at low concentration values for these ions is both an important and topical area of research.²³



Herein, we describe the development of **1.Tb.Na**, a Tb(III)-cyclen based chemosensor possessing an iminodiacetate antenna. The objective is to take advantage of the long-lived and long-wavelength emission properties of Tb(III) in

* Corresponding author. Tel.: + 353 1 896 3459; fax: + 353 1 671 2826; e-mail: gunnlaut@tcd.ie.



Scheme 1. Synthesis of **1.Tb.Na**.

conjunction with the use of a structurally simple and combined antenna/receptor moiety that could selectively recognize Cu(II) and Hg(II) ions in competitive media. While many examples of luminescent lanthanide sensors of group I and II metal ions, as well as several transition metal ions such as Cu(II), Zn(II), *etc.* exist,¹³⁻¹⁷ the sensing of Hg(II), has, to the best of our knowledge, not been achieved previously using lanthanide luminescence, and hence sensor **1.Tb.Na** is the first example of a lanthanide sensor for targeting this ion.

The synthesis of ligand **1** began with the formation of **2**, Scheme 1. This involved the initial di-alkylation of aniline with ethyl bromoacetate using Na_2HPO_4 and KI in refluxing CH_3CN for 24 h, which upon purification by flash silica gel column chromatography (gradient elution of 100 to 90:10 CH_2Cl_2 : CH_3OH), resulted in the isolation of the desired intermediate in 56% yield. Nitration at the *para* position was achieved in acetic acid, at 0 °C, using 70% nitric acid. The desired product was obtained in 93% yield after recrystallisation from EtOH.²⁴ The product was then subjected to catalytic hydrogenation using 10% Pd/C, in DMF, giving **2** in 90% yield. The formation of the α -chloroamide of **2**, was achieved using chloroacetyl chloride, in the presence of Et_3N , at -10 °C in CH_2Cl_2 , which after washing the organic layer with water, 0.1 M HCl and brine, gave **3**, in 86% yield. With the receptor/antenna unit successfully synthesised, the next step involved introduction to the cyclen framework using the mono alkylation procedure recently published by our group,²⁵ involving refluxing four equivalents of cyclen with one equivalent of **3**, in dry CHCl_3 , in the presence of Et_3N for 12 hours. Isolation of **4** was achieved in 90% yield after successful removal of the unreacted cyclen by extraction using 1 M NaOH. The final step involved refluxing **4** with three equivalents of dimethyl acetamide using K_2CO_3 and KI in freshly distilled CH_3CN for five days. Purification of the crude mixture by alumina column chromatography (elution gradient 100 to 80:20 CH_2Cl_2 : CH_3OH) was required to give the desired ligand, in 63% yield. The formation of the Tb(III) complex of ligand **1** was achieved by refluxing overnight with an equivalent amount of Tb(III) triflate $[\text{Tb}(\text{CF}_3\text{SO}_3)_3]$ in freshly distilled CH_3CN , under an inert atmosphere. Upon completion, the solvent was removed under reduced pressure and the resulting residue re-dissolved in MeOH and added slowly into a large volume of dry diethyl ether, resulting in precipitation of **1.Tb** in 80% yield, as a pale orange solid. The final step involved alkaline hydrolysis of the diethyl diester moiety of **1.Tb** in MeOH/ H_2O , which gave **1.Tb.Na** in 81% yield.²⁶

The design of **1.Tb.Na** envisaged sensitization of the Tb(III) metal ion *via* excitation of the covalently attached phenyl chromophore, *i.e.* the antenna (possessing both π - π^* and n- p^* transitions). This would enable sensitization of the Tb(III) $^5\text{D}_4$ excited state, with concomitant characteristic line-like Tb(III) emission; which would then be expected to be modulated upon binding of d-metal ions at the iminodiacetate moiety, possibly involving electron transfer quenching. Indeed, excitation of **1.Tb.Na** in H_2O at 285 nm (λ_{max} for the antenna) gave rise to Tb-centered emissions at 490, 545, 586 and 622 nm, respectively, verifying successful energy transfer from the antennae to the $^5\text{D}_4$ excited state, and deactivation to the $^7\text{F}_j$ ($J = 6, 5, 4, 3$) ground state. The excited state lifetimes were also measured in both H_2O and D_2O , upon excitation at 285 nm. This gave $\tau_{\text{H}_2\text{O}} = 1.469$ ms and $\tau_{\text{D}_2\text{O}} = 1.984$ ms, demonstrating the long excited state lifetime, from which the hydration state q was determined as ~ 1 indicating that **1.Tb.Na** possesses one metal bound water molecule, and therefore the Tb(III) ion being overall nine-coordinate within the complex.

It was clear from the structure of **1.Tb.Na** that possible protonation of the aniline moiety as well as potential deprotonation of the secondary amide (as well as the metal bound water) could perturb the photophysical properties of the Tb(III) complex.²⁷ In order to quantify the pH dependence of the Tb(III) emission from **1.Tb.Na**, we carried out a pH titration, the result of which is shown in Figure 1. Here, the Tb(III) emission corresponding to the $^5\text{D}_4 \rightarrow ^7\text{F}_j$ ($J = 6, 5, 4, 3$) transition occurring at 490 nm, 545 nm, 586 nm and 622 nm, respectively, showed some change with respect to pH environment, Figure 1. Analysis of the Tb(III) emission for the $\Delta J = 5$ transition (see inset in Figure 1) clearly demonstrates that the Tb(III) emission is




Figure 1. The overall changes in the Tb(III) luminescence of **1.Tb.Na** (10 μM) upon excitation of the phenyl antenna ($\lambda_{\text{ex}} = 285$ nm) as a function of pH ($I = 0.1$ M $\text{NEt}_4\text{HClO}_4$ (TEAP)). *Inset*: The changes at 545 nm vs. pH.

'switched on' within the physiological pH range, with any major modulations in emission intensity only occurring below pH 5.5, or above pH 11.5. Hence, there exists a large pH window in which no change in the Tb(III) emission of **1.Tb.Na** is observed, and that this window overlaps with the physiological pH range. Hence, we would anticipate that any modulation in the emission intensity observed within this pH range upon titration with various transition metal ions is solely due to some form of interaction with the complex, and not as a result of a change in the pH of the solution.^{8,15}

Having established the pH behaviour of **1.Tb.Na**, the next step was to investigate its ability to recognise various group I, group II and transition metal ions in solution. Screening a large library of these in water at pH 7, demonstrated that the Tb(III) emission was only affected by the presence of Cu(II) and Hg(II), using either the Cl⁻ or HClO₄⁻ salts of these ions. However, ions such as Co(II) and Ni(II) also gave rise to changes in the Tb(III) emission, but these were due to the appearance of an absorption band corresponding to these ions, and hence, an inner filter effect, and not the direct binding of these ions to the receptor of **1.Tb.Na**.²⁸ Moreover, it was observed that in the presence of either HEPES or TRIS buffers alone, both the Cu(II) and Hg(II) ions showed some form of complexation with the buffers resulting in the formation of a strong absorption band which overlapped with the absorption of the antenna in **1.Tb.Na**. Given the fact that the Tb(III) emission can be considered as pH independent within the physiological pH range, e.g. as demonstrated as an inset in Figure 1, all the titrations discussed herein were carried out in the absence of any buffer in deionised H₂O. Nevertheless, to ensure that there was no pH effect, the pHs of all the samples were recorded before and after carrying out these titrations; the pH was not found to change by more than 0.2 pH units.

The changes in the ground state properties of **1.Tb.Na** upon titration with Cu(II) ions (using the HClO₄⁻ salt) were first investigated, the results of which are shown in Figure 2. The changes observed in the spectra clearly indicates an interaction between the Cu(II) ions in solution and the iminodiacetate moiety of **1.Tb.Na**, where the absorption




Figure 2. The overall changes in the absorption spectra (uncorrected) of **1.Tb.Na** (10 μ M) upon titration with Cu(HClO₄)₂ in deionised H₂O. *Inset:* Changes observed at 285 nm as a function of equivalents of Cu(II) added

band centered at 285 nm was blue-shifted to 260 nm, with the formation of an isobestic point at ca. 266 nm. The titration profile of the absorbance changes at 285 nm versus the number of equivalents of Cu(II) added is shown as an inset in Figure 2, and suggests that the most significant interaction occurs between 0 \rightarrow 1 equivalents of Cu(II), with minor changes occurring between 1 \rightarrow 2 equivalents (0 \rightarrow 20 μ M). A similar response was observed on analysis of the singlet excited state of **1.Tb.Na**, where, in the absence of Cu(II), a broad emission band was observed, centered at 350 nm with a slight shoulder at ca 300 nm. Upon binding to Cu(II), a moderate emission intensity increase of ca. 15% was observed (with a slight hypsochromic shift to 345 nm) being most significant within the addition of 0 \rightarrow 2 equivalents of Cu(II), as seen above in the absorption spectra.

We next monitored changes in the Tb(III) emission upon Cu(II) binding to the receptor which was expected to effect the energy transfer process from the antenna to the lanthanide, and hence, the Tb(III) emission. The luminescence spectrum of **1.Tb.Na**, upon titration with Cu(II) is shown in Figure 3, and clearly demonstrates a gradual decrease in the Tb(III) emission as a function of




Figure 3. Changes in the Tb(III) luminescence spectra of **1.Tb.Na** (10 μ M) upon titration with Cu(HClO₄)₂ (being quenched) in deionised H₂O. (0 \rightarrow 100 μ M) *Inset:* Changes observed at 454 nm as a function of equivalents of Cu(II) added.

increased Cu(II) concentration with ca. 65% quenching being observed at 545 nm, with a similar magnitude of quenching being seen for the other transitions. The Tb(III) excited state lifetime was also shortened upon binding to Cu(II), with $\tau_{\text{H}_2\text{O}} = 1.180$ ms. A plot of the Tb(III) intensity at 545 nm vs. Cu(II) equivalents is shown as an inset in Figure 3, and demonstrates the ability of **1.Tb.Na** to bind Cu(II), with concomitant changes in the emission properties. Analysis of the changes in Figure 3 also showed that the sensor bound the ion in a 1:1 stoichiometry, Figure 4, which demonstrates that the binding occurs over two log units, with an estimated binding constant²⁹ of logK = 5.5 (± 0.2). Figure 4, also shows the response observed for ions such as Zn(II) and Cd(II) as well as the group II ions Mg(II) and Ca(II), which did not give rise to any significant changes in the Tb(III) emission. This also demonstrates the importance of our receptor design, as structurally related sensors have been developed by Parker *et al.*¹⁶ for the




Figure 4. Plot of the changes in the Tb(III) luminescence of **1.Tb.Na** (10 μM) at 545 nm upon titration with $\text{M}(\text{HClO}_4)_2$ or MCl_2 (expressed as $\text{pM} = -\log[\text{M}]$) in deionised H_2O .

selective sensing of Zn(II); which is not detected by **1.Tb.Na**. With the free Cu(II) concentrations in serum being in the range of 12.5 – 21 μM ,³⁰ it is clear from the titration profile in Figure 4, that **1.Tb.Na** can detect Cu(II) within this concentration window. Figure 4, also shows the changes observed in the Tb(III) emission of **1.Tb.Na**, upon titration with Hg(II), the overall spectral changes being shown in Figure 5. A similar response to that seen with




Figure 5. Changes in the Tb(III) luminescence spectra of **1.Tb.Na** (10 μM) upon titration with $\text{Hg}(\text{HClO}_4)_2$ (being quenched) in deionised H_2O (0 \rightarrow 100 μM).

Cu(II) was observed, *i.e.* only the intensity and lifetime were affected, where upon the addition of 10 equivalents (100 μM) of Hg(II) gave rise to *ca* 40% luminescence quenching in the Tb(III) emission, with $\tau_{\text{H}_2\text{O}} = 1.302$ ms. From these changes, we estimated the Hg(II) binding affinity of **1.Tb.Na** as $\log K = 4.6 (\pm 0.2)$. These studies suggest that **1.Tb.Na** shows both high sensitivity and selectivity for both Cu(II) and Hg(II) ions; which was achieved through binding at the iminodiacetate receptor moiety in solution. These results also demonstrate that the sensor is more sensitive to Cu(II) over Hg(II) (*c.f.* Figure 4). To confirm this, we also carried out a test where firstly, an excess amount of Hg(II) was added to a solution of the sensor. This again, gave rise to quenching in the Tb(III) emission to the same degree as seen for the Hg(II) titrations. However, subsequent addition of Cu(II) lead to further quenching in the Tb(III) emission, clearly demonstrating this selectivity. We also investigated the response of **1.Tb.Na** towards Cu(II) and Hg(II) in a

simulated biological environment with 150 mM NaCl, 10 mM KCl, 10 mM CaCl_2 , 10 mM MgCl_2 , 10 mM CdCl_2 and 10 mM ZnCl_2 , at pH 7. As expected, no significant modulation in the singlet excited state properties of the Tb(III) metal was observed in the presence of this mimicked biological background, but addition of both Cu(II) and Hg(II) to the solution resulted in quenching of the Tb(III) metal luminescence, to the same extent as seen in the above titrations.

In summary, we have developed a Tb(III) cyclen-based chemosensor, with the incorporation of an iminodiacetate receptor/antenna unit, for Cu(II) and Hg(II) ions in water at pH 7. While only minor changes were observed in both the absorption and fluorescence spectra of **1.Tb.Na**, upon recognition of these ions, a substantial quenching was observed in the Tb(III) emission. These studies showed that the sensor has high affinity for both of these ions, where Cu(II) is selectively bound. It was also shown that in the presence of other biologically relevant ions, the emission properties of the Tb(III) metal were not modulated, confirming the **1.Tb.Na** selectivity for Cu(II) and Hg(II). We are in the process of developing analogues of **1.Tb.Na**, using other antennae, which will enable us to achieve sensitisation of Tb(III) at longer wavelengths and at the same time preserve the binding ability of the receptor unit, already used for **1.Tb.Na**.

Acknowledgements

We thank TCD, Irish Research Council for Science, Engineering and Technology (IRCSET Postgraduate Award to BKMCM), and Science Foundation Ireland (SFI RFP 2008 and 2009 research grants to TG), for financial support.

Notes and References

- Ryan, G. J.; Quinn, S.; Gunnlaugsson, T. *Inorg. Chem.* **2008**, *47*, 401; Zeglis, B. M.; Pierre, V. C.; Barton, J. K. *Chem. Commun.* **2007**, 4565; Moucheron, C.; Kirsch-De-Mesmaeker, A.; Kelly, J. M. *J. Photochem. Photobiol. B* **1997**, *40*, 91; Hannon, M. J. *Chem. Soc. Rev.* **2007**, *36*, 280; Metcalfe, C.; Thomas, J. A.; *Chem. Soc. Rev.* **2003**, *32*, 215; de Silva, A. P.; McCaughan, B.; McKinney, B. O. F.; Querol, M. *Dalton Trans.* **2003**, 1902; De Silva, A. P.; Fox, D. B.; Huxley, A. J. M.; Moody, T. S. *Coord. Chem. Rev.* **2000**, *205*, 41.
- Filbey, M. H.; Steed, J. W. *Coord. Chem. Rev.* **2006**, *250*, 3200; Redshaw, C. *Coord. Chem. Rev.* **2003**, *244*, 45; Gunnlaugsson, T.; Glynn, M.; Tocci, G. M.; Kruger, P. E.; Pfeffer, F. M. *Coord. Chem. Rev.* **2006**, *250*, 3094; Quinlan, E.; Matthews, S. E.; Gunnlaugsson, T. *J. Org. Chem.* **2007**, *72*, 7497; Rurack, K.; Resch-Genger, U. *Chem. Soc. Rev.* **2002**, *31*, 116.
- de Silva, A. P.; Gunaratne, H. Q. N.; Gunnlaugsson, T.; Huxley, A. J. M.; McCoy, C. P.; Rademacher, J. T.; Rice, T. E. *Chem. Rev.* **1997**, *97*, 1515; Czarnik, A. W. *Acc. Chem. Res.* **1994**, *27*, 302; *Fluorescent Chemosensors for Ion and Molecular Recognition*; Czarnik, A. W., Ed.; ACS Books: Washington, **1993**. Bissell, R. A.; de Silva, A. P.; Gunaratne, H. Q. N.; Lynch, P. L. M.; Maguire, G. E. M.; McCoy, C. P.; Sandanayake, K. R. A. S. *Top. Curr. Chem.* **1993**, *168*, 223.

4. Dai Z.; Canary, J. W. *New J. Chem.* **2007**, *31*, 1708; Gunnlaugsson, T.; Lee, T. C.; Parkesh, R. *Tetrahedron* **2004**, *60*, 11239; Gunnlaugsson, T.; Nieuwenhuyzen, M.; Richard, L.; Thoss, V. *Tetrahedron Lett.* **2001**, *42*, 4725; Balzani, V. *Photochem. Photobiol. Sci.* **2003**, *2*, 459; Balzani, V.; Credi, A.; Venturi, M. *Pure Appl. Chem.* **2003**, *75*, 541; Raymo, F. M. *Adv. Mater.* **2002**, *14*, 401; Raymo, F. M.; Giordani, S. *J. Am. Chem. Soc.* **2004**, *124*, 124; Gunnlaugsson, T.; Leonard, J. P.; Murray, N. S. *Org. Lett.*, **2004**, *6*, 1557; Brown, G. J.; de Silva, A. P.; Pagliari, S. *Chem. Commun.* **2002**, 2461; Ballardini, R.; Balzani, V.; Credi, A.; Gandolf, M. T.; Venturi, M. *Acc. Chem. Res.* **2001**, *36*, 445; Czarnik, A. W. *Fluorescent Chemosensors of Ion and Molecular Recognition*, ACS, Symp. Ser. American Chemical Society: Washington DC, **1993**.
5. dos Santos, C. M. G.; Harte, A. J.; Quinn, S.; Gunnlaugsson, T. *Coord. Chem. Rev.* **2008**, *252*, 2512; Leonard, J. P.; Nolan, C. B.; Stomeo, F.; Gunnlaugsson, T. *Top. Curr. Chem.* **2007**, *281*, 1; Gunnlaugsson, T.; Stomeo, F. *Org. Biomol. Chem.* **2007**, *5*, 1999; Gunnlaugsson, T.; Leonard, J. P. *Chem. Commun.* **2005**, 3114; Leonard, J. P.; Gunnlaugsson, T. *J. Fluoresc.* **2005**, *15*, 585; Comby, S.; Bünzli, J.-C. G. In *Handbook of the Chemistry and Physics of Rare Earths*, Gschneider, K. A.; Bünzli, J.-C. G.; Pecharsky, V. K.; Ed. Elsevier, Amsterdam, **2007**, *37*; Murray, B. S.; New, E. J.; Pal, R.; Parker, D. *Org. Biomol. Chem.* **2008**, *6*, 2085; Liellar, F.; Law, G.-L.; New, E. J.; Parker, D. *Org. Biomol. Chem.* **2008**, *6*, 2256; Chauvin, A.-S.; Comby, S.; Song, B.; Vandevyver, C. D. B.; Thomas, F.; Bünzli, J.-C. G. *Chem. Eur. J.* **2007**, *13*, 9515; Gunnlaugsson, T.; Harte, A. J.; Leonard, J. P.; Nieuwenhuyzen, M. *Supramol. Chem.* **2003**, *15*, 505.
6. Cantuel, M.; Linceneau, C.; Buffeteau, T.; Jonusauskaite, L.; Gunnlaugsson, T.; Jonusauskas, G.; McClenaghan, N. D. *Chem. Commun.*, **2010**, *46*, 2486; Murray, N. S.; Jarvis, S. P.; Gunnlaugsson, T. *Chem. Commun.* **2009**, 4959; Bonnet, C. S.; Massue, J.; Quinn, S. J.; Gunnlaugsson, T. *Org. Biomol. Chem.* **2009**, *7*, 3074; Plush, S. E.; Gunnlaugsson, T. *Dalton Trans.* **2008**, 3801; Massue, J.; Quinn, S. J.; Gunnlaugsson, T. *J. Am. Chem. Soc.* **2008**, *130*, 6900; Plush, S. E.; Gunnlaugsson, T. *Org. Lett.* **2007**, *9*, 1919; dos Santos, C. M. G.; Fernandez, P. B.; Plush, S. E.; Leonard, J. P.; Gunnlaugsson, T. *Chem. Commun.* **2007**, 3389; Harte, A. J.; Jensen, P.; Plush, S. E.; Gunnlaugsson, T. *Inorg. Chem.* **2006**, *45*, 9465;
7. Sénéchal-David, K.; Pope, S. J. A.; Quinn, S.; Faulkner, S.; Gunnlaugsson, T. *Inorg. Chem.* **2006**, *45*, 10040.
8. McCoy, C. P.; Stomeo, F.; Plush, S. E.; Gunnlaugsson, T. *Chem. Mater.* **2006**, *18*, 4336; Gunnlaugsson, T.; McCoy, C. P.; Stomeo, F. *Tetrahedron Lett.* **2004**, *45*, 8403
9. Bünzli, J.-C. G.; Piquet, C. *Chem. Soc. Rev.* **2005**, *34*, 1048; Nonat, A.; Gateau, C.; Fries, P. H.; Mazzanti, M. *Chem. Eur. J.* **2006**, *12*, 7133.
10. Bünzli, J.-C. G. *Chem. Lett.* **2009**, *38*, 104; Song, B.; Vandevyver, D. B.; Deiters, E.; Chauvin, A.-S.; Hemmila, I.; Bünzli, J.-C. G. *Analyst* **2008**, *133*, 1749; Pandya, S.; Yu, J.; Parker, D. *Dalton Trans.* **2006**, 2757; Yu, J. J.; Parker, D.; Pal, R.; Poole, R. A.; Cann, M. J. *J. Am. Chem. Soc.* **2006**, *128*, 2294.
11. McMahon, B.; Mauer, P.; McCoy, C. P.; Lee, T. C.; Gunnlaugsson, T. *J. Am. Chem. Soc.* **2009**, *131*, 17542.
12. Nonat A. M.; Harte, A. J.; Sénéchal-David, K.; Leonard, J. P.; Gunnlaugsson, T. *Dalton Trans.* **2009**, 4703.
13. Andrews, M.; Jones, J. E.; Harding, L. P.; Pope S. J. A. *Chem. Commun.*, **2010**, *46*, DOI: 10.1039/c0cc00210k; Pope, S. J. A.; Laye, R. H. *Dalton Trans.* **2006**, 3108; Hanaoka, K.; Kikuchi, K.; Kojima, H.; Urano, Y.; Nagana, T. *Angew. Chem. Int. Ed.* **2003**, *42*, 2996; Hanaoka, K.; Kikuchi, K.; Kojima, H.; Urano, Y.; Nagana, T. *J. Am. Chem. Soc.* **2004**, *126*, 12470.
14. Armelao, L.; Bottaro, G.; Quici, S.; Cavazzini, M.; Raffo, M. C.; Barigelletti, F.; Accorsi, G. *Chem. Commun.* **2007**, 2911; Li, C.; Law, G. L.; Wong, W. T. *Org. Lett.* **2004**, *6*, 4841; Li, C.; Wong, W. T. *Tetrahedron Lett.* **2004**, *45*, 6055; de Silva, A. P.; Gunaratne, H. Q. N.; Rice, T. E. *Angew. Chem. Int. Ed. Engl.* **1996**, *35*, 2116; de Silva, A. P.; Gunaratne, H. Q. N.; Rice, T. E. *Chem. Commun.* **1997**, 1891; Gunnlaugsson, T.; Parker, D. *Chem. Commun.* **1998**, 511.
15. Gunnlaugsson, T.; Leonard, J. P. *Chem. Commun.* **2003**, 2424; Gunnlaugsson, T.; Leonard, J. P. *Dalton Trans.* **2005**, 3204; Coldwell, J. B.; Felton, C. E.; Harding, L. P.; Moon, R.; Pope, S. J. A.; Rice, C. R. *Chem. Commun.* **2006**, 5048.
16. Reany, O.; Gunnlaugsson, T.; Parker, D. *Chem. Commun.* **2000**, 473. Reany, O.; Gunnlaugsson, T.; Parker, D. *J. Chem. Soc., Perkin Trans 2*, **2000**, 1819.
17. Voutsadaki, S.; Tsikalas, G. K.; Klontzas, E.; Froudakis, G. E.; Katerinopoulos, H. E. *Chem. Commun.* **2010**, *46*, 3292; Li, N.; Xiang, Y.; Tong, A. *Chem. Commun.* **2010**, *46*, 3363; Hanaoka, K.; Kikuchi, K.; Kojima, H.; Urano, Y.; Nagano, T. *J. Am. Chem. Soc.* **2004**, *126*, 12470.
18. Selid, P. D.; Xu, H.; Collins, E. M.; Collins, M.; Zhao, J. X. *Sensors* **2009**, *9*, 5446; Torrado, A.; Walkup, G. K.; Imperiali, B. *J. Am. Chem. Soc.* **1998**, *120*, 609; Beltramello, M.; Gatos, M.; Mancin, F.; Tecilla, P.; Tonellato, U. *Tetrahedron Lett.* **2001**, *42*, 9143; Yoon, J.; Ohler, N. E.; Vance, D. H.; Aumiller, W. D.; Tonellato, U. *Tetrahedron Lett.* **1997**, *38*, 3845.
19. Barja, B. C.; Aramendia, P. F. *Photochem. Photobiol. Sci.* **2008**, *7*, 1391; Liu, Y.; Yu, M.; Chen, Y.; Zhang, N. *Bioorg. Med. Chem.* **2009**, *17*, 3887; Kwon, J. Y.; Hyun Soh, J.; Yoon, Y. J.; Yoon, J. *Supramol. Chem.* **2004**, *16*, 621; Tian, M.; Ihmels, H. *Chem. Commun.* **2009**, 3175; Wanichacheva, N.; Sirirupmoonthum, M.; Kamkaew, A.; Grudpan, K. *Tetrahedron Lett.* **2009**, *50*, 1783; Hasegawa, Y.; Hieda, R.; Nakagawa, T.; Kawai, T. *Helv. Chim. Acta.* **2009**, *92*, 2238.
20. Sarkar, B. *Chem. Rev.* **1999**, *99*, 2535.
21. Lindar, M. C.; Hazegh-Azam, M. *Am. J. Clin. Nutr.* **1996**, *63*, 8218.
22. Sarker, B. *Handbook on Metals in Clinical and Analytical Chemistry*, Marcel Dekker Inc. New York, **1994**.
23. Nolan, E. M.; Lippard, S. J. *Chem. Rev.* **2008**, *108*, 3443; Nolan, E. M.; Lippard, S. J. *J. Am. Chem. Soc.* **2007**, *129*, 5910.
24. Parkesh, R.; Gowin W.; Lee, T. C.; Gunnlaugsson, T. *Org. Biomol. Chem.* **2006**, *4*, 3611.
25. Massue, J.; Plush, S. E.; Bonnet, C. S.; Moore, D. A.; Gunnlaugsson T. *Tetrahedron Lett.* **2007**, *48*, 805
26. Ligand **1** was obtained as an orange crystalline solid. (0.5 g, 63% yield). m.p. 189-192 °C; HRMS (*m/z*, ES⁺) Calculated for C₃₆H₆₁N₉O₈Na *m/z* = 770.4541 [M + Na]. Found *m/z* = 770.4549; ¹H NMR (400 MHz, CDCl₃, δ_H): 10.40 (s, 1H, N-H), 7.70 (d, 2 H, *J* = 9 Hz, Ar-H), 6.47 (d, 2 H, *J* = 9 Hz, Ar-H), 4.17 (q, 4 H, *J* = 7 Hz, NCH₂CO₂CH₂CH₃), 4.08 (s, 4H, NCH₂CO₂CH₂CH₃), 3.30-1.98 (br m, 42H, cyclen-CH₂ + 4.7,10-CH₂CON(CH₃)₂ + 1-CH₂CONH), 1.26 (t, 6 H, *J* = 7 Hz, NCH₂CO₂CH₂CH₃); ¹³C NMR (100 MHz, CDCl₃, δ_C): 170.58 (q), 170.52 (q), 170.24 (q), 169.58 (q), 143.48 (CH), 130.65 (CH), 120.72 (CH₂), 112.06 (CH₂), 60.59 (CH₂), 57.44 (CH₂), 54.42 (CH₂), 53.36 (CH₂), 53.03 (CH₂), 50.13 (CH₂), 35.96 (CH₃), 35.71 (CH₃), 35.22 (CH₃), 35.12 (CH₃), 13.80 (CH₃); IR ν_{max} (cm⁻¹): 2818, 1740, 1646, 1518, 1450, 1400, 1370, 1346, 1297, 1262, 1179, 1102, 1062, 1005, 974, 902, 818, 772, 730, 631.
- Complex **1.Tb** was obtained as a pale orange solid (0.06 g, 80% yield). m.p. decomposed above 200 °C; Calculated for C₄₂H₈₄Cl₆F₉N₉O₁₇S₃Tb: C, 29.17; H, 3.85; N, 7.29. Found C, 29.19; H, 3.85; N, 7.47; HRMS (*m/z*) Calculated for C₃₈H₆₁F₆N₉O₁₄S₂Tb *m/z* = 1204.2937 [M + 2(CF₃SO₃)]. Found *m/z* = 1204.2952; ¹H NMR (400 MHz, CDCl₃, δ_H): 72.47, 61.18, 50.78, 49.44, 46.20, 42.74, 36.48, 35.48, 34.58, 23.40, 19.48, 17.81, 13.79, 7.64, 6.75, 4.25, 3.79, 3.06, 2.76, 2.43, 1.43, 1.34, 0.93. IR ν_{max} (cm⁻¹): 3458, 2941, 1735, 1618, 1560, 1520, 1459, 1411, 1246, 1224, 1158, 1081, 1028, 958, 910, 823, 759, 635.
- Complex **1.Tb.Na** was obtained as a pale yellow solid (0.080 g, 81% yield). m.p. decomposed above 250 °C; HRMS (*m/z*, ES⁺) Calculated for C₃₄H₅₃N₉O₁₄S₂F₆Tb *m/z* = 1148.2311 [M - CF₃SO₃ - 2Na + 2H]⁺. Found *m/z* = 1148.2357; ¹H NMR (400 MHz, D₂O, δ_H): 85.92, 79.51, 69.02, 67.04, 61.52, 58.42, 54.56, 53.79, 52.79, 49.04, 44.07, 44.38, 22.96, 21.53, 20.53, 16.67, 15.77, 14.20, 11.41, 10.37, 8.35, 7.84, 7.19, 6.72, 6.59, 6.43, 6.34, 3.52, 1.30, 1.17, -0.06, -82.05, -85.69, -88.45, -97.73, -101.59, -103.47; IR ν_{max} (cm⁻¹): 2972, 1603, 1438, 1251, 1229, 1168, 1088, 1035, 945, 906, 878, 864, 765, 687, 638.
27. Plush, S. E.; Clear, N. A.; Leonard, J. P.; Fanning, A. M. Gunnlaugsson, T. *Dalton Trans.* **2010**, *39*, 3644.
28. This was verified by carrying out titrations using these ions in the absence of the sensor, which on both occasions, gave rise to large

absorptions with a λ_{\max} of 400 and 505, which partially overlapped with the emission wavelengths of Tb(III).

- 29 We were unable to fit the changes in the Tb(III) accurately using non-linear regression analysis
- 30 Pešák, J.; Opavský, J. *Acta Univ. Palacki. Olomuc., Fac. Med.* **2000**, *143*, 71.

ACCEPTED MANUSCRIPT