


Accepted Manuscript

Title: fMRI activation during response inhibition and error processing: the role of the DAT1 gene in typically developing adolescents and those diagnosed with ADHD.

Authors: Wouter Braet, Katherine A. Johnson, Claire T. Tobin, Ruth Acheson, Caroline McDonnell, Ziarah Hawi, Edwina Barry, Aisling Mulligan, Michael Gill, Mark A. Bellgrove, Ian H. Robertson, Hugh Garavan



PII: S0028-3932(11)00003-0
DOI: doi:10.1016/j.neuropsychologia.2011.01.001
Reference: NSY 3951

To appear in: *Neuropsychologia*

Received date: 26-10-2009
Revised date: 3-12-2010
Accepted date: 1-1-2011

Please cite this article as: Braet, W., Johnson, K. A., Tobin, C. T., Acheson, R., McDonnell, C., Hawi, Z., Barry, E., Mulligan, A., Gill, M., Bellgrove, M. A., Robertson, I. H., & Garavan, H., fMRI activation during response inhibition and error processing: the role of the DAT1 gene in typically developing adolescents and those diagnosed with ADHD., *Neuropsychologia* (2008), doi:10.1016/j.neuropsychologia.2011.01.001

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The DAT1 10R allele was associated with more effortful response inhibition

However, this group showed a reduced response to errors, suggesting less learning

The ADHD group displayed underactivation in parietal, medial and frontal regions

The effects of DAT1 genotype were more pronounced in the ADHD group

Accepted Manuscript

1
2
3
4 **fMRI activation during response inhibition and error processing: the role of the**
5
6 **DAT1 gene in typically developing adolescents and those diagnosed with ADHD.**
7
8
9

10
11 Wouter Braet^{1,2,*}, Katherine A. Johnson¹, Claire T. Tobin¹, Ruth Acheson¹, Caroline
12 McDonnell³, Ziarah Hawi³, Edwina Barry³, Aisling Mulligan³, Michael Gill³, Mark A.
13 Bellgrove⁴, Ian H. Robertson¹, Hugh Garavan¹
14
15
16
17
18

19 *¹School of Psychology and Trinity College Institute of Neuroscience, Trinity College Dublin, Dublin 2,*
20 *Ireland; ²Laboratory of Experimental Psychology, University of Leuven, Belgium; ³ School of Medicine,*
21 *Trinity College Dublin; ⁴ The University of Queensland, School of Psychology and Queensland Brain*
22 *Institute, Australia.*
23
24
25
26

27 ***corresponding author: Wouter.braet@gmail.com**
28
29
30
31

32 Running title: DAT1 effects on response inhibition in ADHD
33
34
35
36
37
38
39
40
41
42

43 Corresponding author:
44
45 Wouter Braet
46 Laboratory of Experimental Psychology
47 University of Leuven
48 Tiensestraat 102
49 B 3000 Leuven
50 Belgium
51 Tel: +32 (0)16 32 59 57
52 Fax: +32 (0)16 32 60 99
53 Email: wouter.braet@gmail.com
54
55
56
57
58
59
60
61
62
63
64
65

Abstract

The DAT1 gene codes for the dopamine transporter, which clears dopamine from the synaptic cleft, and a variant of this gene has previously been associated with compromised response inhibition in both healthy and clinical populations. This variant has also been associated with ADHD, a disorder that is characterised by disturbed dopamine function as well as problems with response inhibition. In the present study we used fMRI to investigate the role of dopaminergic genetic variation on executive functioning by comparing how activation associated with successful and unsuccessful inhibitions differs based on DAT1-genotype and ADHD-diagnosis in adolescents performing a go/nogo task. The results identify regional specificity concerning which functional differences can be attributed to the possession of the high risk DAT1 genotype, the clinical condition or an interaction between the two. During response inhibition, individuals with two copies of the 10-repeat allele showed increased activation in frontal, medial, and parietal regions, which may indicate that inhibition is more effortful for this group. Conversely, this group displayed a reduced error response in the parahippocampal gyrus, suggestive of reduced learning from errors. There were also a number of frontal, parietal, medial and occipital regions, where the relationship between genotype and fMRI-activation differed between the ADHD group and typically developing adolescents. Finally, the ADHD group displayed decreased activation in parietal and (pre)frontal regions during response inhibition, and in frontal and medial brain regions on error trials.

Keywords: response inhibition, fMRI, endophenotype, DAT1, dopamine, error processing, ADHD

Introduction

Controlling and inhibiting actions are important cognitive abilities that allow us to function successfully in a changing and complex environment. Individual differences in executive functioning such as response inhibition and error monitoring have been related to genetic variation (Goldberg & Weinberger, 2004), for example, by studying unaffected family members of clinical groups with deficits in response inhibition (Slaats-Willemse et al., 2003) and by comparing groups who differ in genotype. Earlier studies suggest an important role for the dopamine system in response inhibition: typically developing children homozygous for the 10-repeat allele of DAT1 perform less well on a test of response inhibition (Cornish et al., 2005), though this effect can be modulated by genotypic variation on other dopaminergic genes such as DRD4 (Congdon, Lesch & Canli, 2007). In this study we investigate the role of dopaminergic functioning in response inhibition and error-monitoring, by comparing the effects of variation in DAT1-genotype in typically developing adolescents and in adolescents diagnosed with ADHD.

Attention Deficit/Hyperactivity Disorder (ADHD) is a childhood disorder characterised by symptoms of inattention, hyperactivity and impulsivity (DSM-IV). The disorder is relatively common, with between 3-5% of school-age children affected (Buitelaar, 2002). ADHD symptoms persist into adulthood in about 65% of cases (Faraone, Biederman, & Mick, 2006) and individuals with ADHD often show impairments of executive function, including the ability to control and inhibit behaviour (Nigg, 2005).

ADHD is thought to have a strong genetic component, with additive effects of multiple genes explaining around 80% of an individual's susceptibility to ADHD

1
2
3
4 (Albayrack, Friedel, Schimmelmann, Hinney & Hebebrand, 2008). More than a decade of
5
6 molecular genetics research into ADHD has confirmed associations of small effect size
7
8 for a number of candidate genes, including those for the dopamine transporter (DAT1)
9
10 (Mick & Faraone, 2008). Since individual genetic influences on clinical phenotypes, such
11
12 as ADHD, are likely to be small and potentially heterogeneous, recent studies have
13
14 sought to link DNA variation in candidate genes to objective constructs such as a specific
15
16 psychological process or markers of brain function that are hypothesised to lie
17
18 intermediate between gene and disorder.
19
20
21
22

23
24 Within the cognitive genetics literature of ADHD, multiple lines of evidence
25
26 suggest that executive function measures, such as response inhibition, might index
27
28 susceptibility to ADHD. First, recent twin studies have demonstrated a remarkably strong
29
30 genetic contribution to executive functions, including response inhibition (Friedman et
31
32 al., 2008). Second, familial risk profiles for response inhibition deficits in ADHD have
33
34 also been established. Response inhibition deficits are more pronounced in ADHD
35
36 probands of affected as compared to non-affected parents (Crosbie, Pérusse, Barr &
37
38 Schachar, 2008). Non-affected siblings of children with ADHD have response inhibition
39
40 deficits that fall intermediate between typically developing children and children with
41
42 ADHD (Slaats-Willemse, Swaab-Barneveld, de Sonnevill, van der Meulen & Buitelaar,
43
44 2003). Neuroimaging studies of response inhibition have shown similar hypo-activation
45
46 of prefrontal areas in ADHD probands and their unaffected siblings, compared with
47
48 typically developing children (Durstun, Mulder, Casey, Ziermans, & van Engeland,
49
50 2006). Finally, a number of molecular genetic studies in both children with ADHD and
51
52 non-clinical participants have reported associations between allelic variation in DRD4
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 and DAT1 and measures of response inhibition (Cornish et al., 2005; Johnson et al.,
5
6 2008; though also see Kebir, Tabbane, Sengupta, & Jooper, 2009; Rommelse et al.,
7
8 2008). More recently, Durston and colleagues (2008) found that DNA variation in DAT1
9
10 influenced brain activation during response inhibition in the striatum and cerebellum.
11
12

13
14 In the present study, we used functional magnetic resonance imaging (fMRI) to
15
16 investigate the effects of the DAT1 40 base pair (bp) variable number of tandem repeat
17
18 (VNTR) polymorphism on the neural networks underlying response inhibition, in
19
20 adolescents diagnosed with ADHD as well as typically developing adolescents. The
21
22 dopamine transporter gene (DAT1) codes for the production of the dopamine transporter
23
24 (DAT) which clears dopamine from the synaptic cleft back into the neuron and is the
25
26 primary site of action for psychostimulants. Allelic variation within DAT1 has been
27
28 shown to affect the level of DAT expression, with increased expression associated with
29
30 the 10R-allele (Mill, Asherson, Browes, D'Souza, & Craig, 2002; though also see Van
31
32 Dyck et al., 2005). Thus, it is plausible that tonic levels of available dopamine are lower
33
34 in individuals homozygous for the 10R-allele, compared to other genotypes, and this may
35
36 affect cognitive control processes that rely on dopamine, such as response inhibition and
37
38 error-related processing. Individuals diagnosed with ADHD often show impairments in
39
40 error-monitoring (van Meel, Heslenfeld, Oosterlaan, & Sergeant, 2007) and/or response
41
42 inhibition (Vaidya et al., 1998).
43
44
45
46
47
48
49

50 Comparing the relative contribution of genotype and diagnosis allows us to
51
52 distinguish which brain differences in activation are affected by variation in DAT1
53
54 genotype, and which are likely caused by other (genetic or environmental) factors that
55
56 contribute to the disorder. Consequently, a main effect of genotype in the absence of
57
58
59
60
61
62
63
64
65

1
2
3
4 effects of diagnosis suggests that DAT1 genotype explains most of the variation in
5
6 activation in this region, with little additional contributions from other (genetic or
7
8 environmental) factors associated with ADHD. Similarly, regional brain functioning for
9
10 which ADHD-related impairment is mediated by genotype may be expected to show
11
12 genotype by diagnosis interactions while ADHD-related impairments not mediated by
13
14 genotype might be expected to show just main effects of diagnosis. In total, this approach
15
16 enables us to determine the contribution that the DAT1 gene makes to specific cognitive
17
18 and specific regional functional impairments that might be observed in those with
19
20
21
22
23
24 ADHD.

25 26 27 28 **MATERIAL AND METHODS**

29 30 **Participants**

31
32
33 A total of 58 right-handed adolescents were included in this study: 20 diagnosed
34
35 with ADHD (mean age: 14.1, SD: 2.1; 3 female) and 38 typically developing adolescents
36
37 (mean age: 13.26, SD: 1.98; 7 female); the proportion of genders did not differ between
38
39 the two groups ($p=.476$, Fisher's exact test), nor did their ages ($t(56)=1.5$, $p=.14$) (also
40
41 see Table 1, left panel). The participants with ADHD were referred by consultant
42
43 psychiatrists in Ireland. All participants with ADHD met DSM-IV criteria for ADHD, as
44
45 determined through semi-structured interviews by psychiatrists using the parent form of
46
47 the Child and Adolescent Psychiatric Assessment (CAPA) (Angold et al., 1995) or the
48
49 Parental Account of Children's Symptoms (PACS) (Taylor, 1986). Control participants
50
51 were recruited through local schools. Participants who were taking stimulant medication
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 for ADHD were asked to discontinue their medication for at least 24 hours¹ prior to the
5
6 scan session.
7

8
9 Exclusion criteria included known neurological conditions or pervasive
10 developmental disorders, serious head injuries and below average intelligence (below 70
11 on the WISC (Wechsler's Intelligence Scale for Children)), and MRI-exclusion criteria
12 (metal anywhere in the body (excluding the mouth), pregnancy, claustrophobia). In
13
14 addition, the parents of all participants completed the Conners' ADHD Rating Scale-
15 Revised: Long Version (Conners, 1996). Control adolescents had Conners' Global Index
16 H-scores ≤ 60 , and adolescents with ADHD had scores ≥ 65 . Participants diagnosed with
17 ADHD had lower scores on the WISC, as well as on both spelling and reading subtests
18 of the WRAT (Wide Range Achievement Test), and scored higher on the Conner's scale
19 (all $F \geq 11.6$, $p \leq .001$). There were no significant effects on these variables of genotype (all
20 $F \leq .67$, $p \geq .42$), nor were there significant interactions between genotype and diagnosis (all
21 $F \leq 1.2$, $p \geq .28$). We also present analyses on a subsample of the participants, who were
22 matched for their scores on the WISC and WRAT (all $F < 2.8$, $p \geq .105$), and which
23 included only boys (see Table 1, right panel).
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 After the study was described, participant and parental consent was provided. The
44 study was approved by the local ethics committee, and was carried out in accordance
45 with the Declaration of Helsinki (1964; 2000).
46
47
48
49

50 The data from some of the controls were previously reported in a developmental
51 study of inhibitory control (Braet et al., 2009).
52
53
54
55
56

57 ¹ We chose a relatively brief stimulant wash-out period, as this is less disruptive for participants. Acute
58 effects of stimulant withdrawal have been reported for both behavioural (e.g. Carlson & Kelly, 2003) and
59 neuroimaging data (e.g. Langleben et al., 2002). Our results are broadly consistent with those of a recent
60 study by Bédard et al., 2009, where a two-week stimulant withdrawal period was used.
61
62
63
64
65

Table_1_(a+b)

Genotyping

DNA was extracted from blood, or from saliva using Oragene DNA self-collections kits (DNAgenotek, Canada). Polymerase chain reaction (PCR) amplification and genotyping of the variable number of tandem repeats (VNTR) of the 3' untranslated region (UTR) of the DAT1 gene were conducted as described by Cook and colleagues (1995). All genotypes were in Hardy Weinberg Equilibrium. Children were grouped according to possession of the 10 repeat VNTR: the "high risk" (HR) group possessed two copies and the "low risk" (LR) group possessed one or no copies of the 10 repeat allele [ADHD: 9 HR, 11 LR and Control: 18 HR and 20 LR]. There was no significant association between allele possession for the 3' UTR VNTR DAT1 and clinical diagnosis [$\chi^2(1)=0.03$, $p=0.864$]. The categorisation of the genotypes as either HR or LR was done similar to prior studies (see e.g. Bellgrove, Hawi, Kirley, Gill, & Robertson, 2005), based on statistical grounds.

Sustained attention to response task (SART)

Participants performed the random SART, a go/no-go test that measures both inhibitory function (no-go trials) and sustained attention (go trials) (see O'Connell et al., 2009). In this paradigm, numbers from 1-9 are presented in a random (i.e. non-sequential)

1
2
3
4 order, and participants are asked to make a buttonpress response to each number, except
5
6 '3'. Each trial consisted of the (go or no-go) target (313 ms), after which a mask was
7
8 presented (563 ms, which consisted of 125ms mask, 63ms response cue, and a further
9
10 375ms mask), followed by a fixation cross (563 ms); inter-stimulus interval (ISI) was
11
12 1439 ms. (see Figure 1). The task was presented using E-prime (Psychology Software
13
14 Tools, Pittsburgh, USA), in a single block of 450 trials (consisting of 400 go trials and 50
15
16 no-go-trials), which included two 30s breaks after 150 and 300 trials.
17
18
19
20

21 We chose this design to be optimised to detect activation related to both
22
23 successful and unsuccessful nogo-trials (which are assumed to reflect response inhibition
24
25 and error-monitoring, respectively). To ensure a sufficient number of nogo-trials, while
26
27 keeping their proportion low (compared to the total number of trials, to encourage a
28
29 prepotent tendency to respond), the duration of individual trials was kept relatively short.
30
31 This design allows for reliable estimation of response maps for response inhibition and
32
33 error-monitoring, but is not ideal to investigate activation relating to tonic processes such
34
35 as sustained attention or motor processes given the fast pace of go-trials (faster than the
36
37 scan acquisition time) which means that hemodynamic response curves cannot be fitted
38
39 to go-events. Consequently, the baseline for assessing successful and unsuccessful nogo-
40
41 related activation is comprised of these tonic attentional and motoric processes. An
42
43 alternative approach would be to include a number of fixation-only trials which could
44
45 then serve as a low-level baseline against which tonic attentional and motor activity could
46
47 be assessed. Several factors motivated the choice of our design. First, our main interest
48
49 was response inhibition/error detection rather than sustained attention, and for this we
50
51 only required activation maps for nogo-trials. Similarly, it was not pertinent to our
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 research question to assess go-related activation: Previous studies have shown that go-
5 and nogo-trials lead to distinct maps with minimal overlap (Garavan, Ross & Stein,
6
7 and nogo-trials lead to distinct maps with minimal overlap (Garavan, Ross & Stein,
8
9 1999), and their differential time courses as revealed by ERP studies (e.g. O'Connell et
10 al., 2009) further confirm that both trial types elicit independent cognitive processes.
11
12 Second, the present design ensures that every trial is used in the analysis (either as
13
14 baseline, or as an effect of interest), which reduces the time that participants are in the
15
16 scanner which, in turn, makes it easier for participants (particularly the ADHD group) to
17
18 keep movement to a minimum. Third, response inhibition is commonly investigated by
19
20 comparing activation during nogo-trials to a baseline of go-trials (Braet et al., 2009;
21
22 Durston et al., 2008; Ramautar, Slagter, Kok & Ridderinkhof, 2006) with one advantage
23
24 being that one employs an active baseline condition with similar visual stimuli to the
25
26 nogo trials.
27
28
29
30
31
32
33
34
35

36 Figure_1

37 38 **MRI data acquisition**

39
40
41
42
43 All scanning was conducted on a Philips Intera Achieva 3.0 Tesla MR system.
44
45 Each scanning sequence began with a reference scan to resolve sensitivity variations. 180
46
47 high-resolution T1-weighted anatomic MPRAGE axial images (FOV 230 mm, thickness
48
49 0.9 mm, voxel size $0.9 \times 0.9 \times 0.9$) were then acquired (total duration 325 s), to allow
50
51 subsequent activation localization and spatial normalization.
52
53

54
55 Functional data were collected using a T2*-weighted echo-planar imaging (EPI)
56
57 sequence that acquired 32 non-contiguous (10% gap) 3.5 mm axial slices covering the
58
59
60
61
62
63
64
65

1
2
3
4 entire brain (TE = 35 ms, TR = 2000 ms, FOV 224 mm, 64 × 64 mm matrix size in
5
6 Fourier space). The functional scans had a total duration of 730 s, and were collected in a
7
8 single sequence.
9

10 11 12 13 14 **fMRI analysis**

15
16
17
18
19 The data were analysed under the general linear model using AFNI
20
21 (<http://afni.nimh.gov>) (Cox, 1996). Images were corrected for motion (using a least-
22
23 squares alignment allowing translations and rotations). These motion parameters were
24
25 also included as variables-of-no-interest in the first-level fMRI analyses; there were no
26
27 reliable differences in any of the 6 motion parameters between participants diagnosed
28
29 with ADHD and controls (all $t \leq 1.5$, $p \leq .139$) (no participants were excluded due to
30
31 excessive motion), and activation outside the brain was removed. MR signal drift was
32
33 estimated by fitting a 5th-order polynomial function. Separate impulse response functions
34
35 (IRFs) were estimated for successful inhibitions and commission errors using
36
37 deconvolution techniques (the time course of the two 30s breaks was also included as a
38
39 variable-of-no-interest). Gamma-variate functions were fit, voxelwise, to these IRFs
40
41 using a non-linear regression programme. A percentage signal-change score (%SC) was
42
43 calculated by dividing the area under the curve of these functions by the area under the
44
45 baseline which, in this case, reflects tonic ongoing processes involved in Go trial
46
47 responses². Individual %SC maps were then spatially blurred using a 3mm rms isotropic
48
49
50
51
52
53
54
55
56

57
58 ² Go-trials were not analysed separately, as the timing of our design was optimized to identify activation
59
60 maps for (successful and unsuccessful) nogo trials, rather than for go-trials, given the high frequency
61
62 (typically more than one in a single TR) of go-trials. Nogo-trials (which comprise approximately 10% of
63
64 the total number of trials) were spaced further apart, leading to more reliable IRFs. Prior studies (e.g.
65

1
2
3
4 Gaussian kernel, and transformed into MNI space using the MNI (Montréal Neurological
5
6
7 Institute) 152-brain template.

8
9 For both commission errors and successful inhibitions, activation maps for each
10
11 group of interest were then determined using one-sample t-tests against 0 (i.e. against the
12
13 null hypothesis of no change in activation compared with the baseline). Significant
14
15 voxels passed a voxelwise statistical threshold (ADHD HR: $t=5.04$; ADHD LR: $t=4.78$;
16
17 control HR: $t=3.97$; control LR: $t=3.88$, $p \leq .001$; these t-values varied as the sample sizes
18
19 differed between groups), and were required to be part of a cluster of significant voxels
20
21 with a minimum volume of 135 μl . This minimum cluster-size was determined using
22
23 Monte-Carlo simulations (1000 iterations), resulting in a 0.05 (corrected) probability of a
24
25 cluster surviving due to chance. Separate maps were generated for the four groups, and
26
27 these were subsequently combined into separate maps (containing every voxel that was
28
29 part of a significant cluster in any of the four groups) for successful inhibition and
30
31 commission errors. The clusters of activation in these resulting maps (one map for
32
33 successful inhibitions and one for commission errors) were then used for functionally-
34
35 defined region of interest (ROI) analyses wherein we extracted mean activation values for
36
37 each region for every participant (these same ROIs were used for the analyses on the
38
39 matched sample). These data were analysed as two-way ANOVAs with diagnosis and
40
41 genotype as grouping factors.
42
43
44
45
46
47
48
49

50 This method was chosen (rather than constructing a single map for all
51
52 participants), as it ensures that every region which was activated by at least one of the 4
53
54 groups is included in the ROI-analyses, rather than just regions which were commonly
55
56
57
58

59 Garavan, Ross & Stein, 1999) have investigated activation maps of both go- and nogo-trials, and found
60 relatively little overlap between the two maps.
61
62
63
64
65

1
2
3
4 activated (and which would otherwise likely be determined primarily by the (larger)
5
6 control group). This method tends to be more sensitive to identify group-differences (and
7
8 particularly, interaction-effects between multiple grouping-factors), as activation at the
9
10 ROI-level is more reliable than at the level of individual voxels (due to the reduction of
11
12 noise by averaging over a larger cortical area). While there are differences in the number
13
14 of events (e.g. omission errors) between the adolescents diagnosed with ADHD and the
15
16 controls, we have previously shown that performance-matching (in terms of the number
17
18 of omission and commission errors as well as response variability) between two groups
19
20 has only limited effects on the results of the ROI-analysis (Braet et al., 2009). This makes
21
22 it unlikely that the effects reported here are directly related to differences between the
23
24 groups in the number of errors they make, or even differences in performance on go-trials
25
26 (which were used as the baseline).
27
28
29
30
31
32
33
34
35
36
37

38 **Analysis of behavioural data**

39
40
41
42

43 Errors of commission (responses made on the no-go digit 3, which indicate failure
44
45 of inhibitory processes) and omission (non-responses on the go-trials, believed to reflect
46
47 temporary lapses in attention) and the mean and standard deviation (SD) of the response
48
49 times (RTs) on the go-trials were calculated for each participant. Differences between the
50
51 groups (as well as interactions between genotype and diagnosis) were assessed using
52
53 univariate analyses under the general linear model, with diagnosis (ADHD or control)
54
55 and DAT1-genotype (HR or LR) as fixed factors, and with the alpha level set at .05.
56
57
58
59
60
61
62
63
64
65

RESULTS

Behavioural performance

Adolescents diagnosed with ADHD made significantly more omission errors ($F(1,54)=18.48, p<.001$), and were significantly more variable in RT (higher SD) on go-trials ($F(1,54)=6.46, p=.014$) than typically developing adolescents, as well as displaying a trend for a higher number of commission errors ($F(1,54)=3.36, p=.072$). Participants diagnosed with ADHD and typically developing adolescents had similar RTs for both go trials (485s and 512s, respectively, $p>.05$) and commission errors (425s and 426s, respectively, $p>.05$). There were no interactions between diagnosis and genotype on any of the behavioural measures, nor were there any main effects of genotype.

The matched sample showed the same pattern of results: significantly more omission errors ($F(1,32)=16, p<.001$) and higher variability on go-trial RTs ($F(1,32)=4.8, p=.035$) for adolescents diagnosed with ADHD compared to typically developing adolescents, but there were no other significant main effects or interactions (all $F\leq 2.6, p\geq .117$).

Participants successfully responded to the majority of go trials (93% for adolescents diagnosed with ADHD, and 98% for the control group), suggesting that both groups had a prepotent tendency to respond on each trial. Response inhibition during infrequent nogo trials was therefore likely effortful, which is further confirmed by the

1
2
3
4 relatively high error rates for nogo trials (39% errors for participants from the ADHD
5
6 group, and 30% for the control group) (see Figure 2).
7
8
9

10
11 Figure_2
12
13

14 **fMRI-analysis: successful inhibitions**
15
16
17

18
19 Table_2
20
21
22
23
24
25

26 Table 2 lists the areas that showed significant activity when participants
27
28 successfully inhibited their response on no-go trials (also see Figure 3).
29
30

31 Adolescents with ADHD showed reduced activation, compared with controls, in
32
33 the right cuneus and in the left inferior parietal cortex.
34
35

36 There were also main effects of genotype, with greater activation for HR
37
38 compared with LR in the following regions: the right middle frontal gyrus, the ACC,
39
40 bilateral caudate, the right middle temporal gyrus and the right middle occipital gyrus.
41
42 One region, the left inferior temporal gyrus showed reduced activation for HR
43
44 participants.
45
46
47

48 There were also eight regions where there was an interaction between diagnosis
49
50 and genotype: the left superior frontal gyrus, the ACC, the left cingulate gyrus, the left
51
52 lentiform nucleus, the right supramarginal gyrus, the right precuneus, the right cuneus,
53
54 and the right middle occipital gyrus. In all these regions, there was a larger effect of
55
56 DAT1 genotype for participants with ADHD than for controls. (see Figure 4).
57
58
59
60
61
62
63
64
65

1
2
3
4 The matched sample showed the same pattern of results, though the difference in
5
6 a number of ROIs was no longer significant: there were no longer main effects of
7
8 diagnosis in the right cuneus and in the left inferior parietal cortex; the main effect of
9
10 genotype was no longer reliable in the left inferior temporal gyrus, though the other 5
11
12 clusters still showed increased activation for the HR group; finally, the right cuneus and
13
14 left cingulate gyrus no longer showed a reliable interaction between diagnosis and
15
16 genotype.
17
18
19
20
21
22

23
24 Figure_3
25
26
27

28
29 Figure_4
30
31
32
33
34
35
36
37

38 **fMRI analysis: commission errors**

39
40
41
42

43 Table_3
44
45
46
47

48 Table 3 lists the areas that showed significant activity when participants made a
49
50 commission error (responded on no-go trials) (also see Figure 5).
51
52

53 Adolescents diagnosed with ADHD showed reduced activation in three regions:
54
55 the left superior frontal gyrus, the ACC, and the right insula.
56
57
58
59
60
61
62
63
64
65

1
2
3
4 Participants with the HR-genotype showed reduced activation compared with
5
6 those with the LR-genotype in the following regions: the left parahippocampal gyrus, and
7
8 the left and right postcentral gyri.
9

10
11 Two regions showed an interaction between diagnosis and genotype: the right
12
13 middle frontal gyrus and the left angular gyrus (see Figure 6). In both regions, activation
14
15 levels were similar for the HR adolescents with ADHD and LR controls (both $t \leq 6.3$,
16
17 $p \geq .53$) with both groups showing activations in the right middle frontal gyrus and
18
19 deactivations in the left angular gyrus. The other two groups (LR ADHD and HR
20
21 controls) showed little activation in either region (all $t \leq 1.2$, $p \geq .24$).
22
23
24
25

26 The matched sample showed the same pattern of results, though the differences in
27
28 a number of ROIs were no longer significant: the ACC no longer showed a reliable effect
29
30 of diagnosis; the difference between HR and LR was no longer significant in the right
31
32 postcentral gyrus; the left angular gyrus no longer showed a significant interaction
33
34 between diagnosis and genotype.
35
36
37
38
39
40
41
42

43 Figure_5
44
45
46

47 Figure_6
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Discussion

Behavioural data

In line with findings from other studies, we observed significantly more omission errors made by adolescents with ADHD compared with typically developing adolescents, indicative of impaired sustained attention (Marchetta, Hurks, De Sonneville, Krabbendam, & Jolles, 2007) as well as increased response variability (Castellanos et al., 2005; Johnson et al., 2007). While the results were less clear in terms of response inhibition, there was also a trend for a higher number of commission errors, suggestive of impaired response inhibition for adolescents with ADHD, compared with typically developing adolescents. Impaired response inhibition has been proposed as a core deficit in ADHD (Wodka et al., 2007; but see e.g. Banaschewski et al., 2004 for an opposing view) as well as a potential endophenotype for ADHD (Aron & Poldrack, 2005).

No effects of DAT1 genotype on any of the behavioural performance measures were found, despite previous studies showing effects of DAT1 genotype on both response inhibition (Cornish et al., 2005) and response variability (Bellgrove et al., 2005) (though also see Kebir et al., 2009; Rommelse et al., 2008). Given the small effect size associated with the DAT1 gene, our study may simply lack the statistical power to replicate these effects (the HR-group in our sample did make more commission and omission errors, had higher response variability and slower RTs, but none of these differences were significant).

Genetic influence on functional activation patterns

In contrast to the behavioural effects, the neuroimaging data provided a more sensitive measure of the downstream effects of genetic variation on cognition (also see Greene, Braet, Johnson, & Bellgrove, 2008), insofar as they did reveal effects of genotype in the absence of behavioural effects (e.g. Durston et al., 2008; also see Bédard et al., 2009). During successful inhibitions, there was increased activation for the HR-genotype in the right middle frontal gyrus, the ACC and caudate, as well as the right middle temporal and middle occipital gyri, while there was reduced activation in the left inferior temporal gyrus. The right prefrontal cortex, as well as medial brain regions including the ACC and caudate are often implicated in response inhibition (Fassbender et al., 2004), and increased activation in these regions may imply that response inhibition is more effortful or less efficient for the HR-group.

During commission errors, however, the HR-group displayed reduced activation in the left parahippocampal gyrus, as well as in both the left and right postcentral gyri. The (para)hippocampal region has been associated with the regulation of arousal-mechanisms (Gray & McNaughton, 2000), and error-related activation in this region has been associated with subsequent improvements in performance in a task in which one needed to learn from one's errors (Hester, Barre, Murphy, Silk & Mattingley, 2008). Consequently, activation of this region during commission errors may reflect an arousal-mediated response to error-detection required to engage additional top-down control processes and prevent or reduce further errors. Given the role of dopamine in error-monitoring (Krämer et al., 2007) the present result may reflect a reduced error-response

1
2
3
4 due to the lower tonic levels of dopamine in the HR group, caused by increased
5
6 expression of the transporter associated with the 10-repeat allele.
7
8
9

10 11 **Interaction of genotype and diagnosis status on functional activation patterns** 12

13
14 There were a number of regions where genotype interacted with diagnosis. During
15
16 successful inhibitions, such interactions were observed in frontal (the left superior frontal
17
18 gyrus), medial and subcortical (the ACC, the left cingulate gyrus, and the left caudate),
19
20 and parietal regions (the right supramarginal gyrus, the right precuneus, and the right
21
22 cuneus), as well as in the right middle occipital gyrus. For all these regions we observed a
23
24 similar pattern: DAT1 genotype accounted for significant heterogeneity in
25
26 neurophysiological networks for response inhibition in children with ADHD (where
27
28 higher activation was observed in the HR group compared to LR) but not in controls
29
30 (where there were no differences in activation between the two genotypes). These data
31
32 demonstrate that the gene had greater penetrance at the level of brain activation
33
34 associated with response inhibition in the adolescents with ADHD (suggestive of more
35
36 effortful/ less efficient inhibition) but not in the typically developing adolescents. These
37
38 data further suggest that physiological heterogeneity in response inhibition networks in
39
40 the ADHD group might be predicted by variation in the DAT1. This may indicate that
41
42 the effect that DAT1 genetic variation has on the response inhibition endophenotype in
43
44 those with ADHD is not “self-contained” insofar as the same effect is not observed in the
45
46 control participants. Instead, other genetic or environmental influences particular to the
47
48 ADHD group must combine with DAT1 status to produce the observed brain activation
49
50 differences. We also observed interactions between diagnosis and genotype on brain
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 activation associated with commission errors, in the right middle frontal gyrus and the
5
6 left angular gyrus, though here the pattern was different, with similar engagement of both
7
8 regions for the LR control and the HR ADHD groups, while neither of the other groups
9
10 showed significant activation in these regions. Although the dopaminergic mechanisms
11
12 underlying these effects are unclear, this type of result indicates that dopaminergic
13
14 variation in healthy controls can produce some endophenotypic effects that are similar to
15
16 those with ADHD. This idea is consistent with the notion that ADHD is characterised by
17
18 a multifactorial polygenetic etiology (see Gizer, Ficks & Waldman, 2009), where the
19
20 effects of a large number of genes (on e.g. dopaminergic function) combine to form a
21
22 continuum of phenotypic variation in e.g. response inhibition.
23
24
25
26
27

28
29 It has been argued that the brain regions that show interactions between diagnosis
30
31 and genotype may contribute knowledge to aid development of new treatments for
32
33 ADHD, as a surrogate endpoint in individualized treatments targeting genotype/fMRI
34
35 activation profiles (Durston et al., 2008). The exact relationship between neurotransmitter
36
37 levels and changes in fMRI activation however is still unknown, and different studies
38
39 have often found contradictory effects. For the DAT1 gene, both the 10R and 9R alleles
40
41 have been associated with increased expression of the transporter (Mill et al., 2002 vs.
42
43 Van Dyck et al., 2005) and with deficits in sustained attention (Bellgrove et al., 2005 vs.
44
45 Kim, Kim & Cho, 2006). In a prior study by Durston and colleagues (2008), which also
46
47 investigated the interaction of ADHD diagnosis and DAT1 genotype using a similar
48
49 behavioural paradigm, increased activation was observed during successful inhibitions in
50
51 the vermis of the cerebellum for the 10R compared with the 9R allele. Although we also
52
53 observed higher activation in this region for the HR compared to LR group, this
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 difference was not significant in our sample. Durston and colleagues also observed an
5
6 interaction between diagnosis and genotype in the caudate nucleus in the striatum, but the
7
8 direction of this interaction effect was opposite to our result. It is unclear why our results
9
10 differ from those of Durston and colleagues. One possibility may be related to differences
11
12 in the difficulty of the respective go/nogo tasks that were used in both studies: in our
13
14 paradigm a single trial lasted only 1.4s, compared to 4s, and the proportion of nogo-trials
15
16 was 10%, compared to 25%. It is therefore likely that in the present study, response
17
18 inhibition was more difficult, and successful inhibitions more rewarding, leading to
19
20 significant differences in baseline dopaminergic release in the striatum between the two
21
22 studies. Similar to our own findings, a recent study by Bédard and colleagues (2009) also
23
24 found increased striatal activation in individuals diagnosed with ADHD who possess two
25
26 copies of the 10R allele. Further research will be required to clarify the nature of these
27
28 interactions, as well as their consistency across different study samples.
29
30
31
32
33
34
35

36 There were also a number of regions that showed differences between participants
37
38 diagnosed with ADHD and typically developing adolescents. During successful
39
40 inhibitions, the ADHD group showed reduced activation in two parietal regions, the right
41
42 cuneus and the left inferior parietal cortex, while during commission errors they showed
43
44 reduced activation of the left superior frontal gyrus, as well as the ACC and the right
45
46 insula, two key regions for response-monitoring. This confirms previous findings of
47
48 hypofunction in ADHD in frontal and parietal regions (Durston, 2003). The insula
49
50 projects to striatal regions, where deactivation during response inhibition has also been
51
52 observed in ADHD (Vaidya et al., 1998). The present results suggest that these ADHD-
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 related effects are not attributable to genetic variations in DAT1 but instead may be
5
6 driven by other genetic or environmental effects.
7
8
9

10 11 **Conclusions** 12 13

14 Imaging individuals with known genetic variation provides us with an exciting
15
16 opportunity to understand better the functional effects of DNA variation for brain and
17
18 cognition. Although we lack a full understanding of the pathways leading from gene to
19
20 changes in BOLD signaling, the current data suggest that fMRI indices may improve our
21
22 ability to detect subtle genetic effects compared with behavioural measures of cognition
23
24 alone. Here we have shown that DAT1 genotype influences the physiological substrates
25
26 of response inhibition and error processing, with effects for response inhibition being
27
28 most pronounced in children with ADHD. Our data add to a growing body of evidence
29
30 suggesting that differences in activation in brain regions underlying response inhibition in
31
32 ADHD are driven by genetic factors. By showing which behavioural effects and which
33
34 areas of cortical and subcortical activation are influenced by DAT1 genetic variation,
35
36 ADHD status or the interaction between the two, these results identify the specific
37
38 contribution of the dopamine transporter to executive functions such as response
39
40 inhibition and error-monitoring.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgements

This study was supported by grants from the Science Foundation Ireland, the Irish Health Research Board, the Irish Higher Education Authority's Programme for Research in Third-Level Institutions, the Australian National Health and Medical Research Council Howard Florey Centenary Fellowship, and the FWO Research Foundation - Flanders. Data-analysis was performed on computers of the Trinity Centre for High Performance Computing. The authors would like to thank all participants, as well as their parents.

Legend to the figures and tables:

Figure 1: The Sustained Attention to Response Task (SART): timecourse of a single trial.

Figure 2: Behavioural performance measures. Error bars represent 1 S.E.

Figure 3: Brain regions activated during successful inhibitions. a) superior frontal gyri and ACC, b) ACC, middle frontal gyrus and insulae, c) insulae and parietal lobe, d) parietal lobes. Green: areas activated but showing no diagnosis or genotype differences, Yellow: reduced activation for ADHD participants relative to controls; Dark Blue: increased activation for HR relative to LR; Red: diagnosis x genotype interaction.

Figure 4: Interactions between diagnosis x genotype, in percentage signal change during successful inhibitions.

1
2
3
4 Figure 5: Brain regions activated during commission errors. a) superior and middle
5 frontal gyri and ACC, b) ACC, middle and superior frontal gyri, and left and right
6
7 frontal gyri and ACC, b) ACC, middle and superior frontal gyri, and left and right
8
9 inferior parietal lobes c) insulae and parahippocampal gyrus, d) ACC, left angular gyrus,
10
11 left and right superior frontal gyri, left and right parietal lobes and postcentral gyri.
12
13 Green: areas activated but showing no diagnosis or genotype differences; Yellow:
14
15 reduced activation for ADHD participants relative to controls; Light Blue: increased
16
17 activation for LR relative to HR; Red: diagnosis x genotype interaction.
18
19
20
21
22

23
24 Figure 6: Interactions between diagnosis x genotype, in percentage signal change during
25
26 commission errors.
27
28
29
30

31 Table 1: Demographic information on the ADHD and control adolescents. The full
32
33 sample is displayed on the left, and the matched sample on the right.
34
35
36
37

38 Table 2: Brain regions activated during successful inhibitions. From left to right: location,
39
40 hemisphere, Brodman area(s), cluster-volume, coordinates (MNI), significance levels (*:
41
42 <.05; **:<.001) and direction for main effect of diagnosis, main effect of genotype, and
43
44 the interaction between diagnosis and genotype. Underlined effects were no longer
45
46 significant in the matched sample.
47
48
49
50

51
52
53 Table 3: Brain regions activated during commission errors. From left to right: location,
54
55 hemisphere, Brodman area(s), cluster-volume, coordinates (MNI), significance levels (*:
56
57 <.05; **:<.001) and direction for main effect of diagnosis, main effect of genotype, and
58
59
60
61
62
63
64
65

1
2
3
4 the interaction between diagnosis and genotype. Underlined effects were no longer
5
6 significant in the matched sample.
7
8
9

10 11 **References:**

- 12
13 Albayrak, O., Friedel, S., Schimmelmann, B. G., Hinney, A., & Hebebrand, J. (2008).
14
15 Genetic aspects in attention-deficit/hyperactivity disorder. *Journal of Neural*
16
17 *Transmission*, 115, 305-315.
18
19
20
21 Angold, A., Predergast, M., Cox, A., Harrington, R., Simonoff, E., & Rutter, M. (1995).
22
23 The Child and Adolescent Psychiatric Assessment (CAPA). *Psychological Medicine*, 25,
24
25 739-753.
26
27
28 Aron, R.A., Poldrack, R.A. (2005). The cognitive neuroscience of response inhibition:
29
30 Relevance for genetic research in attention-deficit/hyperactivity disorder. *Biological*
31
32 *Psychiatry*, 57, 1285–1292.
33
34
35
36 Banaschewski, T., Brandeis, D., Heinrich, H., Albrecht, B., Brunner, E., & Rothenberger,
37
38 A. (2004). Questioning inhibitory control as the specific deficit of ADHD – evidence
39
40 from brain electrical activity. *Journal of neural transmission*, 117, 841-864.
41
42
43 Bédard, A. C., Schulz, K. P., Cook, E. H. Jr, Fan, J., Clerkin, S. M., Ivanov, I., Halperin,
44
45 J. M., & Newcorn, J. H. (2009). Dopamine transporter gene variation modulates
46
47 activation of striatum in youth with ADHD. *Neuroimage*,
48
49 doi:10.1016/j.neuroimage.2009.12.041
50
51
52
53 Bellgrove, M. A., Hawi, Z., Kirley, A., Gill, M., & Robertson, I. H. (2005). Dissecting
54
55 the attention deficit hyperactivity disorder (ADHD) phenotype: sustained attention,
56
57
58
59
60
61
62
63
64
65

1
2
3
4 response variability and spatial attentional asymmetries in relation to dopamine
5 transporter (DAT1) genotype. *Neuropsychologia*, 43, 1847-1857.

6
7
8
9 Braet, W., Johnson, K. A., Tobin, C. T., Acheson, R., Bellgrove, M. A., Robertson, I. H.,
10 & Garavan, H. (2009). Functional developmental changes underlying response inhibition
11 and error-detection processes. *Neuropsychologia*, 47, 3143-3151.

12
13
14
15
16 Buitelaar, J. K. (2002). Epidemiology: what have we learned over the last decade? In:
17 Sandberg S. (editor), *Hyperactivity and attention-deficit disorders*. Cambridge, UK:
18 Cambridge University Press.

19
20
21
22
23 Burock, M. A., Buckner, R. L., Woldorff, M. G., Rosen, B. R., & Dale, A. M. (1998).
24 Randomized event-related experimental designs allow for extremely rapid presentation
25 rates using functional MRI. *Neuroreport*, 9, 3735-3739.

26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
Carlson, G. A., & Kelly, K. L. (2003). Stimulant rebound: how common is it and what
does it mean? *Journal of Child and Adolescent Psychopharmacology*, 13, 137-142.

Castellanos, F.X., Sonuga-Barke, E.J.S., Scheres, A., Di Martino, A., Hyde, C. &
Walters, J.R. (2005). Varieties of attention-deficit/hyperactivity disorder-related intra-
individual variability. *Biological Psychiatry*, 57, 1416–1423.

Conners, K. (1996). *Rating scales in ADHD*, Duke University Medical Center.

Cook, E. H., Stein, M. A., Krasowski, M. D., Cox, N. J., Olkon, D. M., Kieffer, J. E., &
Leventhal, B. L. (1995). Association of Attention-Deficit Disorder and the Dopamine
Transporter Gene. *American Journal of Human Genetics*, 56, 993-998.

Congdon, E., Lesch, K. P., & Canli, T. (2007). Analysis of DRD4 and DAT
polymorphisms and behavioral inhibition in healthy adults : implications for impulsivity.
American Journal of Medical Genetics Part B: Neuropsychiatric Genetics, 147B, 27-32.

1
2
3
4 Cornish, K. M., Manly, T., Savage, R., Swanson, J., Morisano, D., Butler, N., Grant, C.,
5
6 Cross, G., Bentley, L., & Hollis, C. P. (2005). Association of the dopamine transporter
7
8 (DAT1) 10/10-repeat genotype with ADHD symptoms and response inhibition in a
9
10 general population sample. *Molecular Psychiatry*, 10, 686-698.

11
12
13
14 Cox, R. W. (1996) AFNI: software for analysis and visualization of functional magnetic
15
16 resonance neuroimages. *Computers and Biomedical Research*, 29, 162-173.

17
18
19 Crosbie, J., Pérusse, D., Barr, C. L., & Schachar, R. J. (2008). Validating psychiatric
20
21 endophenotypes: inhibitory control and attention deficit hyperactivity disorder.
22
23 *Neuroscience and Biobehavioral Reviews*, 32, 40-55.

24
25
26 Durston, S. (2003). A review of the biological bases of ADHD: what have we learned
27
28 from imaging studies? *Mental Retardation and Developmental Disabilities Research*
29
30 *Reviews*, 9, 184–195.

31
32
33 Durston, S., Fossella, J. A., Mulder, M. J., Casey, B. J., Ziermans, T. B., Vessaz, M. N.,
34
35 & Van Engeland, H. (2008). Dopamine transporter genotype conveys familial risk of
36
37 attention-deficit/hyperactivity disorder through striatal activation. *Journal of the*
38
39 *American Academy for Child and Adolescent Psychiatry*, 47, 61-67.

40
41
42 Durston S, Mulder M, Casey BJ, Ziermans T, & van Engeland H. (2006). Activation in
43
44 ventral prefrontal cortex is sensitive to genetic vulnerability for attention-deficit
45
46 hyperactivity disorder. *Biological Psychiatry*, 60, 1062-1070.

47
48
49 Faraone, S. V., Biederman, J., & Mick, E. (2006). The age-dependent decline of attention
50
51 deficit hyperactivity disorder: a meta-analysis of follow-up studies. *Psychological*
52
53 *Medicine*, 36, 159-165.

1
2
3
4 Fassbender, C., Murphy, K., Foxe, J., Wylie, G., Javitt, D.C., Robertson, I.H., &
5
6 Garavan, H. (2004). A Topography of Executive Functions revealed by functional
7
8 Magnetic Resonance Imaging. *Cognitive Brain Research*, 20, 132-143.

9
10
11 Garavan, H, Ross, T. J., & Stein, E. A. (1999). Right hemispheric dominance of
12
13 inhibitory control: an event-related functional MRI study. *Proceedings of the National*
14
15 *Academy of Sciences of the United States of America*, 96, 8301-8306.

16
17
18 Friedman, N. P., Miyake, A., Young, S. E., Defries, J. C., Corley, R. P., & Hewitt, J. K.
19
20 (2008). Individual differences in executive functions are almost entirely genetic in origin.
21
22 *Journal of Experimental Psychology: General*, 137, 201-225.

23
24
25 Gizer, I. R., Ficks, C., & Waldman, I. D. (2009). Candidate gene studies of ADHD: a
26
27 meta-analytic review. *Human Genetics*, 126, 51–90.

28
29
30 Goldberg, T. E., & Weinberger, D. R. (2004). Genes and the parsing of cognitive
31
32 processes. *Trends in Cognitive Sciences*, 8, 325-335.

33
34
35 Gray, J. A. & McNaughton, N. (2000). *The neuropsychology of anxiety: an enquiry into*
36
37 *the functions of the septo-hippocampal system (second edition)*, Oxford University Press,
38
39 Oxford.

40
41
42 Greene, C. M., Braet, W., Johnson, K. A., & Bellgrove, M. A. (2008). Imaging the
43
44 genetics of executive function. *Biological Psychology*, 79, 30-42.

45
46
47 Hester, R., Barre, N., Murphy, K., Silk, T., & Mattingley, J. B. (2008). Human medial
48
49 frontal cortex activity predicts learning from errors. *Cerebral Cortex*, 18, 1933-1940.

50
51
52 Johnson, K. A., Kelly, S. P., Bellgrove, M. A., Barry, E., Cox, M., Gill, M., & Robertson,
53
54 I. H. (2007). Response variability in attention deficit hyperactivity disorder: evidence for
55
56 neuropsychological heterogeneity. *Neuropsychologia*, 45, 630-638.

1
2
3
4 Johnson, K. A., Kelly, S. P., Robertson, I. H., Barry, E., Mulligan, A., Daly, M., Lambert,
5
6 D., McDonnell, C., Connor, T. J., Hawi, Z., Gill, M., & Bellgrove, M. A. (2008).

7
8
9 Absence of the 7-repeat variant of the DRD4 VNTR is associated with drifting sustained
10
11 attention in children with ADHD but not in controls. *American Journal of Medical*
12
13 *Genetics Part B Neuropsychiatric Genetics*, 147B, 927-937.

14
15
16 Kebir, O., Tabbane, K., Sengupta, S., & Joobar, R. (2009). Candidate genes and
17
18 neuropsychological phenotypes in children with ADHD: review of association studies.
19
20
21 *Journal of Psychiatry and Neuroscience*, 34, 88-101.

22
23
24 Kim, J. W., Kim, B. N., & Cho, S. C. (2006). The dopamine transporter gene and the
25
26 impulsivity phenotype in attention deficit hyperactivity disorder: a case-control
27
28 association study in a Korean sample. *Journal of Psychiatric Research*, 40, 730-737.

29
30
31 Krämer, U. M., Cunillera, T., Càmarà, E., Marco-Pallarés, J., Cucurell, D., Nager, W.,
32
33 Bauer, P., Schüle, R., Schöls, L., Rodriguez-Fornells, A., & Münte, T. F. (2007). The
34
35 impact of catechol-O-methyltransferase and dopamine D4 receptor genotypes on
36
37 neurophysiological markers of performance monitoring. *Journal of Neuroscience*, 27,
38
39 14190-14198.

40
41
42
43 Langleben, D. D., Acton, P. D., Austin, G., Elman, I., Krikorian, G., Monterosso, J. R.,
44
45 Portnoy, O., Ridlehuber, H. W., & Strauss, H. W. (2002). Effects of methylphenidate
46
47 discontinuation on cerebral blood flow in prepubescent boys with attention deficit
48
49 hyperactivity disorder. *Journal of Nuclear Medicine*, 43, 1624-1629.

50
51
52
53 Marchetta, N. D., Hurks, P. P., De Sonneville, L. M., Krabbendam, L., & Jolles, J.
54
55 (2007). Sustained and focused attention deficits in adult ADHD. *Journal of Attention*
56
57 *Disorders*, 11, 664-676.

1
2
3
4 Mick, E., & Faraone, S. V. (2008). Genetics of attention deficit hyperactivity disorder.
5
6 *Child and Adolescent Psychiatric Clinics of North America*, 17, 261-284.

7
8
9 Mill, J., Asherson, P., Browes, C., D'Souza, U., & Craig, I. (2002). Expression of the
10
11 dopamine transporter gene is regulated by the 3' UTR VNTR: Evidence from brain and
12
13 lymphocytes using quantitative RT-PCR. *American Journal of Medical Genetics*, 114,
14
15 975–979.

16
17
18 Nigg, J. T. (2005). Neuropsychologic theory and findings in attention-
19
20 deficit/hyperactivity disorder: the state of the field and salient challenges for the coming
21
22 decade. *Biological Psychiatry*, 57, 1424–1435.

23
24
25
26 O'Connell, R. G., Dockree, P. M., Bellgrove, M. A., Turin, A., Ward, S., Foxe, J. J., &
27
28 Robertson, I. H. (2009). Two Types of Action Error: Electrophysiological Evidence for
29
30 Separable Inhibitory and Sustained Attention Neural Mechanisms Producing Error on
31
32 Go/No-go Tasks. *Journal of Cognitive Neuroscience*, 21, 93-104.

33
34
35
36 Ramautar, J. R., Slagter, H. A., Kok, A., & Ridderinkhof, K. R. (2006). Probability
37
38 effects in the stop-signal paradigm: the insula and the significance of failed inhibition.
39
40 *Brain Research*, 1105, 143-154.

41
42
43 Rommelse, N. N., Altink, M. E., Arias-Vásquez, A., Buschgens, C. J., Fliers, E., Faraone,
44
45 S. V., Buitelaar, J. K., Sergeant, J. A., Franke, B., & Oosterlaan, J. (2008). A review and
46
47 analysis of the relationship between neuropsychological measures and DAT1 in ADHD.
48
49 *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, 147B, 1536-
50
51 1546.

52
53
54
55 Slaats-Willemse, D., Swaab-Barneveld, H., de Sonneville, L., van der Meulen, E., &
56
57 Buitelaar, J. (2003). Deficient response inhibition as a cognitive endophenotype of
58
59
60
61
62
63
64
65

1
2
3
4 ADHD. *Journal of the American Academy of Child and Adolescent Psychiatry*, 42, 1242-
5
6 1248.

7
8
9 Taylor, E. A. (1986), Childhood hyperactivity. *British Journal of Psychiatry*, 149, 562-
10
11 573.

12
13
14 Vaidya, C. J., Austin, G., Kirkorian, G., Ridlehuber, H. W., Desmond, J. E., Glover, G.
15
16 H., & Gabrieli, J. D. (1998). Selective effects of methylphenidate in attention deficit
17
18 hyperactivity disorder: a functional magnetic resonance study. *Proceedings of the*
19
20 *National Academy of Sciences of the United States of America*, 95, 14494 –14499.

21
22
23 van Meel, C. S., Heslenfeld, D. J., Oosterlaan, J., & Sergeant, J. A. (2007). Adaptive
24
25 control deficits in attention-deficit/hyperactivity disorder (ADHD): the role of error
26
27 processing. *Psychiatry Research*, 151, 211-220.

28
29
30 Van Dyck, C. H., Malison, R. T., Jacobsen, L. K., Seibyl, J. P., Staley, J. K., Laruelle,
31
32 M., Baldwin, R. M., Innis, R. B., & Gelernter, J. (2005). Increased dopamine transporter
33
34 availability associated with the 9-repeat allele of the SLC6A3 gene. *Journal of Nuclear*
35
36 *Medicine*, 46, 745-751.

37
38
39 Wodka, E.L., Mahone, E.M., Blankner, J.G., Larson, J.C., Fotedar, S., Denckla, M.B., &
40
41 Mostofsky, S. H. (2007). Evidence that response inhibition is a primary deficit in ADHD.
42
43
44
45
46 *Journal of Clinical and Experimental Neuropsychology*, 29, 345–356.

	ADHD HR (n=9)	ADHD LR (n=11)	Controls HR (n=18)	Controls LR (n=20)
Male	6	11	15	16
	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)
Age in years	14.7 (1.6)	13.6 (2.4)	12.8 (1.9)	13.4 (2.1)
IQ	99.7 (14.4)	92.1 (16)	110.7 (13.5)	111.8 (13.1)
WRAT Reading	93.9 (10.7)	96.9 (13.8)	107.3 (15.1)	107.1 (8.7)
WRAT Spelling	92.4 (9.8)	94.7 (14.2)	108.7 (14.3)	108.5 (7.1)
Conners' ADHD Index	76.6 (8.7)	75 (8.5)	42.8 (2.8)	44 (3.1)

	ADHD HR (n=5)	ADHD LR (n=8)	Controls HR (n=10)	Controls LR (n=10)
Male	5	8	10	10
	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)
Age in years	14.2 (1.9)	12.8 (2.4)	13.0 (2.1)	13.1 (2.1)
IQ	104 (12.1)	98.9 (12.8)	106.3 (13.5)	107.2 (12.6)
WRAT Reading	100.2 (9.8)	101.5 (8.5)	104.1 (15.3)	105.8 (8.8)
WRAT Spelling	97.8 (8.3)	100.1 (12)	102.1 (8.4)	106.7 (6.8)
Conners' ADHD Index	75.4 (8.8)	76 (8.6)	43.6 (3.4)	44.8 (3.4)

name	HS	BA	Vol (μl)	X	Y	Z	P(diagnosis)	P(genotype)	P(interaction)
frontal lobes									
ACC	BI	6	28478	3	9	45		*	HR>LR *
superior frontal gyrus	R	9/10	15167	36	39	34			
superior frontal gyrus	L	9/10	6041	-34	44	34			**
inferior frontal gyrus	R	9/6	1011	45	2	42			
middle frontal gyrus	L		365	-28	-8	46			
middle frontal gyrus	R	46/9	176	46	15	31		*	HR>LR
precentral gyrus	R	6	211	48	-9	60			
precentral gyrus	L	4	172	-25	-25	70			
cingulate gyrus	L	24/31	155	-15	-21	37			*
subcortical regions									
PCC	BI	23	3674	0	-29	25			
insula	R	13	16400	34	14	-1			
insula	L	47	8724	-33	18	1			
thalamus	R		285	8	-22	8			
thalamus	L		227	-18	-26	6			
caudate	BI		223	3	0	9		*	HR>LR
lentiform nucleus	L		183	-12	5	5			*
mammillary body	BI		188	0	-11	-8			
cerebellum									
cerebellar vermis	BI		847	2	-27	-25			
parietal lobes									
supramarginal gyrus	R	40	17116	51	-47	32			*
supramarginal gyrus	L	40	247	-54	-49	23			
precuneus	R	7	3824	8	-77	34			*
precuneus	R	31	628	13	-38	40			-
cuneus	R	18	306	11	-72	-5	*	<u>c>a</u>	**
angular gyrus	L	39	607	-33	-61	34			
superior parietal lobule	R	7/39	720	30	-66	33			
inferior parietal lobule	L	40	479	-58	-46	20			
inferior parietal lobule	L	40	168	-50	-45	43	*	<u>c>a</u>	
temporal lobes									
middle temporal gyrus	R	20	862	50	-25	-15		**	HR>LR
inferior temporal gyrus	L	20	398	-48	-24	-22		*	<u>LR>HR</u>
superior temporal gyrus	L	13	153	-45	-41	16		-	
superior temporal gyrus	L	38	152	-55	11	-10			
occipital lobes									
middle occipital gyrus	R		192	50	-74	-17		*	HR>LR **


name	HS	BA	Vol (μ l)	X	Y	Z	P(diagnosis)	P(genotype)	P(interaction)
frontal lobes									
ACC	BI	32	23659	0	14	43	*	<u>c>a</u>	
superior frontal gyrus	L	9/10	2179	-33	41	31	*	c>a	
superior frontal gyrus	R	9/10	1992	31	42	37			
middle frontal gyrus	R	10	179	40	39	22			*
precentral gyrus	R	13	263	45	-15	8			
subcortical regions									
PCC	BI	23	885	0	-24	29			
insula	R	44	18975	42	10	7	*	c>a	
insula	L	13	11383	-37	13	0			
thalamus	L		270	-12	-19	6			
caudate	R		979	10	5	6			
caudate	L		684	-9	4	1			
parahippocampal gyrus	BI		4030	3	-24	-18			
parahippocampal gyrus	L	30	270	-20	-40	-6		*	LR>HR
culmen	L	36	189	-21	-36	-21			
cerebellum									
declive	L	37	157	-49	-55	-25			
parietal lobes									
inferior parietal lobule	L	40	5689	-58	-44	27			
inferior parietal lobule	R	40	4007	56	-49	34			
supramarginal gyrus	L	40	354	-39	-54	36			
cuneus	L	7	162	-12	-78	32			
cuneus	R	18	216	10	-76	19			
precuneus	R	7	198	12	-78	39			
angular gyrus	L	39	152	-46	-77	33			**
postcentral gyrus	L	3	956	-53	-26	42		*	LR>HR
postcentral gyrus	R	40	281	57	-26	32		*	<u>LR>HR</u>
temporal lobes									
superior temporal gyrus	R	22	637	53	-47	12			
transverse temporal gyrus	L	41	544	-54	-22	13			
inferior temporal gyrus	R	20	251	51	-21	-22			

Figure 1
[Click here to download high resolution image](#)

Sustained Attention to Response Task (SART)



Manuscript







Figure 4
[Click here to download high resolution image](#)






Figure 6
[Click here to download high resolution image](#)

Mean % signal change during commission errors

