

SUPINE HYPERTENSION

Supine Hypertension Is Associated With an Impaired Cerebral Oxygenation Response to Orthostasis: Finding From The Irish Longitudinal Study on Ageing

Louise Newman¹, John D. O'Connor², Roman Romero-Ortuno³, Richard B. Reilly⁴, Rose Anne Kenny⁵

ABSTRACT: The cerebrovascular effects of supine hypertension (SH) are still poorly understood. With aging and atherosclerosis of the vascular system, it is not uncommon for SH and non-neurogenic orthostatic hypotension to co-occur. Given evidence for end organ damage and more extreme cerebral dysfunction in those with SH-orthostatic hypotension, we hypothesized that SH would be associated with impaired cerebral autoregulation. The aim of this study was to characterize the cerebrovascular response to orthostasis. Near-infrared spectroscopy was used to quantify the cerebrovascular response. We analyzed data from Wave 3 of TILDA (The Irish Longitudinal Study on Ageing; n=2750). Cerebral oxygenation and blood pressure (BP) were monitored continuously during an active stand. Responses were modeled using multilevel mixed-effects models and adjusted for important covariates such as age, sex, education, antihypertensive medications, and comorbidities. Forty-nine percent of the sample had SH. Those with SH demonstrated an impaired BP response and a slower recovery of BP after standing, graded by severity of SH. The cerebral oxygenation response was similar for both groups, but when standardized to mean arterial BP, the response was impaired in those with SH. A deficit of -0.83% (95% CI, -0.93 to -0.74) remained after 3 minutes of standing. Our study determined that cerebral oxygenation and cerebral autoregulation are impaired in those with SH. In older patients, consideration should be given to measuring SH and screening for orthostatic hypotension. Therapeutic studies are needed to better understand the relationship between cerebral oxygenation, medications, supine BP, and orthostatic hypotension. (*Hypertension*. 2021;78:210–219. DOI: 10.1161/HYPERTENSIONAHA.121.17111.) • **Data Supplement**

Key Words: aging ■ blood pressure ■ cerebral hemodynamics ■ hemodynamics ■ hypotension, orthostatic

Supine hypertension (SH) is a paradoxical elevation in blood pressure (BP) upon assuming a supine position for 5 to 10 minutes. It is a potential risk factor for end organ damage and poor cardiovascular outcomes, but its cerebrovascular effects are still poorly understood.¹ To date, most research on SH has been conducted in the context of neurogenic orthostatic hypotension (OH).¹ However, with aging and atherosclerosis of the vascular system, it is not uncommon for SH and non-neurogenic OH to co-occur. This poses a therapeutic challenge.²

Cerebral autoregulation (CA) is the process that maintains a constant cerebral blood flow and oxygenation

despite changes in systemic BP. CA is essential for normal cognitive function and brain health. The autonomic nervous system and the baroreflex are key in adapting to BP fluctuations, such as those induced by postural change, by modulating cardiovascular resistance via vasoconstriction and vasodilation to ensure a constant flow. Maintaining a constant cerebral blood flow may be a challenge at extremes of BP.³ With age, mechanisms that adapt to maintain peripheral BP can be deficient. These include decreased baroreflex sensitivity⁴ and increased vascular stiffness, resulting in increased pulsatility, as well as decreased muscle pump,⁵ which can lead to extreme BP and challenge CA.

Correspondence to: Louise Newman, The Irish Longitudinal Study on Ageing, Trinity College Dublin, Dublin 2, Ireland. Email louise.newman@tcd.ie

The Data Supplement is available with this article at <https://www.ahajournals.org/doi/suppl/10.1161/HYPERTENSIONAHA.121.17111>.

For Sources of Funding and Disclosures, see page 218.

© 2021 American Heart Association, Inc.

Hypertension is available at www.ahajournals.org/journal/hyp

Novelty and Significance

What Is New?

- Those with supine hypertension demonstrated an impaired blood pressure response and slower recovery in response to orthostasis, graded by severity of supine hypertension.
- Those with supine hypertension displayed deficits in maintaining cerebral oxygenation after standing, when the response was standardized to mean arterial blood pressure.
- Our results suggest cerebral autoregulation is impaired in those with supine hypertension.

What Is Relevant?

- Impaired stabilization is relevant in the context of managing hypertension, as well as orthostatic hypotension, and identifying those at risk for falls.
- Currently supine hypertension is not routinely screened for in the clinic.

Summary

There is evidence of impaired cerebral oxygenation and cerebral autoregulation in this large cohort of older adults. Future clinical studies are needed to investigate the relationship between cerebral oxygenation, supine blood pressure, medications, and orthostatic hypotension.

Nonstandard Abbreviations and Acronyms

BP	blood pressure
CA	cerebral autoregulation
DBP	diastolic blood pressure
MAP	mean arterial pressure
NIRS	near-infrared spectroscopy
OH	orthostatic hypotension
PWV	pulse wave velocity
SBP	systolic blood pressure
SH	supine hypertension
TILDA	The Irish Longitudinal Study on Ageing
TSI	tissue saturation index
TUG	timed up and go
WMH	white matter hyperintensities

OH is defined by consensus statement as a sustained drop ≥ 20 mm Hg in systolic BP (SBP) or ≥ 10 mm Hg in diastolic BP (DBP) upon assuming the upright position.⁶ It is associated with falls,⁷ frailty, cognitive impairment,⁸ dementia,⁹ depression,¹⁰ cardiovascular morbidity, and mortality.¹¹ Impaired CA and cerebral hypoperfusion is a pathway that has been implicated in these outcomes.

In some cases of SH, BP fails to dip at night.¹² This is associated with natriuresis and consequently volume depletion, all of which exaggerate morning OH, increasing the risk of falls.¹³ Nocturnal nondipping is also a risk factor for cardiovascular damage, including cerebrovascular damage.¹⁴

Our group has already demonstrated, both cross-sectionally and longitudinally, that those with SH-OH display poorer cognitive performance alone across multiple cognitive domains than those with SH alone.^{8,15} In a small synucleinopathy cohort ($n=57$; 67% had SH), SH

was associated with increased white matter hyperintensities (WMH), decreased renal function, greater left ventricular hypertrophy, earlier cardiovascular events, and shorter survival times from these events.¹⁶ A previous study by TILDA (The Irish Longitudinal Study on Ageing) using cardiac beat-to-beat recording technology identified a pattern in those with SH, which demonstrated a large drop in BP and nonrecovery to baseline upon standing.¹⁷ This pattern was a predictor of orthostatic intolerance, suggesting a higher hypotensive burden in those with SH. The cerebrovascular response remains to be examined directly.

Given evidence for end organ damage and more extreme cerebral dysfunction in those with SH-OH, we hypothesized that SH would be associated with impaired CA. The aim of this study was to characterize the cerebrovascular response to orthostasis in a large cohort of older adults with SH. Near-infrared spectroscopy (NIRS) was used to quantify the cerebrovascular response.

METHODOLOGY

Data Availability

Due to the sensitive nature of the data collected for this study, requests to access the dataset from qualified researchers should be directed to TILDA, Trinity College Dublin, Dublin 2, via email at tilda@tcd.ie.

Sample

This was a cross-sectional study that utilized data from Wave 3 (2014–2015) of TILDA. TILDA is a large prospective cohort study collecting a range of health, social, and economic data on those aged ≥ 50 years living in Ireland. The sampling and design have been described previously.^{18,19} Data collection began in 2009 with 8504 community-dwelling adults. Participants complete a computer-assisted personal interview and self-completion

questionnaire every 2 years and every ≈ 4 years are invited for a comprehensive health assessment. The study was granted ethical approval by the Research Ethics Committee at Trinity College Dublin and complied with the Declaration of Helsinki. Written informed consent was obtained from all participants. Participants who completed an active stand challenge at Wave 3 were eligible for this analysis. Those with known Alzheimer, dementia, or Parkinson disease were excluded due to known effects on cerebral blood flow and autonomic control.

Active Stand Protocol

Participants lay supine for ≈ 10 minutes in a room at ambient temperature before being asked to stand as quickly as possible. They were aided by a nurse when required. They remained standing quietly for a further 3 minutes. Cerebral oxygenation was continuously recorded by an NIRS Portalite system (Artinis Medical Systems, Zetten, the Netherlands) at 50 Hz. A single probe was placed on the left forehead, approximately at a position corresponding to FP1 of the international 10- to 20-electrode placement system.²⁰ Using 2 different wavelengths (nominally 760 and 850 nm), the device determines concentrations of oxygenated hemoglobin and deoxygenated hemoglobin based on the principle of the modified Beer-Lambert law.²¹ Multiple transmitters with inter-optode distances of 30, 35, and 40 mm allow determination of absolute values of cerebral oxygenation using spatially resolved spectroscopy. Based on inter-optode distances, measurements are at a maximum depth of 20 mm. Spatially resolved spectroscopy minimizes the extracerebral influences.²² The output, tissue saturation index (TSI), was calculated as oxygenated hemoglobin expressed as a percentage of total oxygenated hemoglobin and deoxygenated hemoglobin. BP, heart rate, and cardiac output were continuously monitored using digital photoplethysmography (Finometer; Finapres Medical Systems, Arnhem, the Netherlands) sampled at 200 Hz. The height correction unit, which is part of the Finometer, corrected for hydrostatic pressure differences between the finger cuff and the heart level. At the start of each test, the height sensors were calibrated by nulling.²³ Then one was sensor placed on the finger cuff and one placed on an arm sling at heart level. Mean arterial pressure (MAP) was determined by the Finometer as the true integrated mean pressure between the current and the next upstroke.^{23,24}

Signal Processing

NIRS and Finometer data were resampled to 1 Hz and smoothed using a 10-s moving average filter and an 11-point median filter. These were implemented in MATLAB R2016b. The onset of the stand and time to transition from supine to standing (standing speed) were determined by applying an algorithm using the Finometer height correction unit data.²⁵ Baseline was calculated from the average response 60 to 30 s before stand onset. Data from a number of participants were excluded due to technical problems with equipment or signal processing. These cases can reasonably be assumed to be random occurrences.

Supine Hypertension

SH was first classified as binary yes (SBP ≥ 140 mm Hg and DBP ≥ 90 mm Hg) or no (SBP < 140 mm Hg and DBP < 90

mm Hg) using baseline values. A graded severity classification of none (SBP < 140 mm Hg and DBP < 90 mm Hg), mild (SBP, 140–159 mm Hg and DBP, 90–99 mm Hg), moderate (SBP, 160–179 mm Hg and DBP, 100–109 mm Hg), or severe (SBP ≥ 180 mm Hg and DBP ≥ 110 mm Hg) was also used. These cutoffs are in line with recent guidelines for neurogenic SH²⁶ in the absence of established non-neurogenic SH cutoffs.

Covariates

Selected covariates included age, sex, educational attainment, smoking history, problematic alcohol intake based on the Cut down, Annoyed, Guilty, and Eye-opener questionnaire,²⁷ diabetes, number of cardiovascular conditions (transient ischemic attack, stroke, hypertension, angina, heart attack, congestive heart failure, high cholesterol, heart murmur, or any abnormal heart rhythm), cancer, kidney disease, antihypertensive medications (anatomical therapeutic chemical codes C02, C03, C07, C08, or C09), antidepressant medications (anatomical therapeutic chemical code N06A), benzodiazepines (anatomical therapeutic chemical code N05BA), antipsychotics (anatomical therapeutic chemical code N05A), timed-up-and-go (TUG) test (time taken to stand up from a chair, walk 3 m, turn, and sit back down), height, body mass index, depressive symptoms (short form of the Centre for Epidemiological Studies Depression) scale,²⁸ standing speed, change in cardiac output, and pulse wave velocity (PWV). Carotid-femoral PWV was measured using Vicorder (Skidmore Medical, Ltd, Bristol, United Kingdom) and calculated as an average of 2 measurements. Seated BP was calculated as an average of 2 readings from an automatic oscillometric sphygmomanometer (Omron).

Statistical Modeling

Data were analyzed using Stata 14 (StataCorp, TX). Our analysis includes active stand data at 10-s intervals from 0 to 180s, with time 0 s being the onset of the stand. We used linear multilevel mixed-effects models to examine relationships with SH. Fixed and random effects (random intercept) accounted for repeated measurements within participants. Linear splines were fitted to the data with knots at 0, 10, 20, 30, and 50 s, corresponding to the phases of the response (0–10, 10–20, 20–30, 30–50, and 50–180 s). An autoregression covariance matrix with a lag of 1 was applied to model residual variance over time as time points closer together are more strongly correlated.

First, we examined the BP response (absolute and change from baseline) adjusted for age, sex, education, and height by SH severity. Next, we examined change in TSI from baseline in those with and without SH, adjusted for age, sex, education, and height. To assess CA efficiency, we additionally adjusted for beat-to-beat MAP at all time points during the stand, thus examining the TSI response standardized to MAP. This method is similar to transcranial Doppler and NIRS studies, which assess CA via a CA index, that is, a ratio of cerebral blood flow and BP.^{29,30} A fully controlled model also adjusted for smoking history, alcohol excess, stand transition time, depressive symptoms, antidepressant medications, antihypertensive medications, benzodiazepines, antipsychotics, diabetes, number of cardiovascular conditions, cancer, kidney disease, body mass index, TUG time, change in beat-to-beat cardiac output, seated SBP, and PWV. Finally, we looked at

change in TSI from baseline by SH severity adjusted for the full set of covariates.

Sensitivity Tests

We included sensitivity tests to examine the effect of cardiac output and heart rate on the TSI response. First, we assessed the cardiac output and heart rate responses after standing by SH category. We examined the effect of cardiac output on the TSI response in a model adjusting for age, sex, height, and change in cardiac output only and then additionally adjusting for MAP. Similarly, we examined the effect of heart rate on the TSI response adjusting for age, sex, height, and heart rate only and then additionally adjusting for MAP.

RESULTS

Participant Characteristics

Valid data were available for 2750 participants (Figure 1). Almost half the sample (49%) had SH. Table S1 in the [Data Supplement](#) displays the severity of SH. Those with SH were older, more likely to be women, taking antihypertensive medications, have a higher PWV, and a slower TUG. Supine TSI was not different in those with or without SH. Full characteristics by SH are shown in Table S2. Characteristics of excluded cases are shown in Table S3. Compared with the final sample, they were older, had a higher percentage of current smokers, and were more likely to rate their physical health as fair or poor.

BP Response

Those with SH maintained significantly higher absolute levels of MAP throughout the stand, graded by severity of SH (Figure 2, top). The change in MAP from baseline after standing revealed a graded pattern of impaired and slower recovery in those with SH (Figure 2, bottom). Those with SH experienced a larger initial drop after standing (at 10 s), which is also graded by severity of SH. After 180 s (3 minutes) of standing, those without SH and those with mild/moderate SH had returned to baseline MAP levels (Figure 2, bottom), but those with severe SH had not recovered to baseline level. Those without SH quickly returned to baseline (by 30 s).

Cerebral Oxygenation Response

Those with SH maintained TSI and recovered in a similar manner to those without SH (with only minor differences observed at 10- and 20-s time points; Figure 3, top). However, when the TSI response was standardized to orthostatic MAP at each time point, a graded pattern of impaired recovery emerged (Figure 3, middle), that is, when presented with the same MAP deficit (held at mean MAP values at each time point over the stand), TSI was impaired in those with SH, indicating impaired cerebral regulation. As reported above, those

with SH have differing MAP profiles, and we wanted to assess TSI independent of the MAP response. Adjusting for orthostatic MAP in this manner is conceptually similar to asking the question “What if these two groups had the same MAP response on standing, how would their TSI recover?”

After 3 minutes of standing, a deficit of -0.83% (95% CI, -0.93 to -0.74) remained in those with SH. This relationship persisted in the fully adjusted model (Figure 3, bottom). Examining the TSI response by SH severity, those with the most severe levels of SH had the largest initial drop. Overall, there is some evidence of a gradient by severity of SH, particularly in the earlier stages of the response (Figure 4).

Sensitivity Tests

Those with SH exhibited a greater increase in cardiac output upon standing but stabilized at lower levels than those without SH (Figure S1, top). There was no difference in maximum increase of heart rate upon standing between the groups; however, those with SH stabilized at lower levels than those without SH (Figure S1, bottom).

Adjusting for change in cardiac output (Figure S2) or heart rate (Figure S3) did not alter the TSI response compared with previous models (Figure 3).

DISCUSSION

This study demonstrated that in a large cohort of community-dwelling older adults, SH was associated with a larger drop and impaired recovery of cerebral oxygenation upon standing when standardized to orthostatic MAP. This suggests that CA is impaired and less efficient in those with SH. These effects persisted after adjusting for important covariates including age, cardiovascular conditions, and medications, as well as heart rate and change in cardiac output. TSI stabilized more quickly in those without SH or with mild SH than those in moderate/severe categories of SH.

In agreement with previous studies, we have shown that the orthostatic BP response is influenced by baseline BP levels,³¹ that the magnitude of the drop is related to baseline supine BP,^{32,33} and that the slowest/most impaired BP recovery is in those with the highest baseline BP.¹⁷

This is the first study to describe the cerebrovascular response to orthostasis and CA efficacy in those with SH. SH may share underlying mechanisms with essential hypertension. Previous studies have reported mixed results as to whether essential hypertension impairs CA efficacy.^{34–37} In chronic hypertension, the cerebral autoregulatory curve is moved to the right. Thus, tolerance to hypotension is impaired.³ Some people with OH have an expanded autoregulatory range and, therefore, can tolerate large changes in BP.³⁸ However, it is generally




Figure 1. Data available for analysis.
BP indicates blood pressure.

observed in those with neurogenic OH, as evidenced by severe autonomic dysfunction. Lassen first proposed a pressure-passive relationship within the cerebral autoregulatory range of ≈ 60 to 150 mmHg MAP.³⁹ However, later studies indicate a smaller plateau³ and even suggest that CBF and oxygenation are not independent of BP changes.⁴⁰

Future Problems

Poor regulation of cerebral oxygenation upon standing may increase symptoms of orthostatic intolerance, falls, or syncope in the short term, all affecting quality of life. Chronic hypotensive burden may lead to functional and structural brain changes such as increased WMH and cognitive impairment and may also be a risk for hypoperfusion of other end organs. A recent study by Palma et al¹⁶ examined longitudinal changes

in those with SH. The study reported associations with WMH in SH. However, there were no differences in cognition as measured by MOCA scores, although the study was not designed/powerd to detect cognitive changes. An earlier TILDA study reported poor cognitive performance in those who had both OH and SH, compared with those who had OH only,¹⁵ as well as cognitive decline over a 4-year period in those with SH-OH.⁸ This current study suggests that those with SH may have deficits in cerebral regulation, which in turn may be a pathway for cognitive impairment. Future work should explore these links.

Possible Mechanisms

The SH group had a significantly higher PWV (10.0 versus 10.9 m/s), indicating increased arterial stiffness. In chronic hypertension, arterial stiffness increases, thus




Figure 2. Conditional means and 95% CIs for absolute mean arterial pressure by supine hypertension (SH) category, estimated from mixed-effects models, adjusted for age, sex, education, and height.

Absolute values (top) and change from baseline (bottom) are shown. As baseline is calculated as an average of readings 60 to 30 s before standing, it may not always be at 0.

increased pulsatility in the vessels⁴¹ leads to remodeling of the vasculature and long-term changes in brain structure and function such as WMH. Evidence of remodeling in SH in the Palma study demonstrates association with increased left ventricular hypertrophy as well as WMH. In this study, we included PWV as a covariate in our fully controlled models, but this did not explain the differences in the TSI response in those with SH. We also included TUG—an indicator of sarcopenia⁴² and thus muscle pump function—as a covariate. Although the SH group had a slower TUG, this did not explain the differences we observed in the TSI response. Likewise, the heart rate response or change in cardiac output did not account for the differences we observed. In fact, the increase in cardiac output was

greater in those with SH, suggesting that it may be a sufficient response, although stabilization in the recovery phase was at a lower level.

METHODOLOGICAL CONSIDERATIONS

CA and CA efficiency are typically assessed by calculation of indexes that are standardized to BP. For example, Mol et al³⁰ defined CA as the drop in oxygenated hemoglobin divided by MAP within 60 s of postural change. However, assessment of CA based on such ratios represents a loss of information, with a reduction to a single quantitative value that does not capture the rapid, transient, or more subtle patterns of impairment in the response. Here, we balanced the use of continuous data




Figure 3. Conditional means and 95% CIs for change in tissue saturation index from baseline by supine hypertension (SH) estimated from mixed-effects models. Models are adjusted for age, sex, education, and height (top), additionally adjusted for beat-to-beat mean arterial pressure (middle), and additionally adjusted for smoking history, alcohol excess, stand transition time, depressive symptoms, antidepressant medications, antihypertensive medications, benzodiazepines, antipsychotics, diabetes, number of cardiovascular conditions, cancer, kidney disease, body mass index, timed-up-and-go time, change in beat-to-beat cardiac output, seated systolic blood pressure, and pulse wave velocity (bottom). As baseline is calculated as an average of readings 60 to 30 s before standing, it may not always be at 0.




Figure 4. Conditional means and 95% CIs for change in tissue saturation index from baseline adjusted for age, sex, education, height, beat-to-beat mean arterial pressure, smoking history, alcohol excess, stand transition time, depressive symptoms, antidepressant medications, antihypertensive medications, benzodiazepines, antipsychotics, diabetes, number of cardiovascular conditions, cancer, kidney disease, body mass index, timed-up-and-go time, change in beat-to-beat cardiac output, seated systolic blood pressure, and pulse wave velocity by severity of supine hypertension (SH) estimated from mixed-effects models.

As baseline is calculated as an average of readings 60 to 30 s before standing, it may not always be at 0.

and the practicalities of statistical comparisons by utilizing mixed-effects models to examine multiple time points over the full time course of the response. This method also allowed us to standardize the TSI response to BP and adjust for multiple covariates of interest.

Clinical Implications

We found a remaining difference of almost 1% in TSI after 3 minutes in those with SH. Differences of this magnitude have previously been shown to be associated with clinical outcomes such as depression.⁴³ Bachus et al⁴⁴ also found drops in cerebral oxygenation of $\approx 5\%$ (range, -2% to -10%) relating to syncope in tilt-table tests. Together, these suggest small differences in cerebral oxygenation are clinically relevant.

Treating those with SH is challenging when it is accompanied by OH, and clinical recommendations are to prioritize nonpharmacological management when possible.¹ Previous work reported that a large drop in BP and nonrecovery was associated with the use of β -blockers in particular, and antidepressants, as well as being a predictor of orthostatic intolerance.^{17,45} Treatment interventions such as salt-retaining fludrocortisone used to treat OH may exasperate or even cause SH.

There is some concern that lowering BP may lead to lower cerebral blood flow and oxygenation. There is evidence from smaller studies that the cerebral autoregulatory curve is restored after treatment for hypertension and cerebral hemodynamics remain intact after lowering BP.⁴⁶ A recent study of treated chronic hypertensive patients shows that some CA measures derived from transcranial Doppler and NIRS recordings (transfer function estimates of coherence, gain, phase, and transit times) were comparable to controls.⁴⁷ However, duration of hypertension may also be a factor, and cerebrovascular

resistance was increased in this cohort. Studies from the SPRINT (Systolic Blood Pressure Intervention Trial) show no increase in falls or OH prevalence after intensive lowering BP.⁴⁸ For some people, stability of BP may reduce OH. A recent SPRINT study also reported that OH was not associated with an increased risk of falls, syncope, or CVD events, even in the intensive treatment group.⁴⁹ This suggests antihypertensive treatment improves OH and the BP response, which may in turn influence the TSI response. To better understand the implications for treatment of OH and CA, it will be necessary to examine all of these components in trials and derive more precise phenotyping to guide treatments.

Current hypertension guidelines refer to seated and standing measurements for monitoring BP. However, our study highlights the importance of screening for SH, particularly in older adults, which may have additional diagnostic value in managing OH and identifying those at risk for falls. This is even more important as BP can appear lower when sitting.⁵⁰ Also, a previous study comparing seated office BP, supine BP, and ambulatory BP ($n=280$) reported that supine BP had greater sensitivity and specificity for diagnosing hypertension than seated BP,⁵¹ although 24-hour ambulatory BP monitoring is a reasonable approach for capturing SH.⁵²

Strengths and Limitations

Strengths of this study include the large sample size consisting of community-dwelling older adults and the comprehensive data on health behaviors, comorbidities, medication use, and demographics, which allowed for adjustment of potential confounders. Another strength is the noninvasive continuous assessment of both peripheral BP and cerebral oxygenation. Limitations

include the cross-sectional nature of the study. Blood hemoglobin and CO₂ were not measured due to practical limitations in such a large cohort. CO₂ vasoreactivity is reduced in individuals with hypertension.⁵³ A small reduction in CO₂ is also expected with postural change⁵⁴ contributing to a reduction in cerebral flow, evident in controls and orthostatic intolerant patients.⁵⁵ However, the relationship is complex, and cerebral CO₂ reactivity alone does not account for this drop.⁵⁶ While collection of NIRS data may be subject to interoperator, bias all TILDA health assessment data are collected by a team of rigorously trained nurses who follow a detailed protocol, minimizing such biases. Due to practicalities, participants were not asked to fast or refrain from caffeine, smoking, alcohol, or medications before the health assessment; however, this is more representative of daily living.

Conclusions

In this large population study, we determined that cerebral oxygenation and CA is impaired in those with SH, and this impairment is graded by severity of SH. The implications are that in older patients, consideration should be given to measuring SH and screening for OH. Therapeutic studies are needed to better understand the relationship between cerebral oxygenation, medications, supine BP, and OH.

Perspectives

Current routine clinical BP assessment relies on seated BP readings, but there may be benefit to screening for SH, particularly for older adults who may be at risk for OH and falls.

ARTICLE INFORMATION

Received February 1, 2021; accepted May 4, 2021.

Affiliations

The Irish Longitudinal Study on Ageing (L.N., J.D.O., R.R.-O., R.A.K.), Trinity Centre for Biomedical Engineering (R.B.R.), and Discipline of Medical Gerontology, School of Medicine (R.R.-O., R.B.R., R.A.K.), Trinity College Dublin, Ireland. Mercer's Institute for Successful Ageing, St. James' Hospital, Dublin, Ireland (R.R.-O., R.A.K.).

Acknowledgments

We would like to acknowledge the ongoing contribution of the TILDA (The Irish Longitudinal Study on Ageing) participants and research staff.

Sources of Funding

Financial support for TILDA (The Irish Longitudinal Study on Ageing) was provided by the Irish Government Department of Health, the Atlantic Philanthropies, Irish Life plc, and Science Foundation Ireland, grant number 18/FRL/6188. The funders had no involvement in the study design, collection, analysis, and interpretation of data or preparation of this article. Any views expressed in this report are not necessarily those of the Department of Health and Children or of the Minister for Health.

Disclosures

None.

REFERENCES

- Jordan J, Fanciulli A, Tank J, Calandra-Buonaura G, Cheshire WP, Cortelli P, Eschboeck S, Grassi G, Hilz MJ, Kaufmann H, et al. Management of supine hypertension in patients with neurogenic orthostatic hypotension: scientific statement of the American Autonomic Society, European Federation of Autonomic Societies, and the European Society of Hypertension. *J Hypertens*. 2019;37:1541–1546. doi: 10.1097/HJH.0000000000002078
- Chisholm P, Anpalahan M. Orthostatic hypotension: pathophysiology, assessment, treatment and the paradox of supine hypertension. *Intern Med J*. 2017;47:370–379. doi: 10.1111/imj.13171
- Willie CK, Tzeng YC, Fisher JA, Ainslie PN. Integrative regulation of human brain blood flow. *J Physiol*. 2014;592:841–859. doi: 10.1113/jphysiol.2013.268953
- Monahan KD. Effect of aging on baroreflex function in humans. *Am J Physiol Regul Integr Comp Physiol*. 2007;293:R3–R12. doi: 10.1152/ajpregu.00031.2007
- Verma AK, Garg A, Xu D, Bruner M, Fazel-Rezaei R, Blaber AP, Tavakolian K. Skeletal muscle pump drives control of cardiovascular and postural systems. *Sci Rep*. 2017;7:45301. doi: 10.1038/srep45301
- Freeman R, Wieling W, Axelrod FB, Benditt DG, Benarroch E, Biaggioni I, Cheshire WP, Chelimsky T, Cortelli P, Gibbons CH, et al. Consensus statement on the definition of orthostatic hypotension, neurally mediated syncope and the postural tachycardia syndrome. *Clin Auton Res*. 2011;21:69–72. doi: 10.1007/s10286-011-0119-5
- Finucane C, O'Connell MD, Donoghue O, Richardson K, Savva GM, Kenny RA. Impaired orthostatic blood pressure recovery is associated with unexplained and injurious falls. *J Am Geriatr Soc*. 2017;65:474–482. doi: 10.1111/jgs.14563
- McNicholas T, Tobin K, Carey D, O'Callaghan S, Kenny RA. Is baseline orthostatic hypotension associated with a decline in global cognitive performance at 4-year follow-up? Data from TILDA (The Irish Longitudinal Study on Ageing). *J Am Heart Assoc*. 2018;7:e008976. doi: 10.1161/JAHA.118.008976
- Wolters FJ, Mattace-Raso FU, Koudstaal PJ, Hofman A, Ikram MA; Heart Brain Connection Collaborative Research Group. Orthostatic hypotension and the long-term risk of dementia: a population-based study. *PLoS Med*. 2016;13:e1002143. doi: 10.1371/journal.pmed.1002143
- Briggs R, Carey D, Kennelly SP, Kenny RA. Longitudinal association between orthostatic hypotension at 30 seconds post-standing and late-life depression. *Hypertension*. 2018;71:946–954. doi: 10.1161/HYPERTENSIONAHA.117.10542
- Fedorowski A, Stavenow L, Hedblad B, Berglund G, Nilsson PM, Melander O. Orthostatic hypotension predicts all-cause mortality and coronary events in middle-aged individuals (The Malmo Preventive Project). *Eur Heart J*. 2010;31:85–91. doi: 10.1093/eurheartj/ehp329
- Okamoto LE, Gamboa A, Shibao C, Black BK, Diedrich A, Raj SR, Robertson D, Biaggioni I. Nocturnal blood pressure dipping in the hypertension of autonomic failure. *Hypertension*. 2009;53:363–369. doi: 10.1161/HYPERTENSIONAHA.108.124552
- Jordan J, Shannon JR, Pohar B, Paranjape SY, Robertson D, Robertson RM, Biaggioni I. Contrasting effects of vasodilators on blood pressure and sodium balance in the hypertension of autonomic failure. *J Am Soc Nephrol*. 1999;10:35–42.
- Yano Y, Kario K. Nocturnal blood pressure and cardiovascular disease: a review of recent advances. *Hypertens Res*. 2012;35:695–701. doi: 10.1038/hr.2012.26
- Frewen J, Finucane C, Savva GM, Boyle G, Kenny RA. Orthostatic hypotension is associated with lower cognitive performance in adults aged 50 plus with supine hypertension. *J Gerontol A Biol Sci Med Sci*. 2014;69:878–885. doi: 10.1093/gerona/glt171
- Palma JA, Redel-Traub G, Porciuncula A, Samaniego-Toro D, Millar Vernetti P, Lui YW, Norcliffe-Kaufmann L, Kaufmann H. The impact of supine hypertension on target organ damage and survival in patients with synucleinopathies and neurogenic orthostatic hypotension. *Parkinsonism Relat Disord*. 2020;75:97–104. doi: 10.1016/j.parkreldis.2020.04.011
- Romero-Ortuno R, Finucane C, O'Connell M, Kenny RA. Insights into the clinical management of the syndrome of supine hypertension-orthostatic hypotension (Sh-Oh): the Irish longitudinal study of ageing. *Age Ageing*. 2013;42:22–22.
- Cronin H, O'Regan C, Finucane C, Kearney P, Kenny RA. Health and aging: development of the Irish Longitudinal Study on Ageing health assessment. *J Am Geriatr Soc*. 2013;61(suppl 2):S269–78. doi: 10.1111/jgs.12197
- Whelan BJ, Savva GM. Design and methodology of the Irish longitudinal study on ageing. *J Am Geriatr Soc*. 2013;61(suppl 2):S265–8. doi: 10.1111/jgs.12199

20. Jasper HH. The ten twenty electrode system of the international federation. *Electroencephalogr Clin Neurophysiol*. 1958;10:371–375.
21. Ferrari M, Quaresima V. Near infrared brain and muscle oximetry: from the discovery to current applications. *J Near Infra Spectr*. 2012;20:1–14. doi: 10.1255/jnirs.973
22. Ferrari M, Quaresima V. A brief review on the history of human functional near-infrared spectroscopy (fNIRS) development and fields of application. *Neuroimage*. 2012;63:921–935. doi: 10.1016/j.neuroimage.2012.03.049
23. FMS. *Finometer User's Guide*. Finapres Medical Systems BV; 2005.
24. Guelen I, Westerhof BE, Van Der Sar GL, Van Montfrans GA, Kiemeneij F, Wesseling KH, Bos WJ. Finometer, finger pressure measurements with the possibility to reconstruct brachial pressure. *Blood Press Monit*. 2003;8:27–30. doi: 10.1097/00126097-200302000-00006
25. O'Connor JD, O'Connell MDL, Nolan H, Newman L, Knight SP, Kenny RA. Impact of standing speed on the peripheral and central hemodynamic response to orthostasis. *Hypertension*. 2020;75:524–531. doi: 10.1161/HYPERTENSIONAHA.119.14040
26. Fanciulli AI, Jordan J, Biaggioni I, Calandra-Buonaura G, Cheshire WP, Cortelli P, Eschlboeck S, Grassi G, Hilz MJ, Kaufmann H, et al. Consensus statement on the definition of neurogenic supine hypertension in cardiovascular autonomic failure by the American Autonomic Society (AAS) and the European Federation of Autonomic Societies (EFAS): endorsed by the European Academy of Neurology (EAN) and the European Society of Hypertension (ESH). *Clin Auton Res*. 2018;28:355–362. doi: 10.1007/s10286-018-0529-8
27. Bush B, Shaw S, Cleary P, Delbanco TL, Aronson MD. Screening for alcohol abuse using the CAGE questionnaire. *Am J Med*. 1987;82:231–235. doi: 10.1016/0002-9343(87)90061-1
28. O'Halloran AM, Kenny RA, King-Kallimanis BL. The latent factors of depression from the short forms of the CES-D are consistent, reliable and valid in community-living older adults. *Euro Geriatr Med*. 2014;5:97–102. doi: 10.1016/j.eurger.2013.12.004
29. Panerai RB. Transcranial Doppler for evaluation of cerebral autoregulation. *Clin Auton Res*. 2009;19:197–211. doi: 10.1007/s10286-009-0011-8
30. Mol A, Maier AB, van Wezel RJA, Meskers CGM. Multimodal monitoring of cardiovascular responses to postural changes. *Front Physiol*. 2020;11:168. doi: 10.3389/fphys.2020.00168
31. Imholz BP, Dambrink JH, Karemaker JM, Wieling W. Orthostatic circulatory control in the elderly evaluated by non-invasive continuous blood pressure measurement. *Clin Sci (Lond)*. 1990;79:73–79. doi: 10.1042/cs0790073
32. van Dijk JG, Tjon-A-Tsien AM, Kamzoul BA, Kramer CG, Lemkes HH. Effects of supine blood pressure on interpretation of standing up test in 500 patients with diabetes mellitus. *J Auton Nerv Syst*. 1994;47:23–31. doi: 10.1016/0165-1838(94)90062-0
33. Fedorowski A, Burri P, Melander O. Orthostatic hypotension in genetically related hypertensive and normotensive individuals. *J Hypertens*. 2009;27:976–982. doi: 10.1097/hjh.0b013e3283279860
34. Eames PJ, Blake MJ, Panerai RB, Potter JF. Cerebral autoregulation indices are unimpaired by hypertension in middle aged and older people. *Am J Hypertens*. 2003;16(9 pt 1):746–753. doi: 10.1016/s0895-7061(03)00947-6
35. Lipsitz LA, Mukai S, Hamner J, Gagnon M, Babikian V. Dynamic regulation of middle cerebral artery blood flow velocity in aging and hypertension. *Stroke*. 2000;31:1897–1903. doi: 10.1161/01.str.31.8.1897
36. Serrador JM, Sorond FA, Vyas M, Gagnon M, Iloputaife ID, Lipsitz LA. Cerebral pressure-flow relations in hypertensive elderly humans: transfer gain in different frequency domains. *J Appl Physiol (1985)*. 2005;98:151–9. doi: 10.1152/jappphysiol.00471.2004
37. Traon AP, Costes-Salon MC, Galinier M, Fourcade J, Larrue V. Dynamics of cerebral blood flow autoregulation in hypertensive patients. *J Neurol Sci*. 2002;195:139–144. doi: 10.1016/s0022-510x(02)00010-2
38. Novak V, Novak P, Spies JM, Low PA. Autoregulation of cerebral blood flow in orthostatic hypotension. *Stroke*. 1998;29:104–111. doi: 10.1161/01.STR.29.1.104
39. LASSER NA. Cerebral blood flow and oxygen consumption in man. *Physiol Rev*. 1959;39:183–238. doi: 10.1152/physrev.1959.39.2.183
40. Lucas SJ, Tzeng YC, Galvin SD, Thomas KN, Ogoh S, Ainslie PN. Influence of changes in blood pressure on cerebral perfusion and oxygenation. *Hypertension*. 2010;55:698–705. doi: 10.1161/HYPERTENSIONAHA.109.146290
41. Najjar SS, Scuteri A, Shetty V, Wright JG, Muller DC, Fleg JL, Spurgeon HP, Ferrucci L, Lakatta EG. Pulse wave velocity is an independent predictor of the longitudinal increase in systolic blood pressure and of incident hypertension in the Baltimore Longitudinal Study of Aging. *J Am Coll Cardiol*. 2008;51:1377–1383. doi: 10.1016/j.jacc.2007.10.065
42. Martinez BP, Gomes IB, Oliveira CS, Ramos IR, Rocha MD, Forgiarini Júnior LA, Camelier FW, Camelier AA. Accuracy of the Timed Up and Go test for predicting sarcopenia in elderly hospitalized patients. *Clinics (Sao Paulo)*. 2015;70:369–372. doi: 10.6061/clinics/2015(05)11
43. Briggs R, Carey D, Claffey P, McNicholas T, Newman L, Nolan H, Kennelly SP, Kenny RA. The association between frontal lobe perfusion and depressive symptoms in later life. *Br J Psychiatry*. 2019;214:230–236. doi: 10.1192/bjp.2018.288
44. Bachus E, Holm H, Hamrefors V, Melander O, Sutton R, Magnusson M, Fedorowski A. Monitoring of cerebral oximetry during head-up tilt test in adults with history of syncope and orthostatic intolerance. *Europace*. 2017;20:1535–1542. doi: 10.1093/europace/eux298
45. Canney M, O'Connell MD, Murphy CM, O'Leary N, Little MA, O'Seaghdha CM, Kenny RA. Single agent antihypertensive therapy and orthostatic blood pressure behaviour in older adults using beat-to-beat measurements: the Irish longitudinal study on ageing. *PLoS One*. 2016;11:e0146156. doi: 10.1371/journal.pone.0146156
46. Zhang R, Witkowski S, Fu Q, Claassen JA, Levine BD. Cerebral hemodynamics after short- and long-term reduction in blood pressure in mild and moderate hypertension. *Hypertension*. 2007;49:1149–1155. doi: 10.1161/HYPERTENSIONAHA.106.084939
47. Müller M, Österreich M, Lakatos L, Hessling Av. Cerebral macro- and micro-circulatory blood flow dynamics in successfully treated chronic hypertensive patients with and without white matter lesions. *Sci Rep*. 2020;10:9213. doi: 10.1038/s41598-020-66317-x
48. Wright JT Jr, Williamson JD, Whelton PK, Snyder JK, Sink KM, Rocco MV, Reboussin DM, Rahman M, Oparil S, Lewis CE, et al; SPRINT Research Group. A randomized trial of intensive versus standard blood-pressure control. *N Engl J Med*. 2015;373:2103–2116. doi: 10.1056/NEJMoA1511939
49. Juraschek SP, Taylor AA, Wright JT Jr, Evans GW, Miller ER III, Plante TB, Cushman WC, Gure TR, Haley WE, Moinuddin I, et al; SPRINT Research Group. Orthostatic hypotension, cardiovascular outcomes, and adverse events: results from SPRINT. *Hypertension*. 2020;75:660–667. doi: 10.1161/HYPERTENSIONAHA.119.14309
50. Cicolini G, Pizzi C, Palma E, Bucci M, Schioppa F, Mezzetti A, Manzoli L. Differences in blood pressure by body position (supine, Fowler's, and sitting) in hypertensive subjects. *Am J Hypertens*. 2011;24:1073–1079. doi: 10.1038/ajh.2011.106
51. Krzesiński P, Stańczyk A, Gielera G, Piotrowicz K, Banak M, Wójcik A. The diagnostic value of supine blood pressure in hypertension. *Arch Med Sci*. 2016;12:310–8. doi: 10.5114/aoms.2016.59256
52. Ghazi L, Drawz PE, Pajewski NM, Juraschek SP. The association of orthostatic hypotension with ambulatory blood pressure phenotypes in SPRINT. *Am J Hypertens*. 2020. doi: 10.1093/ajh/hpaa184
53. Lavi S, Gaitini D, Milloul V, Jacob G. Impaired cerebral CO₂ vasoreactivity: association with endothelial dysfunction. *Am J Physiol Heart Circ Physiol*. 2006;291:H1856–H1861. doi: 10.1152/ajpheart.00014.2006
54. Sorond FA, Serrador JM, Jones RN, Shaffer ML, Lipsitz LA. The sit-to-stand technique for the measurement of dynamic cerebral autoregulation. *Ultrasound Med Biol*. 2009;35:21–29. doi: 10.1016/j.ultrasmedbio.2008.08.001
55. Novak V, Spies JM, Novak P, McPhee BR, Rummans TA, Low PA. Hypocapnia and cerebral hypoperfusion in orthostatic intolerance. *Stroke*. 1998;29:1876–1881. doi: 10.1161/01.str.29.9.1876
56. Immink RV, Secher NH, Roos CM, Pott F, Madsen PL, van Lieshout JJ. The postural reduction in middle cerebral artery blood velocity is not explained by PaCO₂. *Eur J Appl Physiol*. 2006;96:609–614. doi: 10.1007/s00421-006-0136-6