

Paper published as:

Senge, M. O.; Shaker, Y. M.; Pintea, M.; Ryppa, C.; Hatscher, S. S.; Ryan, A.; Sergeeva, Y. (2010):
Synthesis of meso-Substituted ABCD-type Porphyrins via Functionalization
Reactions.
European Journal of Organic Chemistry, 237–258.

Synthesis of meso-Substituted ABCD-type Porphyrins via Functionalization Reactions

Mathias O. Senge,^{*,[a,b]} Yasser M. Shaker,^[a] Monica Pinteá,^[a] Claudia Ryppa,^[c] Sabine S. Hatscher,^[c] Aoife Ryan^[a] and Yulia Sergeeva^[a]

Keywords: Porphyrinoids / C–C coupling / Organolithium / Tetrapyrroles / Conformation analysis

Considerable progress has been made in recent years in the search for synthetic methods leading to functionalized porphyrins, esp. for modification of either the β - or meso positions. For the latter, total synthesis based on condensation methods or partial synthesis through functionalization of preformed porphyrin have emerged as possible methods. The increasing number of possible technical and medicinal applications for unsymmetrically meso substituted porphyrins requires straightforward methods for the preparation of the so-called ABCD-porphyrins, i. e., porphyrins with up to four different meso substituents. Here, we describe new strategies for the synthesis of ABCD-type porphyrins based on porphyrin reactions with organolithium reagents and the use of Pd-catalyzed coupling reactions. Using the whole repertoire of contemporary functionalization methods a comprehensive analysis and comparison of the various strategies for A-, AB-, A₂B-, ABC-,


A₂BC- and ABCD-type porphyrins is given. In addition we report on the synthesis of new functionalized derivatives for some of these porphyrin classes. In practical terms and taking an applied science-oriented approach the synthesis of unsymmetrically meso substituted porphyrins is best accomplished by a combination of well-developed condensation methods with subsequent functionalization via organolithium compounds or transition metal catalyzed coupling protocols. The methods described are suitable for the preparation of porphyrins for many divergent applications ranging from amphiphilic porphyrins for photodynamic therapy, push-pull systems for optical applications, chiral systems useful in catalysis to donor-acceptor systems suitable for electron transfer studies. (© WILEY-VCH Verlag GmbH & Co. KGaA, 69451 Weinheim, Germany, 2009)

- [a] School of Chemistry, SFI Tetrapyrrole Laboratory, Trinity College Dublin, Dublin 2, Ireland
Fax: +353-1-896-8536
E-mail: sengem@tcd.ie
- [b] Medicinal Chemistry, Institute of Molecular Medicine, Trinity Centre for Health Sciences, Trinity College Dublin, St James's Hospital, Dublin 8, Ireland
- [c] Institut für Chemie, Universität Potsdam, Karl-Liebknecht-Str. 24-25, D-14476 Golm, Germany

Introduction

Porphyrins are a unique class of compounds that are ubiquitous in nature and function in a wide variety of roles ranging from oxygen transport, electron transfer and oxidation catalysts to photosynthesis. Thus, these widely distributed and important tetrapyrrolic cofactors play crucial roles in many biochemical and technical processes. These range from impaired biosynthesis (porphyrias), photodynamic cancer therapy (PDT), malaria, neuropsychiatric disorders to industrial applications as sensors, optical materials, oxidation catalysis and in supramolecular chemistry. Overall, porphyrins are probably the most widely studied class of natural products whose studies have impacted almost any modern scientific field.^[1] From a chemical viewpoint, their importance is related to their chemical properties, i.e., their photochemical (energy and exciton transfer), redox (electron transfer, catalysis), and coordination properties (metal and axial ligand binding), and their conformational flexibility (functional control).^[2]


The various applications and their biochemical relevance are related by a common denominator in the underlying chemistry of these compounds. Current applications require the synthesis of unsymmetrically substituted porphyrins, i.e. systems with different substituents in a regiochemically defined manner. Next to the naturally occurring β -substituted porphyrins (e.g., the chlorophylls and hemes) typical examples are the so-called ABCD-porphyrins (**1**), where four different meso substituents are present (**2**).^[3] The type and arrangement of these substituents then defines different applications. To name a few, residues with functional groups suitable for small molecule binding may be used for catalysis, a mixture of polar and nonpolar residues yields amphiphilic porphyrins suitable for photodynamic therapy and a mix of electron donating and withdrawing groups will give push-pull porphyrins for applications in nonlinear optics.^[4]



Future progress in this area obviously depends on the synthetic accessibility of porphyrins. While – theoretically – almost any

desired β -substituted porphyrin can be prepared based on historical contributions from eminent chemists such as Willstätter, Fischer, Woodward, Eschenmoser and Smith^[5a] there were still limitations in the preparation of meso substituted porphyrins. Due to their symmetry and simple preparation most of the applications currently under study use 5,10,15,20-tetrasubstituted porphyrins (**3**) and a significant body of methods has been developed for the modification of the β -positions in such systems.^[5b]

In principle, there are three ways to access the more interesting unsymmetrical ABCD-type compounds (Scheme 1). These are the disconnection into a bilane **5** or any combination of pyrrole building blocks **6** that can be used in [2+2] or [3+1] condensation reactions.^[6] Theoretically possible is also a mixed condensation using pyrrole **4** and various aldehydes which is still used by many researchers for systems with lower symmetry. Here, the number of regioisomers formed is too large and the necessary purification and separation workup is too cumbersome if possible at all. Note, that most of these reactions involve acid-catalyzed condensation reactions, often resulting in significant scrambling of the pyrrole units thus limiting the type of substituents that can be used.



Scheme 1. Retrosynthetic analysis of ABCD-type porphyrins.

The most significant contributions in this area of ABCD-porphyrin "total synthesis" were made by the group of Prof. Lindsey.^[6b] Examples for new developments are the synthesis of 1-bromo-19-acylbilanes by acid-catalyzed condensation of 1-acyldipyrromethanes and 9-bromodipyrromethane-1-carbinols followed by intramolecular cyclization of the 1-bromo-19-acylbilanes in the presence of a metal salt in a noncoordinating solvent and the synthesis of 1-protected 19-acylbilanes by acid-catalyzed condensation 1-acyldipyrromethanes and 9-protected dipyrromethane-1-carbinols with subsequent transformation of the 1-protected 19-acylbilane under basic, metal-templating conditions to give the corresponding metalloporphyrins.^[7]

We have taken an alternative approach focusing on partial synthesis starting with preformed porphyrins, i.e. envisioning a


step-wise introduction of the four meso substituents in **2**. The last decades have seen a surge in novel functionalization reactions of porphyrins, many based on C-C coupling reactions which has put this approach within possibility.^[8] By now many A₂BC- and A₃B-type porphyrins have been prepared starting with the easily accessible 5,15-disubstituted porphyrins^[3,9-12] and from brominated precursor compounds often based on Heck-type reactions. In this context we have developed the use of organolithium reagents for the synthesis of various tetrapyrrole classes.^[13-16]

In a strategic sense the preparation of ABCD-type porphyrins via functionalization requires the use of the less substituted precursors, e.g. **7**, and ultimately use of unsubstituted porphyrin **8**. The latter then allows for preparation of a mono-substituted A-type porphyrin (**7**, $\text{R}^2 = \text{R}^3 = \text{H}$) followed by subsequent regiochemical introductions of the B-, C- and D-type residues. With recent advances in the synthesis of **8**^[17] and the preparation of A-type porphyrins through condensation and functionalization reactions^[18] all necessary parts for partial synthesis of ABCD porphyrins are now in hand. Based on our current interest in the application of porphyrins for photodynamic cancer therapy (PDT) and optical applications^[19] we here present a comprehensive analysis of the application of the currently available synthetic strategies for the meso functionalization of porphyrins. This study aims to evaluate the applicability and limitations of these methods for the preparation of ABCD-type porphyrins in general. We also report on the synthesis of new functionalized derivatives for some of the different ABCD porphyrin classes.

Results and Discussion

A preparation of ABCD-type porphyrins through functionalization requires a step-wise approach, i.e., starting from A-porphyrins and through sequential introduction of more residues going to the ABCD-type systems. Thus, for practical purposes, we start in our discussion with the least substituted systems.

Synthesis of A-type porphyrins




These porphyrins are accessible either via substitution of porphyrin ("porphine") with RLi or via mixed condensation.^[18] One example for the latter is the mixed condensation of dipyrromethane, 2-hydromethylpyrrole and methyl 4-formylbenzoate to give **9** in 8 % yield. Another example is the preparation of compound **13** by condensation of the 5-substituted

dipyrromethane **11** (derived from the aldehyde **10**) with dipyrromethane **12** and trimethyl orthoformate in 4 % yield. Typically such condensations give 2-12 % yield and the formation of the A-type porphyrin is accompanied by the the respective 5,15-A₂-type system, which can be easily removed by chromatography.^[18] Preparation from *in situ* generated porphyrin and reaction with RLi is attractive but works well best for reactions involving ~100 mg of starting material. Large scale syntheses are still best performed via mixed condensations.


Synthesis of A₂-type porphyrins

The synthesis of 5,15-disubstituted A₂-type porphyrins is well established by now. Their preparation is easily accomplished by a [2+2] condensation using dipyrromethane **12** and the respective aldehyde carrying the meso substituent.^[3,9-11,20] An example for the synthesis of a 5,15-disubstituted porphyrin with new functional groups is the synthesis of the bisdioxaborolan porphyrin **14** from dipyrromethane and the respective aldehyde in 58 % yield. Metallation with nickel acetylacetonate gave the respective metal complex **15**. The related 5,10-disubstituted derivatives are accessible via similar methods but typically give much lower yields.^[21] Here compound **16** could be prepared in very low yield (0.4%). Total syntheses have been reported for e.g. 5,10-diphenylporphyrin.^[22] 5,15-Disubstituted porphyrins are also often derived from mixed condensation such as AB-type porphyrin syntheses (*vide infra*).




Synthesis of AB-type porphyrins

Two possibilities exist for the preparation of 5,15-AB porphyrins. One involves classic mixed condensation reactions. In practical terms this is still a facile reaction and for certain residues allows quick generation of the desired compounds and can even be controlled to give only the target compound as the main product.^[11,23] In simple approaches, often product mixtures are encountered. Nevertheless, the different polarities mostly allow for a quick chromatographic separation of the compounds. E.g., the



Scheme 2. Synthesis of AB-porphyrins via condensation reactions.

More elegant might be a synthesis using C–C coupling reactions. The use of Heck-type coupling reactions is still problematic due to limited access to the necessary monobromo derivatives.^[24] An alternative might be the use of reactions with LiR. However, initially we were not too convinced, as studies with 2,3,7,8,12,13,17,18-octaethylporphyrin and other systems had shown a preference for formation of the 5,10-AB derivatives over the 5,15-AB regioisomers.^[13–16] Nevertheless, we attempted some reactions of 5-substituted aryl and alkylporphyrins with LiR reagents under our standard conditions (Scheme 3).




The ethylpropyl porphyrin **33** gave quite satisfactory yields with hexyl (59%) and phenyllithium (69%) but showed a clear preference for the formation of the 5,10-regioisomer (**42** and **43**, respectively). We tried to optimize the synthesis of **42** by variation of the number of equivalents of hexyllithium. Depending on the number of equivalents used (1.5, 2.5, 3.5, 4.5, 5.5) the yields varied over a broad range (42, 52, 59, 29, 44 %, respectively) with 3.5 equiv. being the optimum which makes sense as 2 equiv. will be required to neutralize the N-H. Use of the aryllithium reagent 4-dimethylaminophenyllithium gave a mixture of the two regioisomers **44** and **45** accompanied by the higher substituted porphyrins **46** and **47**. Similarly, in the case of the 5-arylporphyrins **34** and **35** the formation of both regioisomers was observed with a clear preference for the 5,10 isomer.

Clearly, the reaction of porphyrins with RLi for the preparation of AB-type porphyrins is difficult to control and offers advantages only for very specific cases. Note, that for the reaction of 5-alkylporphyrins with RLi the yields are comparable to those obtained from mixed condensations. Here, the latter offer the advantage of larger scale and easier chromatographic separation. The main benefit of the present method lies in an easier access to the otherwise inaccessible 5,10-regioisomer. Alternatively, use of an excess of alkyllithium reagent offers an entry into A₂B porphyrins from the monosubstituted precursors. E.g., reaction of **36** with hexyllithium gave the trisubstituted porphyrin **55** in 54 % yield. A similar reactivity was observed for **37**, although in this case only the 5,15-A₂B porphyrin **57** could be isolated from the product mixture.



5,15-A₂-type porphyrins are conveniently prepared from aldehydes and dipyrromethanes and present useful starting materials for the investigation of reactions at the meso position.^[9] Using S_NAr reactions they can easily be converted into the respective A₂B-porphyrins without any regiochemical problems.^[13b,15] For example, using **58**^[11] as starting material the two porphyrins **59** and **60** could be prepared easily in 71 and 77 % yield, respectively.




Synthesis of ABC-type porphyrins

First we investigated the reaction of 5,15-AB-type porphyrins with RLi reagents to yield ABC-porphyrins (Scheme 5). Similar to the situation for A₂B porphyrins only one product can be formed and the reactions typically proceeded in very good yields. Using various alkyl and aryllithium reagents the ABC porphyrins **61–64** and **67–71** were obtained in yields ranging from 47 to 90 %. Some of these free bases were converted into the respective manganese(III) complexes for later oxidation catalyst studies.^[25] As one example for possible applications we prepared the 5,10-meso-meso-strapped porphyrin derivative **72** in good yields


through reaction of **20** with 2-methoxyphenyllithium, deprotection of the methyl ether with BBr₃ followed by reaction with 1,12-dibromododecane.

Obviously, 5,10-type AB porphyrins can be used for the preparation of ABC-porphyrins as well. A comparison of the reactivity of **44** and **45** shows that the 5,10-AB porphyrins do react but give lower yields in line with the stability of the intermediate anion.^[16] Note, that such porphyrins could also be derived from the attempted syntheses of ABCD-type porphyrins through reaction of AB-porphyrins with R¹Li/R²I (*vide infra*).



Scheme 5. Synthesis of ABC-porphyrins via S_NAr.

An alternative approach to ABC porphyrins is the use of AB-type systems, attempt a selective monobromination of one meso position and then use Pd-catalyzed reactions for introduction of the C-residue. Both iodination and bromination reactions have been applied to the meso position of porphyrins.^[26] In order to access the monobromo derivatives of the AB-type porphyrins we used 1.1 eq. of NBS for the bromination. Nevertheless, as shown in Scheme 6 the reaction is difficult to control and both the mono- and dibrominated derivatives were formed roughly in a 1:1 ratio with overall yields of 80-90 %. However, their polarities are quite different and they are easily separated by column chromatography.



Scheme 6. Synthesis of ABC-porphyrins via Suzuki reactions.

Pd-catalyzed reactions have found widespread use in porphyrin chemistry. These coupling reaction offers several advantages: availability of the reagents, mild reaction conditions, tolerance for


a broad range of functional groups, small amount of catalysts and application in one pot synthesis and appropriate conditions have been developed for porphyrins.^[27]

4-Carboxymethylphenyl boronic acid was reacted with porphyrins **22** and **31** and afforded the desired products **82** and **86** in 87 % and 89% yield, respectively. Likewise, the styrene type derivative **84** was prepared in 46 % yield. Reaction of **31** with 3-nitrophenylboronic acid gave **85** in 82 % yield. Reaction of the same boronic acid with porphyrin **27** yielded product **83** in 46 % yield. Note, that 3-nitrophenyl residues in AB-diarylporphyrins have recently been identified by Banfi et al. as having potential for PDT and thus appropriate systems were ABCD-type targets for us.^[28] Note, that the 5,15-dibromo derivatives (**77,79,81**) offer an additional entry into 5,15-A₂BC compounds, which show promise as push-pull systems for optical applications.^[19b]

Synthesis of A₂BC-type porphyrins


Use of A₂B porphyrins as starting materials offers again two possibilities: Pd-type couplings of the bromo derivatives and direct substitution with LiR reagents. The latter strategy has been employed quite successfully before.^[11,12,14]

Reaction of A₂B porphyrins with RLi: In extension of our earlier studies we investigated some additional reagents and A₂B-type porphyrins (Scheme 7). While porphyrin **90** reacted in good to excellent yields with various alkyl and aryllithium reagents attempts to use RLi with electron withdrawing groups such as *p*-nitro- and *p*-bromophenyl lithium only recovered starting material. Likewise, *p*-methoxy lithium or use of the *in situ* prepared lithiated ester of ethyl-4-bromobutyrate gave no reaction. Similar results were found for porphyrin **89** and additionally, we could show that reaction with 4-ethynylphenyl lithium gave the porphyrin **91** in low yield.




Scheme 7. Synthesis of A₂BC-porphyrins via S_NAr.

Porphyrins such as **89** bearing a *p*-ethynyl group have potential uses in superstructured materials, porphyrin arrays, light harvesting systems and porphyrin based materials such as push-pull porphyrins *via* Glaser or Heck coupling.^[29,30] For example, reaction of **94** under Glaser conditions gave the bisporphyrins **95** in excellent yield (Scheme 8). Couplings of compounds such as **94** and *trans*-diiodides could give new unsymmetrical, linear trisubstituted porphyrins which may be used as model compounds for investigations of larger array systems.



Scheme 8. Glaser coupling of **94** to **95**.


The studies shown in Scheme 7 were then extended to the utilization of meso-trisubstituted porphyrin **88** to carry different alkyl residues in the “C” position using butyllithium reagents with an increasing degree in steric demand. Reactions of *n*-, *sec*-, and *tert*-butyllithium reagents with **88** were utilized to study the scope of their reactivity towards the meso position under the same reaction conditions. *n*- And *sec*-butyllithium reacted in a similar manner to form the unsymmetrical porphyrins **90**^[11] and **91** in yields of 31 and 25 %, respectively. Reaction with the more hindered *tert*-butyllithium proceeded in a different manner although the attack was still directed to the more reactive meso position. Here, phlorin **96** was formed in 20 % yield instead of porphyrin **92**. The NMR data are typical for a nonaromatic conjugated system with pyrrolic signals in the $\delta = 6\text{--}7$ ppm range. The formation of phlorin **96** indicates that the free meso position in **88** is more reactive towards *t*BuLi than β -positions as no chlorin was separated in this case. A similar behavior was found for **59** which gave the phlorin **98** in 28 %. These results are in line with results described by Krattinger and Callot.^[32] The phlorins are stable against oxidants and this reaction opened a route for the conversion of free base porphyrins to phlorins with a variety of different meso substituents.^[33]



Next we investigated the reactivity of A₂B porphyrins with aryl groups in the B position. Typically, attacking a meso position opposite to one carrying an aryl group is difficult due to steric hindrance of the mesomeric benzylic anion stabilization.^[16] While the synthesis of some A₂BC-type porphyrins by this method was successful the yields obtained of these A₂BC-porphyrins were lower than those of A₂BC-porphyrins formed when B was an alkyl group. For example, porphyrin **60** did not react with *p*-ethynylphenyllithium or *n*-butyllithium. Likewise, the use of


electron rich porphyrins gave mixed results. For example, 5-(4-aminophenyl)-10,20-diphenylporphyrin reacted readily with 4-dimethylaminophenyl or butyllithium^[11] while 5-(4-aminophenyl)-10,20-bis(3-methoxyphenyl)porphyrin reacted only with butyllithium^[11] but not with *p*-hydroxyphenyllithium or 2-methoxyphenyllithium.

Reaction of A₂ porphyrins with R¹Li/R²I: Earlier we had shown that the nickel^{II} complexes of A₂ porphyrins can be accomplished through the addition of electrophilic reagents such as alkyl iodide “after the hydrolysis step” of the RLi reaction followed by oxidation with atmospheric oxygen permits the preparation of functionalized unsymmetric tetrasubstituted porphyrins.^[14] This method was then extended to the respective free base porphyrins.^[12] While the yields were low to moderate in the range of 20 to 45 %, the method is still advantageous when compared to related mixed condensations or multi-step syntheses. Attempts to extend the reaction to 5,15-dialkylporphyrins confirmed these limitations (Scheme 9).



Scheme 9. Synthesis of A₂BC-porphyrins via anion trapping.

Porphyrin **98** was formed in only 8 % yield, while **99** and **100** were obtained in 18 and 21 %, respectively. In all three cases the target compounds were formed as the minor product. The main products (24–28 %) were the respective trisubstituted A₂B porphyrins formed by substitution with only R² from the organolithium reagent. A similar reaction was observed for reaction of **18** with 4-aminophenyllithium and propyl iodide. Here the A₂BC porphyrin **101** was formed in 20 % yield accompanied by porphyrin **102** in 16 % yield. Clearly, this reaction is only useful in certain cases.




Scheme 10. Synthesis of an 5,10-A₂BC-porphyrin via anion trapping.

Use of an 5,15-AB-type porphyrin also allowed the formation of an 5,10-A₂BC-type porphyrin in low yield. Here, reaction of hexyllithium at the meso position and the carbonyl position of **17** followed by reaction with propyl iodide gave the porphyrin **103** (18 %) accompanied by **104** (14 %) (Scheme 10).

Synthesis of ABCD-type porphyrins

Based on the outline given above three approaches were investigated. These include one-pot anion trapping reactions, sequential RLi reactions and Pd-catalyzed reactions of bromoporphyrins.

One pot reactions: First we investigated the synthesis of ABCD-type porphyrins via a one pot reaction. As described, this was based on the reaction of porphyrins with RLi under formation of an anionic intermediate^[16] that can be trapped with electrophiles followed by oxidation.^[12,14] For example, the 5,15-disubstituted AB-type porphyrin (5-hexyl-15-(4-methoxyphenyl)-porphyrin) **51** was reacted with phenyllithium in dry THF under formation of an anion that could be trapped with alkyl iodides (*n*-propyl iodide or *n*-pentyl iodide) to yield the tetra-meso-substituted free base ABCD-porphyrins **105** and **106** in yields of 25 and 14 %, respectively (Scheme 11). In a similar manner, reaction of **51** with *n*-butyllithium/*n*-propyl iodide gave porphyrin **108** and reaction with *n*-butyllithium/*n*-pentyl iodide gave compound **107** in 13 and 18 % yield, respectively (Scheme 11).




Scheme 11. Reactions of **51** with R¹Li/R²I combinations.

Typically, three to five equivalents of the organolithium reagents were used and more than ten equivalents of alkyl iodides. Complete trapping of the anionic complex formed with electrophiles required long reaction times under heating (12-24 h at 70 °C) after addition of RI. Nevertheless, the reaction appears difficult to control. Reaction of **51** with *sec*-butyllithium/*n*-propyl iodide or with *sec*-butyllithium/*n*-pentyl iodide in dry THF and subsequently hydrolysis with water and oxidation with DDQ yielded the new free base ABC-porphyrin **111** in 14 % yield without formation of the meso tetrasubstituted porphyrin. Similarly, **51** reacted with phenyllithium in dry THF followed by addition of 3-iodopropionic acid or with phenyllithium followed by addition of 3-(iodomethyl)pyridine hydrogen iodide under the same conditions

to the ABC-porphyrin **110** in 45 % yield, again without any formation of ABCD-porphyrins. Mechanistically the formation of the ABC-porphyrins can be explained by a shift in the equilibrium from the carbanionic intermediate to a phlorin type intermediate. This is more easily hydrolyzed and oxidized to the ABC-porphyrin than undergoing an electrophilic attack.


Treatment of **51** with *p*-hydroxyphenyllithium/*n*-pentyl iodide or with *p*-hydroxyphenyllithium/*n*-propyl gave no reaction. In contrast reaction with phenyllithium/5-iodouracil (dissolved in 2 ml DMF) or with *p*-hydroxyphenyllithium/3-(iodomethyl)pyridine hydrogen iodide in dry THF and boiling the reaction mixture resulted in the formation of polar blue and black materials, indicating ring-opening reactions.

Similar observations were made with porphyrin **112**. For example, reaction with *n*-butyllithium/*n*-propyl iodide or with *n*-butyllithium/*n*-pentyl iodide gave only the ABC-porphyrin **113** in 15 % yield. Various other combinations of organolithium reagents and alkyl, aryl or heterocyclic iodides together with a wide variety of AB-porphyrins with different functional groups were tested for the synthesis of ABCD-porphyrins but failed to do so. In some cases this is probably due to the inability of the *in situ* formed ABC-porphyrin intermediate to react with the electrophilic iodide reagents.




Sequential reactions with RLi: Next, we attempted the preparation of ABCD-type porphyrins via two-step synthesis through functionalization of the respective AB-porphyrins by two S_NAr reactions with organolithium reagents, first to the ABC-porphyrins (see above) and then to ABCD-type porphyrins. Target compounds were biologically relevant ABCD-porphyrins carrying both precursors for *p*-hydroxyphenyl groups and various alkyl chains.

Here, porphyrin **51** was reacted to the ABC-porphyrin **114** via addition of *n*-butyllithium. The compound was not characterized but added directly to a solution of *p*-hydroxyphenyllithium followed by addition of water and DDQ to form the ABCD-porphyrin 5-butyl-10-hexyl-15-*p*-hydroxyphenyl-20-(4-methoxyphenyl)-porphyrin **115** in 60 % yield (Scheme 12). Using a similar sequence with porphyrin **112**, PhLi and 4-hydroxyphenyllithium, porphyrin **117** was obtained in 37% yield. Similar results were obtained when using ABC-type porphyrins for conversion to the respective ABCD-type systems (see below). The results indicate that the two-step synthesis might be the method of choice for the preparation of functionalized ABCD-porphyrin in acceptable yields as the complete synthetic pathway gave higher overall yields compared to the one-pot procedure.



Scheme 12. Transformation of AB-porphyrins to ABCD-porphyrins.


Reaction of ABC porphyrins with RLi: Obviously another possibility is the use of ABC-type porphyrins for reactions with RLi, formally as the last step in a sequence of four reactions starting with porphyrin to yield ABCD porphyrins. Scheme 13 outlines some of the results. For example, preparation of the ABCD-type porphyrin **118** was possible through reaction of **62** with 4-methoxyphenyllithium in 40 % yield. In a similar manner the porphyrins **119** to **122** were accessible in varying yields. In order to prevent β -addition reactions the number of RLi equivalents has to be kept at a minimum for the conversion $\text{ABC} \rightarrow \text{ABCD}$. To some extent, this accounts for the relatively low yields encountered in these reactions.



Scheme 13. Transformation of ABC-porphyrins to ABCD-porphyrins via $\text{S}_{\text{N}}\text{Ar}$.

Nevertheless, these reactions show that, in principle, it is possible to sequentially substitute porphyrin with four different residues using RLi reagents. Admittedly, this method works much better, i.e. with higher yields, for the more soluble octaethylporphyrins where introduction of the first 3 meso residues can be accomplished in quantitative yield and only introduction of the D residue gives lower yields.^[34]

Synthesis of ABCD-porphyrins through Pd-catalyzed reactions: Having the ABC-type porphyrins in hand, another possibility for the construction of ABCD-type porphyrins was obvious. This involved bromination of the ABC porphyrins, followed by Pd-catalyzed C–C-couplings. The bromination of ABC porphyrins was achieved easily in good yields. For example, compounds **123** and **124** were obtained in 71 and 77 % yield, respectively (Scheme 14). Upon reaction of **82** with NBS a new reaction was observed. The product **125** showed that both meso bromination and bromination of the 2,4,6-trimethoxyphenyl residue had occurred in 54 % yield. Similar reactions have been observed in the literature for other systems.^[35]



Scheme 14. Transformation of ABC-porphyrins to ABCD-porphyrins via Suzuki reactions.

The bromo-ABC porphyrins were then subjected to Suzuki coupling conditions similar to those used for the preparation of ABC porphyrins from AB-type systems (see above). As shown in Scheme 14 the reactions work well, although the yields vary significantly and depend on the porphyrin and boron coupling partner used. In some cases (e.g. **127**) debromination of the starting

material (to **85**) was observed. The dibromo derivative **125** was also subjected to the coupling conditions to investigate the reactivity of the two different bromo residues. Reaction with 3-nitrophenyl boronic acid gave a mixture of the doubly substituted product **134** and compound **133** where only the porphyrin meso bromo residue had reacted.

Structural studies

Despite many attempts only two crystals of sufficient quality for single crystal X-ray determination could only be grown for two compounds. These were the closely related 5,15-dibromo compounds **77** (Fig. 1) and **81** (Fig. 2) which differ only in the steric demand of one meso aryl residue (2,4,6-trimethoxyphenyl in **77** versus 3,4,5-trimethoxyphenyl in **81**). As expected for an unsymmetrical compound^[36] **77** forms weakly π -stacked dimers in the crystal with a hexyl chain on the same face as a neighbouring aryl group, i.e. anti-parallel layers. In compound **81** the intermolecular separation is much larger and parallel layers are formed. The closest contact observed is a Br-CH₂ interaction (Br2...H10G, 2.883 Å). As shown in Table 1 both structures exhibit only very minor deviations from planarity. The largest deviations are observed for the b-positions (0.03–0.27 Å in **77** and 0.15–0.32 Å in **81**). These deviations are in line with crystal packing and minor steric effects to be expected for this class of compounds.^[38]




Fig. 1. View of the molecular structure of **77** in the crystal. Hydrogen atoms have been omitted for clarity; thermal ellipsoids are for 50 % occupancy.

In conformational terms the main difference between both compounds is the tilt angle between the trimethoxyphenyl residue and the least-squares-plane of the macrocycle. In compound **81** the tilt angle is 94.4°, i.e. the residue is orthogonal to the molecular plane. In the compound **77** the tilt angle is 70.6° which is more in line with those of other meso arylporphyrins.^[36a]




Fig. 2. View of the molecular structure of **81** in the crystal. Hydrogen atoms have been omitted for clarity; thermal ellipsoids are for 50 % occupancy.

Table 1. Selected structural and conformation parameters for **77** and **81** [Å].

Compound	77	81
Σ [a]	2.064	2.056
Ξ [b]	0.013	0.039
$\Delta 24$ [c]	0.06	0.15
ΔC_m [d]	0.06	0.09
δC_5 [e]	0.04	0.11
δC_{10} [e]	0.11	0.08
δC_{15} [e]	0.07	0.09
δC_{20} [e]	0.02	0.07

[a] Core size, average vector length from the geometric center of the four nitrogen atoms to the nitrogen atoms. [b] Core elongation parameter defined as the difference between the vector lengths ($|N21-N22|+|N23-N24|$)/2- ($|N22-N23|+|N21-N24|$)/2. [c][37] Average deviation of the 24 macrocycle atoms from their least-squares plane. [d] Average deviation of the C_m carbon atoms from the 4N-plane. [e] Average deviation of the C_m carbon atom from the 4N-plane.

Conclusions

A full analysis of the synthesis of all members of the ABCD-type porphyrin series using porphyrin reactions with organolithium reagents and Pd-catalyzed coupling reactions has been given. As shown the synthesis of the 5,15-A₂ starting materials is still best accomplished via condensation methods. Reaction of A-porphyrins with RLi works but is somewhat problematic due to the formation of regioisomers and multiply alkylation products. Here additional synthetic developments are required.^[24] Nevertheless, the AB-porphyrins are suitable starting materials for ABC-type systems through nucleophilic substitution and the reactions give good yield. Additionally, Pd-catalyzed reactions of appropriately brominated systems can be used but they require the separation of the mono- and dibromo starting materials. The ABC porphyrins can be transformed with RLi reagents to the respective ABCD-type systems but the yields are somewhat unsatisfactory and for sterically hindered systems side reactions may be observed. Similarly AB-systems can be converted via RLi/RI to ABCD-porphyrins, again with low yields. Here use of Pd-catalyzed reactions is preferred although the RLi method gives easier access to alkyl substituted systems. Monobromination of ABC-porphyrins proceeds well and the subsequent coupling reactions give acceptable yields. In order to optimize the overall yields a combination of both methods is necessary. Overall, a use of either method for the individual steps in the sequence A $\rightarrow$ AB $\rightarrow$ ABC $\rightarrow$ ABCD porphyrins can give almost any desired ABCD porphyrins.

Experimental Section

General methods: All chemicals used were of analytical grade and purified before use. Dichloromethane was dried over phosphorous pentoxide followed by distillation; THF was dried over sodium followed by distillation. Silica gel 60 (Merck) was used for column chromatography unless otherwise noted. Analytical thin-layer chromatography (TLC) was carried out using silica gel 60 plates (fluorescence indicator F₂₅₄; Merck). Melting points are uncorrected and were measured with a Reichert Thermovar instrument. NMR spectra were recorded using a Bruker AM 270 instrument (270 MHz), a Bruker DPX 400 (400.13 MHz for ¹H NMR and 100.61 MHz for ¹³C NMR) or a Bruker AV 600 (600.13 MHz for ¹H NMR and 150.90 MHz for ¹³C NMR) instrument. Chemical shifts are given in ppm and referenced to the TMS signal as internal standard. The assignment of the signals was confirmed by 2D spectra (COSY, HMBC, HMQC) except for those porphyrins with low solubility. Electronic absorption spectra were recorded with a Specord S10 instrument (Zeiss) using CH₂Cl₂ as solvent. Mass spectra were recorded using a Varian MAT 711 or MAT 112 S mass spectrometer using the EI technique with a direct insertion probe and an excitation energy of 80 eV. FAB spectra were recorded with CH-5 DF instrument from Varian. HRMS data were

determined using a Micromass TOF instrument fitted with an EI probe. Elemental analyses were performed with a Perkin-Elmer 240 analyzer. Data for single crystal X-ray determinations were collected using Rigaku Saturn-724 system complete with CCD detector utilizing Mo-K α radiation.

Starting materials: 5-(*tert*-butyl)porphyrin **32**, **33**, **34**, **39**,^[18] 5,15-bis(3-methoxyphenyl)porphyrin **58**,^[11] **48**,^[11] **87**,^[11] **88**,^[11] 5-(4-ethynylphenyl)-10,20-diphenylporphyrin **96**,^[15] 5,15-dihexylporphyrin **18**,^[11] 5-hexyl-15-(4-methoxyphenyl)porphyrin **51**,^[11] and 5-hexyl-15-(3,5-dimethoxyphenyl)porphyrin **112**,^[11] were prepared following published procedures.

5-(4-Methoxycarbonylphenyl)porphyrin (9): Dry dichloromethane (2 l) was placed in a three-necked-flask equipped with magnetic stirrer, gas inlet (argon) and a reflux condenser. Dipyrromethane (600 mg, 4 mmol), 2-hydroxymethyl pyrrole (680 μ L, 7.8 mmol), and methyl 4-formylbenzoate (630 mg, 3.8 mmol) were added. The flask was shielded from ambient light and then 1.40 ml (1.8 mmol) of trifluoroacetic acid were added and the reaction mixture was stirred for 18 h at 20 $^{\circ}$ C. After this time, 2.72 g (12 mmol) of DDQ suspended in 100 mL of dry dichloromethane were added and the mixture was stirred for 1 h. Next, 3 ml of triethylamine were added and the reaction mixture was stirred for 15 min. The reaction mixture was filtered through followed by chromatography on silica using dichloromethane as eluent. The first fraction was porphyrin (~10 mg), followed by the title compound and traces of 5,15-bis(4-methoxycarbonylphenyl)porphyrin. Recrystallization from CH₂Cl₂/MeOH gave 128 mg of red crystals (0.29 mmol; 8 %). M.p. >300 $^{\circ}$ C. R_f = 0.77 (CH₂Cl₂). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -3.60 (s, br, 2H, NH), 4.15 (s, 3H, COOCH₃), 8.33 (m, 2H, phenyl-H), 8.46 (m, 2H, phenyl-H), 9.06 (d, J = 4 Hz, 2H, β -pyrrole-H), 9.41 (d, J = 4 Hz, 2H, β -pyrrole-H), 9.50 (m, 4H, β -pyrrole-H), 10.25 (s, 1H, C15-H), 10.33 (s, 2H, C10-H and C20-H). UV/Vis (CH₂Cl₂): λ_{max} (log ϵ) = 400 (4.85), 422 (3.77), 586 (3.54). MS (EI, 80 eV, 140 $^{\circ}$ C) m/z (%) = 444 (100) [M⁺], 429 (1) [M⁺ - CH₃], 413 (1) [M⁺ - CH₃O], 385 (9) [M⁺ - C₂H₃O₂], 222 (1) [M²⁺]. HRMS calcd. for C₂₈H₂₀N₄O₂ 444.15863; found 444.15649. C₂₈H₂₀N₄O₂ (444.49): calcd. C 75.66, H 4.54, N 12.44; found C 75.98, H 4.76, N 12.23.

3,5-Di-(*tert*-butyl)benzaldehyde (10): Prepared in 58 % yield following the procedure given by Newman and Lee.^[39] M.p. 84 $^{\circ}$ C; lit. 84-85 $^{\circ}$ C. ¹H NMR (250 MHz, CDCl₃, TMS): δ = 9.99 (1H, s, CHO), 7.71 (3H, m, Ar-H_o, Ar-H_p), 1.35 ppm (18H, s, C(CH₃)₃). ¹³C NMR (63 MHz, CDCl₃): δ = 193.16, 151.87, 136.24, 128.87, 124.13, 34.97, 31.31 ppm. MS (40 $^{\circ}$ C, 80 eV): m/z = 219 (3 %, [M + 1]⁺), 218 (16 %, [M]⁺), 203 (100 %, [M - CH₃]⁺), 175 (3 %, [M - C₃H₇]⁺), 57 (3 %, [M - C₁₁H₁₃O]⁺).

5-(3,5-Di(*tert*-butyl)dipyrromethane (12): Prepared in 48 % yield following the procedure given by Beavington and Burn.^[40] ¹³C NMR (63 MHz, CDCl₃): δ = 150.80, 140.84, 132.94, 122.67, 120.81, 116.96, 108.25, 107.10, 44.45, 34.79, 31.44 ppm.

5-(3,5-Di(*tert*-butyl)porphyrin (13): Dipyrromethane (0.34 g, 2.3 mmol) and the 5-substituted dipyrromethane **12** (2.3 mmol) were dissolved in 1.2 L dichloromethane and degassed under Ar for 10 min. Next, trimethyl formate (38 mL, 0.35 mmol) was added and the solution treated dropwise with a solution of trichloroacetic acid (17.65 g, 108 mmol) in 460 mL dichloromethane. The mixture was stirred for 4 h at rt and then mixed with pyridine (31.2 mL, 0.39 mol) and stirring was continued for another 17 h. The solution was purged with oxygen for 10 min for oxidation and filtered through silica gel. After removal of the solvent *in vacuo* column chromatography (*n*-hexane:ethyl acetate = 5:1, v/v) gave a purple solid as the main fraction after removal of the solvent (23 mg, 0.05 mmol, 4 %). M.p. 273 $^{\circ}$ C. ¹H NMR (400 MHz, CDCl₃, TMS): δ = 10.32 (2H, s, meso-H_{10,20}), 10.24 (1H, s, meso-H₁₅), 9.49 (2H, AB, ³ $J_{12,13}$ = 4.6 Hz, ³ $J_{17,18}$ = 4.6 Hz, β -pyrrole-H_{12,18}), 9.46 (2H, AB, ³ $J_{12,13}$ = 4.6 Hz, ³ $J_{17,18}$ = 4.6 Hz, β -pyrrole-H_{13,17}), 9.41 (2H, AB, ³ $J_{2,3}$ = 4.6 Hz, ³ $J_{7,8}$ = 4.6 Hz, β -pyrrole-H_{2,8}), 9.14 (2H, AB, ³ $J_{2,3}$ = 4.6 Hz, ³ $J_{7,8}$ = 4.6 Hz, β -pyrrole-H_{3,7}), 8.12 (2H, m, Ar-H_o), 7.83 (1H, m, Ar-H_p), 1.55 (18H, s, C(CH₃)₃), -3.58 ppm (bs,). ¹³C NMR (126 MHz, CDCl₃): δ = 149.00, 140.70, 131.71, 131.17, 130.99, 130.10, 121.15, 104.56 (C10, C20), 103.30 (C15), 35.08 (C31, C35), 31.76 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ϵ) = 401 (5.33), 497 (4.19), 528 (3.44), 569 (3.70), 623 (2.78). MS (160-170 $^{\circ}$ C, 80 eV): m/z = 499 (36 %, [M + 1]⁺), 498 (100 %, [M]⁺). HRMS: calcd. for C₃₄H₃₄N₄ 498.27835; found 498.27444.

5,15-Bis(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)porphyrin (14): To a 2 L three-necked flask containing dichloromethane (1 l), dipyrromethane (1.38 g, 9.25 mmol) and 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzaldehyde (2.25 g, 9.71 mmol) were added under an Ar atmosphere. After 30 min TFA (0.1 g, 0.925 mmol) was added and the mixture was stirred for 12 h. Next two equiv.

DDQ (4.2 g, 18.5 mmol) were added and stirring was continued for 60 min at rt followed by addition of 1 mL triethylamine. Without evaporation of solvent, the crude reaction mixture was passed through a filter containing silica gel, using dichloromethane as eluent. The solvent was evaporated and the resulting residue was recrystallized from dichloromethane/ methanol to give purple crystals (1.92 g, 58.2%). M.p. > 300 $^{\circ}$ C. ¹H NMR (400 MHz, CDCl₃, TMS): δ = 10.35 (s, 2H, meso-H), 9.42 (d, 4H, J = 4.64 Hz, β -pyrrole-H), 9.1 (d, 4H, J = 4.64 Hz, β -pyrrole-H), 8.3 (dd, 8H, J = 9.065 Hz, Ph-H), 1.55 (s, 24H, CH₃), -3.1 ppm (br s, 2H, NH). ¹³C NMR (400 MHz, CDCl₃): δ = 133.97, 132.86, 131.24, 130.57, 104.90, 83.70, 28.91, 24.62, 24.42 ppm. UV/vis (THF): λ_{max} (log ϵ) = 406 (4.42), 502 (4.21), 535(3.95), 576(3.84), 631 nm (364). HRMS calcd. for C₄₄H₄₄B₂N₄O₄ 714.4666; found 715.3600.

[5,15-Bis(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)porphyrinato]nickel(II) (15): Free base porphyrin **14** (626.5 mg, 0.877 mmol) and nickel acetylacetonate (900 mg, 3.51 mmol) were dissolved in toluene (200 mL) and stirred for 4 days at 110 $^{\circ}$ C. The crude reaction mixture was passed through a filter containing silica gel, using CH₂Cl₂ as eluent. The solvent was evaporated and the resulting residue was recrystallized from dichloromethane/methanol to yield purple crystals (0.75 mmol, 86 %). M.p. >300 $^{\circ}$ C. ¹H NMR (400 MHz, CDCl₃, TMS): δ = 9.97 (s, 2H, meso-H), 9.21 (m, 2H, β -pyrrole-H), 8.95 (m, 2H, β -pyrrole-H), 8.2 (d, 2H, J = 7 Hz, β -pyrrole-H), 8.11 (d, 2H, J = 7 Hz, β -pyrrole-H), 7.97 (d, 2H, J = 7.6 Hz, Ph-H), 7.89 (d, 2H, J = 7.6 Hz, Ph-H), 1.53 (s, 24H, CH₃). UV/vis(THF): λ_{max} (log ϵ) = 400 (5.2), 515 (4.17), 546 nm (3.94). HRMS calcd. for C₄₄H₄₂B₂N₄O₄Ni 770.2746; found 770.2750.

5,10-Bis(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)porphyrin (16): Tripyrrane (2 g, 8.8 mmol), pyrrole (0.6 g, 8.88 mmol) and 4-(4,4, 5, 5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzaldehyde (4.22 g, 18.2 mmol) were added under an Ar atmosphere at room temperature to a 2L three-necked flask containing 1 L of dichloromethane. After 30 min, TFA (0.127 g, 0.888 mmol) was added, and mixture was stirred for 12 h. Then four equiv. of DDQ (8 g, 35.2 mmol) were added and stirring was continued for 60 min at rt followed by addition of 1 mL NEt₃. The crude reaction mixture was passed through a filter containing silica gel, eluting with CH₂Cl₂. The solvent was evaporated to near dryness and then absorbed onto 5g of silica gel. The absorbed sample was added to the top of the column containing silica gel and with *n*-hexane/acetone (1:1, v/v). The product was isolated as purple crystals in 0.4 % yield (27 mg). M.p. >300 $^{\circ}$ C. ¹H NMR (400 MHz, CDCl₃, TMS): δ = 10.27 (s, 2H, meso-H), 9.49 (s, 2H, β -pyrrole-H), 9.37 (d, 2H, J = 4.08 Hz, β -pyrrole-H), 9.04 (d, 2H, J = 4.68, β -pyrrole-H), 8.93 (s, 2H, β -pyrrole-H), 8.26 (dd, 8H, J = 7.6 Hz, Ph-H), 1.58 (s, 24H, CH₃), -3.34 ppm (br s, 2H, NH). ¹³C NMR (400 MHz, CDCl₃): δ = 24.61, 29.27, 83.7, 119.5, 103.84, 132.57, 133.73, 144.58 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ϵ) = 407 (4.55), 503 (3.28), 576 (2.79), 630 nm (2.30). HRMS calcd. for C₄₄H₄₄B₂N₄O₄ 714.4666; found 715.3685.

General Procedure for the synthesis of AB-type porphyrins (mixed condensation): A three-necked-flask equipped with magnetic stirrer, gas inlet (argon) and a reflux condenser was charged with 2L dichloromethane. Dipyrromethane (1200 mg, 8.2 mmol), the appropriate aldehyde (4.14 mmol), and the other appropriate aldehyde (4.15 mmol) were added. Then 140 μ L (1.8 mmol) TFA were added and the reaction mixture was stirred for 16 h at 20 $^{\circ}$ C. Next DDQ (2.77 g, 12.2 mmol) in 100 mL of dry dichloromethane was added and the mixture was stirred for 1 h. Subsequently, 6 mL of NEt₃ were added and the reaction mixture was filtered through 600 mL of silica, eluting with dichloromethane. Column chromatography eluting with dichloromethane gave the fractions which were recrystallized from CH₂Cl₂/MeOH to yield purple crystals.

5-Hexyl-15-(4-methoxycarbonylphenyl)porphyrin (17): The procedure followed that given above using dipyrromethane (1200 mg, 8.2 mmol), methyl 4-formylbenzoate (680 mg, 4.14 mmol), and heptanal (0.580 mL, 4.15 mmol). Column chromatography eluting with dichloromethane gave three fractions which were recrystallized from CH₂Cl₂/MeOH to yield purple crystals. The first fraction was 5,15-dihexylporphyrin **18** (181 mg, 0.378 mmol, 18 %), the second fraction 5-hexyl-15-(4-methoxycarbonylphenyl)porphyrin **17** (525 mg, 0.99 mmol; 23 %), and the third fraction 5,15-bis(4-methoxycarbonylphenyl)porphyrin **19** (168 mg, 0.29 mmol, 14 %). **17:** M.p. 256 $^{\circ}$ C. R_f = 0.37 (CH₂Cl₂). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -3.05, -3.04 (each s, 1H, NH), 0.92 (t, J = 7 Hz, 3H, CH₂CH₃), 1.41 (m, 2H, CH₂CH₃), 1.53 (m, 2H, CH₂CH₂CH₃), 1.80 (m, 2H, CH₂CH₂CH₂CH₃), 2.59 (m, 2H, CH₂CH₂CH₂CH₂CH₃), 4.11 (s, 3H, COOCH₃), 5.01 (t, J = 8 Hz, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 8.33 (m, 2H, Ph-H), 8.43 (m, 2H, Ph-H), 8.93 (d, J = 5 Hz, 2H, β -pyrrole-H), 9.36 (d, J = 5 Hz, 2H, β -pyrrole-H), 9.43 (d, J = 5 Hz, 2H, β -pyrrole-H), 9.61 (d, J =

5 Hz, 2H, β -pyrrole-*H*), 10.22 (s, 2H, meso-*H*). UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 407 (5.18), 503 (4.07), 536 (3.63), 576, 640 nm (3.55). HRMS calcd. for $\text{C}_{34}\text{H}_{32}\text{N}_4\text{O}_2$ 528.2525; found 528.2749. $\text{C}_{34}\text{H}_{32}\text{N}_4\text{O}_2$ (528.65): calcd. C 77.25, H 6.10, N 10.60; found C 77.21, H 6.49, N 10.99.

5,15-Bis(4-methoxycarbonylphenyl)porphyrin (19): Obtained from the synthesis of **17**. M.p. >300 °C. R_f = 0.11 (CH_2Cl_2). ^1H NMR (400 MHz, CDCl_3 , TMS): δ = 3.13 (s, 2H, *NH*), 4.12 (s, 6H, COOCH_3), 8.37 (*m*, 4H, *Ph-H*), 8.45 (*m*, 4H, *Ph-H*), 9.13 (*d*, J = 5 Hz, 4H, β -pyrrole-*H*), 9.38 (*d*, J = 5 Hz, 4H, β -pyrrole-*H*), 10.18 ppm (s, 2H, meso-*H*). UV/Vis (CH_2Cl_2): λ_{max} = 405, 506, 534, 574, 639 nm. HRMS calcd. for $\text{C}_{36}\text{H}_{26}\text{N}_4\text{O}_4$ 578.1954; found 578.2314. $\text{C}_{36}\text{H}_{26}\text{N}_4\text{O}_4$ (578.63): calcd. C 74.73, H 4.53, N 9.68; found C 74.51, H 4.78, N 9.57.

General procedure for the synthesis of 5,15-AB-type porphyrins (organolithium reaction): E.g., 5-(*tert*-Butyl)porphyrin **32** (100 mg, 0.272 mmol) was dissolved in 80 mL dry THF under argon atmosphere and the reaction mixture was cooled down to -78 °C. *n*BuLi (0.435 mL, 1.088 mmol) was added dropwise over 15 minutes *via* syringe. The cold bath was removed and stirring continued 1.5 hours at room temperature. 0.5 mL H_2O was added and stirring continued 15 min. Then DDQ (243.3 mg, 1.088 mmol) was added in 10 mL THF and the reaction mixture was stirred for an additional hour. The reaction mixture was filtered through silica gel, followed by evaporation of the solvent.

5-(*n*-Butyl)-15-(*tert*-butyl)porphyrin (38): Prepared according to the general procedure using 5-(*tert*-Butyl)porphyrin **32** (100 mg, 0.272 mmol), *n*BuLi (0.435 mL, 1.088 mmol) and DDQ (243.3 mg, 1.088 mmol). The crude reaction mixture was purified by column chromatography eluting with *n*-hexane/ CH_2Cl_2 (2:1, v/v). The first fraction was 5-(*n*-butyl)-15-(*tert*-butyl)porphyrin **38** (20 mg, 0.047 mmol, 17 %) as purple crystals, the second fraction 5-(*n*-butyl)-10-(*tert*-butyl)porphyrin **39** (8 mg, 0.018 mmol, 7 %) as purple crystals, followed by starting material (10 mg, 0.027 mmol, 10 %). Compound **38**: M.p. > 300 °C. R_f = 0.6 (*n*-hexane/ CH_2Cl_2 = 1:1, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): δ = -1.97 (s, 2H, *NH*), 1.15 (t, 3H, J = 7.0 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.81 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.50 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.61 (s, 9H, $\text{C}(\text{CH}_3)_3$), 4.94 (t, 2H, J = 7.6 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 9.24 (d, 2H, J = 4.68 Hz, β -pyrrole-*H*), 9.34 (d, 2H, J = 4.1 Hz, β -pyrrole-*H*), 9.50 (d, 2H, J = 4.68 Hz, β -pyrrole-*H*), 9.86 (d, 2H, J = 4.68 Hz, β -pyrrole-*H*), 10.06 ppm (s, 2H, meso-*H*). ^{13}C -NMR (100 MHz, CDCl_3 , TMS): δ = 13.7, 23.1, 29.2, 33.4, 39.8, 40.0, 40.3, 103.9, 119.1, 127.1, 129.7, 130.7, 131.0, 141.4, 143.0, 146.6, 147.7 ppm. UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 405 (4.68), 509 (3.62), 542 (3.52), 542 (3.52), 636 nm (3.22). MS (EI, 250 °C, 80 eV): m/z = 450 (100 %, $[\text{M}]^{+}$), 435 (72 %, $[\text{M} - \text{CH}_3]^{+}$), 394 (10 %, $[\text{M} - \text{C}_4\text{H}_9 + \text{H}]^{+}$), 379 (7 %, $[\text{M} - \text{C}_5\text{H}_{11}]^{+}$), 323 (17 %, $[\text{M} - \text{C}_4\text{H}_9 + \text{H} - \text{C}_5\text{H}_{11}]^{+}$), 225 (3 %, $[\text{M}]^{2+}$). HRMS calcd. for $\text{C}_{28}\text{H}_{31}\text{N}_4$ 423.2549; found 423.2551.

5-(*n*-Butyl)-10-(*tert*-butyl)porphyrin (39): Derived from the synthesis of **38**. M.p. > 300 °C. R_f = 0.4 (*n*-hexane/ CH_2Cl_2 = 1:1, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): δ = -2.41 (s, 2H, *NH*), 1.15 (t, 3H, J = 7.01 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.81 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.52 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.46 (s, 9H, $\text{C}(\text{CH}_3)_3$), 4.93 (t, 2H, J = 7.6 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 9.11 (d, 1H, J = 4.67 Hz, β -pyrrole-*H*), 9.2 (d, 1H, J = 4.67 Hz, β -pyrrole-*H*), 9.25 (*m*, 3H, β -pyrrole-*H*), 9.5 (d, 1H, J = 4.68 Hz, β -pyrrole-*H*), 9.69 (d, 1H, J = 5.26 Hz, β -pyrrole-*H*), 9.79 (*m*, 2H, J = 4.68 Hz, β -pyrrole-*H*, meso-*H*), 10.06 ppm (s, 1H, meso-*H*). ^{13}C -NMR (100 MHz, CDCl_3 , TMS): δ = 13.6, 13.7, 19.2, 22.2, 22.5, 23.2, 26.6, 28.9, 29.2, 29.7, 30.5, 31.4, 32.3, 35.0, 36.6, 36.9, 40.2, 102.0, 103.3, 120.2, 125.5, 127.4, 128.3, 129.1, 129.2, 130.0, 130.4, 130.7, 131.0, 167.2 ppm. UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 412 (4.81), 513 (3.80), 551 (3.32), 591 (3.47), 645 nm (3.10). HRMS calcd. for $\text{C}_{28}\text{H}_{31}\text{N}_4$ 423.2549; found 423.2553.

5-*tert*-Butyl-15-[4-(*N,N*-dimethylamino)phenyl]porphyrin (41): A 250 mL Schlenk flask was charged with 4-bromo-*N,N*-dimethyl aniline (11 equiv.) dissolved in 20 mL diethyl ether. A 2.5 M solution of *n*BuLi in hexane (1 equiv.) was added dropwise over 45 min under Ar at -78 °C. The cold bath was removed and the solution stirred for 2 h at rt. A 100 mL Schlenk flask was charged with a solution of the porphyrin **32** (53 mg, 0.14 mmol, 1 equiv.) in 50 mL THF and cooled to -70 °C. This solution was then added to the solution of the *in situ* generated LiR reagent (cooled to -40 - -70 °C, 11 equiv.). The color changed from red to green-brown. The cold bath was removed and the solution stirred for 2 h at rt. Next 5 mL water in 5 mL THF was added and this was accompanied by a color change to green. The solution was stirred for 30 min followed by addition of 10 equiv. DDQ which resulted in a color change to red. The mixture was filtered through silica gel and then neutral alumina eluting with dichloromethane followed by removal of the solvent *in vacuo*. Column chromatography on silica gel (CH_2Cl_2 /*n*-hexane = 1:3, v/v and CH_2Cl_2 /*n*-hexane = 1:7, v/v) gave 5-*n*-butyl-15-*tert*-butylporphyrin **38** (2 mg (< 0.01

mmol, 5 %) as the first fraction. The title compound **41** followed as the second fraction to yield purple crystals after recrystallization from CH_2Cl_2 /MeOH (19 mg (0.04 mmol, 36 %). M.p. 270 °C (decomp.). R_f = 0.4 (CH_2Cl_2 : *n*-hexane = 9:1, v/v). ^1H NMR (300 MHz, CDCl_3 , TMS): δ = 10.07 (s, 2H, meso-*H*), 9.85 (*AB*, 2H, 3J = 4.8 Hz, β -pyrrole-*H*_{3,7}), 9.26 (*AB*, 2H, 3J = 4.7 Hz, β -pyrrole-*H*_{12,18}), 9.23 (*AB*, 2H, 3J = 4.8 Hz, β -pyrrole-*H*_{2,8}), 9.10 (*AB*, 2H, 3J = 4.7 Hz, β -pyrrole-*H*_{13,17}), 8.12 (*AB*, 2H, 3J = 8.2 Hz, Ar-*H*_o), 7.15 (*AB*, 2H, 3J = 8.2 Hz, Ar-*H*_m), 3.23 (s, 6H, $\text{N}(\text{CH}_3)_2$), 2.60 (s, 9H, $\text{C}(\text{CH}_3)_3$), -1.93 ppm (*br s*, 2H, *NH*). ^{13}C NMR (75 MHz, CDCl_3): δ = 150.04, 148.10, 147.61, 143.90, 142.10, 135.96, 131.09, 131.06, 130.62, 130.13, 128.78, 126.66, 120.14, 111.38, 104.69, 40.90, 40.73, 40.41 ppm. UV/vis (CH_2Cl_2): λ_{max} (log ϵ) = 413 (4.78), 513 (4.07), 554 (4.07), 581 (3.89), 641 nm (3.70). MS (250 °C, 80 eV): m/z = 485 (92 %, $[\text{M}]^{+}$), 470 (100 %, $[\text{M} - \text{CH}_3]^{+}$), 429 (25 %, $[\text{M} - \text{CH}_3 - \text{C}_6\text{H}_5\text{N}]^{+}$), 243 (23 %, $[\text{M}]^{2+}$), 295 (30 %, $[\text{M} - \text{CH}_3]^{2+}$). HRMS: calcd. for $\text{C}_{32}\text{H}_{31}\text{N}_5$ 485.2579, found 485.2562.

General procedure for the synthesis of 5,10-AB-type porphyrins: A 250 mL Schlenk flask was charged with the porphyrin (1 equiv.) in 50 mL THF under Ar and cooled to -70 °C. The LiR reagent (hexyllithium, 3.5 equiv. of a 2.5 M solution in hexane) was added dropwise over 15 min and the color changed from red to green-brown. The cold bath was removed and the solution stirred for 30 min at rt. Next water (1 mL in 5 mL THF) was added and the color changed to green. The solution was stirred for 15 min and then oxidized with DDQ followed by stirring for a further 30 min accompanied by a color change to red. The mixture was filtered through silica gel eluting with dichloromethane and the solvents removed *in vacuo*.

5-(1-Ethylpropyl)-10-hexylporphyrin (42): Prepared using the general procedure given above using porphyrin **33** (1 equiv.) and hexyllithium (3.5 equiv. of a 2.5 M solution in hexane). Column chromatography on silica gel eluting with CH_2Cl_2 /*n*-hexane (1:1, v/v) gave a first fraction containing a mixture of di- and higher alkylated species followed by the title compound as purple crystals after recrystallization from CH_2Cl_2 /MeOH/ H_2O (37.1 mg, 0.08 mmol, 59 %). M.p. 135 °C. R_f = 0.46 (*n*-hexane/ CH_2Cl_2 = 1:1, v/v). ^1H NMR (500 MHz, CDCl_3 , TMS): δ = 10.06 (s, 1H, meso-*H*₁₅), 10.01 (s, 1H, meso-*H*₂₀), 9.79 (*m*, 1H, β -pyrrole-*H*₇), 9.73 (*m*, 1H, β -pyrrole-*H*₃), 9.65 (*m*, 1H, β -pyrrole-*H*₈), 9.62 (*AB*, 1H, $^3J_{\text{H12,H13}}$ = 4.7 Hz, β -pyrrole-*H*₁₂), 9.33 (*m*, 4H, β -pyrrole-*H*_{2,13,17,18}), 5.14 (*m*, 1H, *m*, $\text{CH}(\text{CH}_2)_2$), 5.09 (*t*, 2H, 3J = 8.2 Hz, $\text{CH}_2\text{C}_5\text{H}_{11}$), 3.00, 2.84 (*m*, 4H, $\text{CH}(\text{CH}_2)_2$), 2.60 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{C}_4\text{H}_9$), 1.87 (*m*, 2H, $\text{C}_2\text{H}_4\text{CH}_2\text{C}_3\text{H}_7$), 1.55 (*m*, 2H, $\text{C}_3\text{H}_6\text{CH}_2\text{C}_2\text{H}_5$), 1.43 (*m*, 2H, $\text{C}_4\text{H}_8\text{CH}_2\text{CH}_3$), 0.97 (*t*, 3H, 3J = 7.4 Hz, $\text{C}_3\text{H}_{10}\text{CH}_3$), 0.96 (*t*, 6H, 3J = 7.3 Hz, CH_2CH_3), -3.22 ppm (*br s*, 2H, *NH*). ^{13}C NMR (126 MHz, CDCl_3): δ = ~130 (C2, C3, C7, C8, C12, C17, C18), 123.21 (C5), 120.14 (C10), 103.11 (C15), 102.84 (C20), 50.53, 39.06, 36.10, 34.83, 31.94, 30.40, 22.78, 14.13 ppm. UV/vis (CH_2Cl_2): λ_{max} (log ϵ) = 406 (5.58), 504 (4.14), 534 (3.37), 579 (3.60), 634 nm (3.08). MS (EI, 80 eV): m/z = 464 (100 %, $[\text{M}]^{+}$), 435 (93 %, $[\text{M} - \text{C}_2\text{H}_5]^{+}$), 393 (20 %, $[\text{M} - \text{C}_5\text{H}_{11}]^{+}$), 364 (11 %, $[\text{M} - \text{C}_5\text{H}_{11} - \text{C}_2\text{H}_5]^{+}$), 335 (10 %, $[\text{M} - \text{C}_5\text{H}_{11} - 2 \text{C}_2\text{H}_5]^{+}$), 232 (11 %, $[\text{M}]^{2+}$). HRMS calcd. for $\text{C}_{31}\text{H}_{37}\text{N}_4$ 465.3018; found 465.3035.

General procedure for the synthesis of 5,15-AB₂/A₂-type porphyrins: A 25 mL Schlenk flask was charged with dried porphyrin (0.05 mmol) dissolved in 10 mL THF. The solution was cooled to -80 °C and treated with 0.2 mmol of a 2.5 M solution of hexyllithium (80 μL). The color changed from red to green-brown and after 10 min changed to purple. The cold bath was removed and the solution stirred for 15 min at rt. Next water (1 mL) was added and stirring was continued for 15 min. This was accompanied by a color change to yellow-brown. Then DDQ (0.1 mmol, 23 mg) in 0.4 mL THF was added and the mixture stirred for 15 min until the color had changed to dark red. The mixture was filtered through silica gel and the solvent removed *in vacuo*.

5-(3,5-Di-*tert*-butylphenyl)-10,15-dihexylporphyrin (56): Prepared following the above general procedure using porphyrin **13** (23 mg, 0.05 mmol) and a 2.5 M solution of hexyllithium (80 μL , 0.2 mmol). Column chromatography eluting with *n*-hexane/ CH_2Cl_2 = 1:1, v/v) gave a purple solid (6 mg, 0.01 mmol, 18 %). M.p. 192 °C. ^1H NMR (500 MHz, CDCl_3 , TMS): δ = 10.00 (s, 1H, meso-*H*₂₀), 9.61, 9.57 (*AB*, 2H, 3J = 4.8 Hz, β -pyrrole-*H*_{12,13}), 9.54 (*AB*, 1H, 3J = 4.6 Hz, β -pyrrole-*H*₁₇), 9.50 (*AB*, 1H, 3J = 4.6 Hz, β -pyrrole-*H*₈), 9.31 (*AB*, 1H, 3J = 4.6 Hz, β -pyrrole-*H*₁₈), 9.21 (*AB*, 1H, 3J = 4.7 Hz, β -pyrrole-*H*₂), 8.95 (*AB*, 2H, 3J = 4.7 Hz, β -pyrrole-*H*_{3,7}), 8.05 (d, 2H, J = 1.9 Hz, Ar-*H*_o), 7.79 (t, 1H, J = 1.9 Hz, Ar-*H*_p), 5.05, 5.00 (t, 4H, J = 8.1 Hz, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.57 (*m*, 4H, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.84 (*m*, 4H, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.55 (*m*, 4H, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.53 (s, 9H, $\text{C}(\text{CH}_3)_3$), 1.41 (*m*, 4H, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 0.94 (t, 6H, J = 7.3 Hz, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), -2.91 ppm (s, 2H, *NH*). ^{13}C -NMR (63 MHz, CDCl_3 , TMS): δ = 148.85, 140.93, 132.52, 131.27, 130.80, 129.85, 128.47,

127.91, 127.91, 128.24, 120.96, 120.42, 119.29, 103.38, 39.06, 38.80, 36.17, 35.33, 35.06, 31.76, 30.34, 22.77, 14.14 ppm. UV/Vis (CH₂Cl₂): $\lambda_{\max}$ (log ϵ) = 413 (5.50), 511 (4.19), 545 (3.78), 588 (3.65), 644 nm (3.54). MS (EI, 80 eV): m/z = 668 (46 %), [M + 1]⁺, 667 (100 %), [M]⁺, 610 (9 %), [M - C₄H₉]⁺, 596 (14 %), [M - C₅H₁₁]⁺, 539 (12 %), [M - C₉H₂₀]⁺, 525 (30 %), [M - C₁₀H₂₂]⁺. HRMS calcd. for C₄₁H₄₈N₄ 596.38790; found 596.38421.

5,15-Bis(3-methoxyphenyl)-10-phenylporphyrin (59): Phenyllithium (2 ml of a 1.8 M solution in hexane, 0.06 mmol) was slowly added (ca. 1 h) under an argon atmosphere to a 100 ml Schlenk flask charged with a solution of 5,15-bis(*m*-methoxyphenyl)porphyrin **58** (200 mg, 0.38 mmol) in 40 ml of dry THF at -80 °C. The reaction mixture was stirred for 5 h (TLC control). Subsequently, a mixture of 2 ml of water in 3 ml of THF was added for hydrolysis. After stirring of the mixture for 20 min, a solution of 10 equivalents of DDQ in THF (0.03 M) was added and the reaction mixture was stirred for another 60 min at room temperature. Subsequently, the mixture was filtered through silica gel and the organic solvent was removed under vacuum. Final purification was achieved by column chromatography and elution with ethyl acetate/*n*-hexane (1:4, v/v) yielding the title compound (163 mg, 0.27 mmol, 71 %) as purple crystals. M.p. >300 °C. R_f = 0.52 (ethyl acetate/*n*-hexane, 1:4, v/v). ¹H NMR (300 MHz, CDCl₃, TMS): δ = -2.91 (*s*, 2H, *NH*), 4.02 (*s*, 6H, OCH₃), 7.41 (*m*, 2H, Ph-*H*), 7.71-7.92 (*m*, 9H, Ph-*H*), 8.29 (*m*, 2H, Ph-*H*), 8.96 (*d*, 2H, *J* = 5.0 Hz, β -pyrrole-*H*), 9.02 (*d*, 2H, *J* = 5.0 Hz, β -pyrrole-*H*), 9.12 (*d*, 2H, *J* = 5.0 Hz, β -pyrrole-*H*), 9.33 (*d*, 2H, *J* = 5.0 Hz, β -pyrrole-*H*), 10.19 ppm (*s*, 1H, meso-*H*). ¹³C NMR (300 MHz, CDCl₃): δ = 55.06, 104.34, 113.09, 118.94, 120.21, 126.12, 127.19, 130.96, 134.05, 142.11, 142.64, 157.66 ppm. UV/Vis (CH₂Cl₂): $\lambda_{\max}$ (log ϵ) = 413 nm (5.17), 441 (3.94), 508 (4.23), 541 (3.73), 582 (3.77), 637 nm (3.52). MS (EI, 80 eV): m/z (%): 598 (100) [M]⁺, 567 (10) [M⁺ - 2 × CH₃ - H], 522 (8) [M⁺ - C₆H₄], 491 (8) [M⁺ - 2 × CH₃ - C₆H₅], 299 (86) [M]⁺. HRMS calcd. for C₄₀H₃₀N₄O₄ 598.2368; found 598.2356.

5-(4-Hydroxyphenyl)-10,20-bis(3-methoxyphenyl)porphyrin (60): *n*-Butyllithium (4 ml of a 2.5 M solution in hexane, 10 mmol) was added under an argon atmosphere to a 100 ml Schlenk flask charged with a solution of *p*-bromophenol (0.87 g, 5 mmol) in 10 ml of dry diethyl ether at 0 °C. After addition of *n*-butyllithium the cold bath was removed and stirring was continued for 18 h at room temperature. To the vigorously stirred mixture was added rapidly a solution of 5,15-bis(3-methoxyphenyl)porphyrin **58** (200 mg, 0.38 mmol) in 40 ml of dry THF under an argon atmosphere. Further workup followed the procedure for **59**. Final purification by column chromatography on silica gel and elution with ethyl acetate/*n*-hexane (1:1, v/v) gave the title compound (182 mg, 0.29 mmol, 77 %) as purple crystals. M.p. >300 °C. R_f = 0.54 (ethyl acetate/*n*-hexane, 2 : 1, v/v). ¹H NMR (300 MHz, CDCl₃, TMS): δ = 4.01 (*s*, 6H, OCH₃), 6.89 (*d*, 2H, *J* = 7.5 Hz, Ph-*H*), 7.37 (*d*, 2H, *J* = 7.5 Hz, Ph-*H*), 7.66 (*t*, 2H, Ph-*H*), 7.88 (*m*, 4H, Ph-*H*), 7.90 (*s*, 1H, OH), 7.97 (*d*, 2H, *J* = 7.5 Hz, Ph-*H*), 8.91 (*d*, 2H, *J* = 5.0 Hz, β -pyrrole-*H*), 9.00 (*d*, 2H, *J* = 5.0 Hz, β -pyrrole-*H*), 9.12 (*d*, 2H, *J* = 5.0 Hz, β -pyrrole-*H*), 9.32 (*d*, 2H, *J* = 5.0 Hz, β -pyrrole-*H*), 10.17 ppm (*s*, 1H, meso-*H*). ¹³C NMR (300 MHz, CDCl₃): δ = 55.06, 104.22, 112.95, 118.91, 120.25, 127.38, 131.01, 134.33, 135.06, 142.67, 154.87, 157.62 ppm. UV/Vis (CH₂Cl₂): $\lambda_{\max}$ (log ϵ) = 414 (5.17), 445 (4.06), 508 (4.17), 542 (3.61), 582 (3.64), 655 nm (3.39). MS (EI, 80 eV): m/z (%): 614 (38) [M]⁺, 307 (40) [M]⁺. HRMS calcd. for C₄₀H₃₀N₄O₅ 614.2317; found 614.2288.

General procedure for the synthesis of ABC-type porphyrins: Porphyrin (0.1 mmol) was dissolved in THF and cooled to -78 °C. Hexyllithium was added dropwise over the course of 15 min. The cold bath was removed and stirring was continued for another 15 min. This was followed by addition of 0.5 mL water in 5 mL THF, stirring for 15 min and addition of 6 mL of a solution of DDQ in THF. The mixture was stirred for 20 min and then filtered through neutral alumina washing with dichloromethane.

5-Hexyl-10-(4-methylphenyl)-20-(2,4,6-trimethoxyphenyl)porphyrin (61): Prepared following the above standard procedure using porphyrin **21** (55 mg, 0.1 mmol). Column chromatography eluting with neat dichloromethane gave the title compound as the single product as a purple solid (59 mg, 0.09 mmol, 90 %). M.p. >300 °C. ¹H NMR (500 MHz, CDCl₃, TMS): δ = 10.09 (*s*, 1H, meso-*H*), 9.60 (*m*, 2H, β -pyrrole-*H*), 9.30 (*m*, 2H, β -pyrrole-*H*), 8.90 (*m*, 4H, β -pyrrole-*H*), 8.21 (*d*, 2H, ³*J* = 7.35 Hz, tolyl-*H*_m), 7.56 (*d*, 2H, ³*J* = 7.35 Hz, tolyl-*H*_o), 6.68 (*s*, 2H, Ar-*H*_o), 5.04 (*t*, 2H, ³*J* = 8.09 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 4.11 (*s*, 3H, *p*-OCH₃), 3.50 (*s*, 6H, *m*-OCH₃), 2.70 (*s*, 3H, C₆H₄-CH₃), 2.56 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.83 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.54 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.43 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.00 (*t*, 3H, ³*J* = 7.30 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), -3.05 ppm (*bs*, 2H,

NH). ¹³C NMR (126 MHz, CDCl₃): δ = 161.99, 161.12, 148.41-145.14, 142.371, 139.80, 137.14, 134.50, 131.02-130.50, 127.25, 119.12-118.56, 104.00, 102.87, 91.00, 56.06, 55.64, 38.81, 35.51, 31.93, 30.34, 22.75, 21.51, 14.14 ppm. UV/vis (CH₂Cl₂): $\lambda_{\max}$ (log ϵ) = 413 (5.45), 511 (4.53), 543 (4.09), 585 (4.03), 641 nm (3.81). MS (270 °C, 8 kV): m/z = 650.0 (100.00 %, [M]⁺), 579.0 (71.47 %, [M - C₅H₁₁]⁺), 325.0 (10.91 %, [M]⁺). HRMS calcd. for C₄₂H₄₂N₄O₃ 650.32569; found 650.32575.

5-(2-Methoxyphenyl)-10-(4-methylphenyl)-20-(2,4,6-trimethoxyphenyl)porphyrin (63): The free base porphyrin **21** (113 mg, 0.2 mmol) was dissolved in 40 mL abs. THF and cooled to -40 °C. The LiR reagent was added under vigorous stirring, the cold bath removed and the mixture stirred for 1h. The LiR reagent was prepared by dissolving 1 g bromoanisole (5.4 mmol) in 15 mL abs. diethylether in a Schlenk tube. Next 2.2 mL BuLi (2.5 M in *n*-hexane, 5.5 mmol) were added dropwise over 30 min at -78 °C. The cold bath was removed and the mixture stirred for 1 h at rt. After mixing the two components the red mixture slowly turned brown. Next, 5 mL water in 5 mL THF were added and stirring continued for 30 min. Oxidation was achieved by stirring under air for 16h. The mixture was filtered through neutral alumina and washed with dichloromethane. Column chromatography eluting with CH₂Cl₂/*n*-hexane (1:2, v/v) yielded the target compound as purple crystals after recrystallization from CH₂Cl₂/MeOH (90 mg, 0.13 mmol, 67 %). M.p. > 320 °C. ¹H NMR (500 MHz, CDCl₃, TMS): δ = 10.23 (*s*, 1H, meso-*H*), 9.33 (*m*, 4H, β -pyrrole-*H*), 9.08 (*d*, 2H, ³*J* = 4.54 Hz, β -pyrrole-*H*), 9.02 (*d*, 2H, ³*J* = 4.54 Hz, β -pyrrole-*H*), 8.16 (*d*, 2H, ³*J* = 7.81 Hz, tolyl-*H*_o), 7.61 (*d*, 2H, ³*J* = 7.45 Hz, tolyl-*H*_m), 6.63 (*s*, 2H, OMe-Ar-*H*_m), 4.12 (*s*, 3H, *p*-OCH₃), 3.53 (*s*, 6H, *o*-OCH₃), 2.71 (*s*, 3H, C₆H₄-CH₃), -3.05 ppm (*bs*, 2H, *NH*). ¹³C NMR (126 MHz, CDCl₃): δ = 162.03, 161.21, 159.29, 148.84-146.42, 139.13, 137.20, 135.42, 134.60, 131.56-130.08, 127.46, 119.62, 111.95, 104.02, 91.09, 56.00, 55.59, 55.47, 21.47 ppm. UV/Vis (CH₂Cl₂): $\lambda_{\max}$ (log ϵ) = 407 (5.23), 504 (4.03), 539 (3.60), 575 (3.55), 630 nm (3.01). MS (FAB⁺, 3 eV): m/z = calcd. for C₄₃H₃₆N₄O₄ 672.27366; found 673.0 (100.00 %, [M+H]⁺). C₄₃H₃₆N₄O₄ (672.78): calcd. C 76.77, H 5.39, N 8.33; found C 76.55, H 5.70, N 8.57.

General procedure for the metallation with manganese chloride: Free base porphyrin (100mg, 0.14 mmol) was dissolved together with 150 mg MnCl₂·4 H₂O in 25 mL acetic acid (96 %) and 7 mL acetic anhydride. The solution was heated for 4 h at 110 °C. After cooling to rt dichloromethane was added and the organic phase washed several times with water. The organic solvent was evaporated and the residue dried under high vacuum.

Chloro[5-hexyl-10-(4-methylphenyl)-20-(2,4,6-trimethoxyphenyl)porphyrinato]manganese(III) (65): Prepared following the above general procedure using porphyrin **61** (100 mg, 0.14 mmol) and MnCl₂·4 H₂O (150 mg). Column chromatography eluting with dichloromethane gave a minor compound followed by the metal complex. Evaporation of the solvent gave a green-red residue (99 mg, 0.12 mmol, 88 %). M.p. >300 °C. UV/Vis (CH₂Cl₂): $\lambda_{\max}$ (log ϵ) = 370 (4.38), 405 (4.40), 479 (4.60), 581 (3.71), 614 (3.66), 643 nm (3.55). MS (FAB⁺, 3 kV): m/z = calcd. for C₄₂H₄₀ClMnN₄O₃ 738.21694; found 738.0 (9.0 %, [M]⁺), 703.0 (100.0 %, [M - Cl]⁺).

5,10-Bis(2-hydroxyphenyl)-15-(4-methylphenyl)porphyrin (71): A solution of **70** (200 mg of the atropisomeric mixture, 0.33 mmol) in 500 mL dichloromethane was treated dropwise with 1.3 mL BBr₃ solution (13 mmol) over the course of 1 h. The mixture was stirred for 16 h at rt and then poured onto ice and treated with NaHCO₃ until neutral. The solution was diluted with dichloromethane and the organic phase extracted three times with water. After drying of Na₂SO₄ the solvent was removed *in vacuo*. Column chromatography on silica gel eluting with neat dichloromethane gave one product fraction which yielded a red-brown solid (120 mg, 0.21 mmol, 62 %). M.p. > 300 °C. ¹H NMR (500 MHz, CDCl₃, TMS): δ = 10.14 (*s*, 1H, meso-*H*), 9.29 (*m*, 2H, β -pyrrole-*H*_{2,18}), 8.99 (*m*, 1H, β -pyrrole-*H*₇), 8.90 (*m*, 5H, β -pyrrole-*H*_{3,7,8,12,13}), 8.27 (*m*, 2H, tolyl-*H*_o), 8.10 (*m*, 2H, Ar-*H*_o), 7.90 (*m*, 2H, Ar-*H*_p), 7.06 (*m*, 2H, tolyl-*H*_m), 7.21 (*m*, 4H, Ar-*H*_m), -2.70 ppm (*bs*, 2H, *NH*). UV/Vis (CH₂Cl₂): $\lambda_{\max}$ (log ϵ) = 413 (5.21), 509 (3.81), 544 (3.12), 584 (3.05), 639 nm (2.42). MS (290 °C, 80 eV): m/z = 584.0 (100.00 %, [M]⁺). HRMS: calcd. for C₃₀H₂₈N₄O₂ 584.22123; found 584.22125.

5,10-[2,2'-(Dodecamethyleneoxy)diphenyl]-15-(4-methylphenyl)porphyrin (72): A solution of 100 mg of the dihydroxyporphyrin **71** (0.2 mmol) and 400 mg K₂CO₃ in 25 mL DMF was heated to 100 °C. Over the course of 4 h 81.5 mg 1,12-dibromodecane (0.25 mmol dissolved in 10 mL DMF) were added dropwise and heating continued for another 3h. The solution was cooled and treated with saturated aq. ammonium chloride. After addition of 50 mL dichloromethane the organic phase was washed three times with water and dried over Na₂SO₄. The solvent was removed in high vacuum and column

chromatography with CH₂Cl₂/*n*-hexane (1:2, v/v) gave the title compound as a purple solid (90 mg, 0.12 mmol, 60 %). M.p. 300 °C. ¹H NMR (500 MHz, CDCl₃, TMS): δ = 10.14 (s, 1H, meso-*H*), 9.26 (*m*, 2H, β-pyrrole-*H*_{2,18}), 8.97 (*AB*, 1H, ³*J* = 4.41 Hz, β-pyrrole-*H*₁₇), 8.89 (*AB*, 1H, ³*J* = 5.50 Hz, β-pyrrole-*H*₁₃), 8.83 (*AB*, 2H, ³*J* = 4.41 Hz, β-pyrrole-*H*_{3,12}), 8.75 (*AB*, 2H, ³*J* = 4.41 Hz, β-pyrrole-*H*_{7,8}), 8.09 (*m*, 4H, Ar-*H*_o), 7.76 (*m*, 2H, Ar-*H*_p), 7.56 (*d*, 2H, ³*J* = 8.09 Hz, tolyl-*H*_m), 7.34 (*m*, 4H, Ar-*H*_m), 3.86 (*m*, 4H, alkyl chain), 0.86 (*m*, 6H, alkyl chain), -0.81 (*m*, 8H, alkyl chain), -1.11 (*m*, 6H, alyl chain), -2.92 ppm (*bs*, 2H, NH). UV/Vis (CH₂Cl₂): λ_{max} (log ε) = 415 (5.39), 511 (4.22), 545 (3.82), 582 (3.71), 635 nm (3.36). MS (300 °C, 80 eV): *m/z* = 750.0 (100.00 %, [M]⁺), 375.0 (9.62 %, [M]²⁺). HRMS calcd. for C₅₁H₅₀N₄O₂ 750.39338; found 750.39341. C₅₁H₅₀N₄O₂ (750.98): calcd. C 81.57, H 6.71, N 7.46; found C 81.23, H 6.80, N 7.59.

Chloro[5,10-[2,2'-(dodecamethyleneoxy)diphenyl]-15-(4-methylphenyl)porphyrinato]manganese(III) (73): The free base porphyrin **72** (100 mg, 0.13 mmol) was dissolved in 50 mL DMF and heated with 60 mg MnCl₂·4 H₂O to reflux. After cooling to rt the mixture was filtered through neutral alumina (Brockman grade III). Column chromatography eluting with dichloromethane gave several minor fractions and the metal complex as the main fraction. Evaporation of the solvent followed by column chromatography eluting with neat dichloromethane gave the title compound (60 mg, 0.07 mmol, 55 %). M.p. >300 °C. UV/Vis (CH₂Cl₂): λ_{max} (log ε) = 380 (4.40), 405 (3.77), 487 (4.90), 597 (3.95), 629 nm (2.91). MS (FAB⁺, 3 kV): *m/z* = calcd. for C₅₁H₄₈ClMnN₄O₂ 838.28463; found 838.0 (7.0 %, [M]⁺), 803.0 (100.0 %, [M - Cl]⁺).

General procedure for bromination of 5,15-AB-type porphyrins: The porphyrin (1 equiv.) was dissolved in 100 mL of chloroform and cooled to 0 °C. Pyridine (1.0 mL) was added to act as an acid scavenger. NBS (1.1 equiv.) was added directly to the flask, and the reaction was followed by TLC. When the reaction reached completion it was quenched with 10 mL of acetone. The solvents were evaporated under reduced pressure, and the product was dissolved in CH₂Cl₂ followed by filtration through silica gel.

5,15-Dibromo-10-(*n*-butyl)-20-(2,4,6-trimethoxyphenyl)porphyrin (77): Prepared following the general bromination procedure given above using 5-butyl-15-(2,4,6-trimethoxyphenyl)porphyrin **22** (160 mg, 0.3 mmol), NBS (58.51 mg, 0.33 mmol) and acetone (10 mL). The products were separated using column chromatography (*n*-hexane/CH₂Cl₂ = 2:1, v/v). The first fraction was 5,15-dibromo-10-butyl-20-(2,4,6-trimethoxyphenyl)porphyrin **77** (90 mg, 0.13 mmol, 43 %) as purple crystals, the second fraction gave 5-bromo-10-butyl-20-(2,4,6-trimethoxyphenyl)porphyrin **76** (60 mg, 0.098 mmol, 33 %) as purple crystals. Data for **77**: M.p. > 300 °C. *R*_f = 0.5 (*n*-hexane/CH₂Cl₂, 1:1, v/v). ¹H NMR (600 MHz, CDCl₃, TMS): δ = -2.75 (s, 2H, NH), 1.13 (t, 3H, *J* = 7.53 Hz, CH₂CH₂CH₂CH₃), 1.77 (*m*, 2H, CH₂CH₂CH₂CH₃), 2.41 (*m*, 2H, CH₂CH₂CH₂CH₃), 3.57 (s, 6H, OCH₃), 4.15 (s, 3H, *p*-OCH₃), 4.72 (t, 2H, *J* = 8.28 Hz, CH₂CH₂CH₂CH₃), 6.62 (s, 2H, Ar-*H*), 8.82 (s, 2H, β-pyrrole-*H*), 9.31 (s, 2H, β-pyrrole-*H*), 9.56 (*d*, 2H, *J* = 4.52 Hz, β-pyrrole-*H*), 9.59 ppm (*d*, 2H, *J* = 4.15 Hz, β-pyrrole-*H*). ¹³C NMR (150 MHz, CDCl₃): δ = 14.0, 23.4, 29.5, 34.9, 40.7, 55.4, 55.8, 56.2, 90.6, 91.4, 91.8, 102.5, 111.4, 112.6, 121.7, 157.3, 160.3, 160.8, 162.0 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ε) = 422 (4.95), 522 (4.14), 556 (4.01), 603 (3.53), 661 nm (3.74). HRMS (ES⁺) calcd. for C₃₃H₃₁N₄O₃Br₂ [M+H]⁺ 691.0740; found 691.0742.

General procedure for Suzuki coupling: A Schlenk flask was charged with K₃PO₄ (20 equiv.) and anhydrous THF (60 mL) under an argon atmosphere. Then the porphyrin (1 equiv.), arylboronic acid or arylboronic ester (10 equiv.) and Pd(PPh₃)₄ (0.1 equiv.) were added. The reaction was heated at reflux temperature for 12-24 hours (TLC control) and protected from light. After completion, the solvent was removed under reduced pressure and the residue was dissolved in CH₂Cl₂. The mixture was washed with saturated NaHCO₃, H₂O, and brine and then dried over Na₂SO₄. The organic solvent was evaporated and the crude product was purified by flash chromatography.

5-(*n*-Butyl-10-(4-methoxycarbonylphenyl)-15-(2,4,6-trimethoxyphenyl)porphyrin (82): Using the above general Suzuki coupling procedure 5-bromo-10-(*n*-butyl)-20-(2,4,6-trimethoxyphenyl)porphyrin **76** (150 mg, 0.245 mmol), 4-methoxycarbonylphenyl boronic acid (440.9 mg, 2.45 mmol), Pd(PPh₃)₄ (28.31 mg, 0.0245 mmol) and K₃PO₄ (1041 mg, 4.9 mmol) in THF (50 mL) were used. Silica gel column chromatography (*n*-hexane/CH₂Cl₂ = 1:1, v/v) afforded the product as purple crystals (145 mg, 0.217 mmol, 89 %). M.p. > 300 °C. *R*_f = 0.42 (*n*-hexane/ethyl acetate, 10:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.83 (s, 2H, NH), 1.15 (t, 3H, *J* = 7.02 Hz, CH₂CH₂CH₂CH₃), 1.85 (*m*, 2H, CH₂CH₂CH₂CH₃), 2.56 (*m*, 2H, CH₂CH₂CH₂CH₃), 3.54 (s, 6H, OCH₃), 4.13 (s, 3H, *p*-OCH₃), 4.14 (s, 3H, OCH₃), 5.06 (t, 2H, *J* = 8.18 Hz, CH₂CH₂CH₂CH₃), 6.62 (s, 2H, Ar-*H*),

8.31 (*d*, 2H, *J* = 8.18 Hz, Ar-*H*), 8.45 (*d*, 2H, *J* = 8.19 Hz, Ar-*H*), 8.73 (*d*, 1H, *J* = 5.26 Hz, β-pyrrole-*H*), 8.86 (*d*, 2H, *J* = 4.68 Hz, β-pyrrole-*H*), 8.96 (*d*, 1H, *J* = 4.1 Hz, β-pyrrole-*H*), 9.11 (*m*, 2H, Ar-*H*), 9.27 (*d*, 1H, *J* = 4.1 Hz, β-pyrrole-*H*), 9.38 (*d*, 1H, *J* = 4.67 Hz, β-pyrrole-*H*), 9.51 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.61 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 10.10 ppm (s, 1H, meso-*H*). ¹³C NMR (100 MHz, CDCl₃): δ = 13.7, 23.2, 29.2, 34.5, 40.4, 51.9, 55.2, 90.4, 103.7, 110.6, 111.4, 117.2, 119.8, 126.8, 127.1, 127.7, 128.8, 129.7, 133.9, 147.5, 160.6, 161.5, 167.0 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ε) = 414 (5.00), 510 (4.00), 544 (3.52), 584 (3.60), 640 nm (3.42). HRMS (ES⁺) calcd. for C₄₁H₃₉N₄O₅ [M+H]⁺ 667.2920; found 667.2938.

5-(4-Ethynylphenyl)-15-hexyl-10,20-bis(3-methoxyphenyl)porphyrin (89): *n*-Butyllithium (2 mL of a 2.5 M solution in *n*-hexane, 2.5 mmol) was added under an argon atmosphere to a 50 mL Schlenk flask charged with a solution of *p*-bromophenylethyne (0.45 g, 2.5 mmol) in 10 mL of dry diethyl ether at -70 °C. The reaction mixture was then warmed to -40 °C and THF was added dropwise until the aryllithium was formed as a white-bright pink suspension. To this vigorously stirred mixture was added rapidly a solution of 5-hexyl-10,20-bis(3-methoxyphenyl)porphyrin **87** (50 mg, 0.082 mmol) in 30 mL of dry THF under an argon atmosphere. The mixture changed from deep purple to brown within 30 min. The reaction mixture was stirred for 2 h (TLC control). Subsequently, a mixture of 2 mL of water in 3 mL of THF was added for hydrolysis. After stirring of the mixture for 20 min, a solution of 10 equivalents of DDQ in THF (0.03 M) was added and the reaction mixture was stirred for another 60 min at room temperature. Subsequently, the mixture was filtered through silica gel and the organic solvent was removed under vacuum or washed with enough *n*-hexane. Final purification was achieved by column chromatography and elution with ethyl acetate/*n*-hexane (1 : 10, v/v) and yielded the title compound (7 mg, 0.1 mmol, 12 %) as purple crystals. M.p. >300 °C. *R*_f = 0.49 (ethyl acetate/*n*-hexane, 1:5, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.72 (s, 2H, NH), 0.94 (t, *J* = 7.2, 3H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.28 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.56 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.84 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.61 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 3.34 (s, 1H, HC≡C), 4.02 (s, 6H, OCH₃), 5.05 (t, 2H, *J* = 8.1, CH₂CH₂CH₂CH₂CH₂CH₃), 7.38 (*m*, 2H, Ph-*H*), 7.69 (*m*, 2H, Ph-*H*), 7.82 (*m*, 4H, Ph-*H*), 7.89 (*d*, 2H, *J* = 7.6, Ph-*H*), 8.19 (*d*, 2H, *J* = 7.6, Ph-*H*), 8.78 (*d*, 2H, *J* = 5.0 Hz, β-pyrrole-*H*_{2,8}), 8.88 (*d*, 2H, *J* = 5.0 Hz, β-pyrrole-*H*_{12,18}), 8.98 (*d*, 2H, *J* = 5.0 Hz, β-pyrrole-*H*_{3,7}), 9.51 ppm (*d*, 2H, *J* = 5.0 Hz, β-pyrrole-*H*_{13,17}). UV/Vis (CH₂Cl₂): λ_{max} (log ε) = 449 (5.10), 440 (3.97), 486 (3.32), 560 (2.15), 581 (3.05), 656 nm (4.29). HRMS calcd. for C₄₈H₄₂N₄O₂ 706.3308; found [M+1] 707.3365. C₄₈H₄₂N₄O₂ (706.89): calcd. C 81.56, H 5.99, N 7.93; found C 81.41, H 6.35, N 8.17.

5-(*sec*-Butyl)-15-hexyl-10,20-diphenylporphyrin (91): *sec*-Butyllithium (1 mL of a 2.5 M solution in hexane, 2.5 mmol) was added under an argon atmosphere to a 50 mL Schlenk flask charged with a solution of 5-hexyl-10,20-diphenylporphyrin **88** (50 mg, 0.09 mmol) in 30 mL of dry THF at -80 °C. The mixture changed from deep purple to brown within 30 min. The reaction mixture was stirred for 3 h (TLC control). Further workup followed the procedure for **89** and column chromatography and elution with ethyl acetate/*n*-hexane (1:8, v/v) yielded the title compound (13 mg, 0.02 mmol, 25 %) as purple crystals. M.p. >300 °C. *R*_f = 0.75 (ethyl acetate/*n*-hexane, 1:5, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.56 (s, 2H, NH), 0.91 (t, 3H, *J* = 7.6, CH₂CH₂CH₂CH₂CH₂CH₃), 1.10-1.31 (*m*, 5H, CH(CH₃)(CH₂CH₃), CH₂CH₂CH₂CH₂CH₂CH₃), 1.44 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.88 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.42 (*d*, 3H, *J* = 7.4 Hz, CH(CH₃)(CH₂CH₃)), 2.52 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 3.75 (*m*, 2H, CH(CH₃)(CH₂CH₃)), 4.98 (t, 2H, *J* = 7.6 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 5.32 (*m*, 1H, CH(CH₃)(CH₂CH₃)), 7.77 (*m*, 6H, Ph-*H*), 8.22 (*m*, 4H, Ph-*H*), 8.86 (*d*, 4H, *J* = 5.0 Hz, β-pyrrole-*H*_{2,8,12,18}), 9.43 (*d*, 2H, *J* = 5.0 Hz, β-pyrrole-*H*_{13,17}), 9.57 ppm (*d*, 2H, *J* = 5.0 Hz, β-pyrrole-*H*_{3,7}). UV/Vis (CH₂Cl₂): λ_{max} (log ε) = 441 (5.08), 486 (3.38), 514 (3.15), 581 (3.42), 656 nm (4.17). HRMS calcd. for C₄₂H₄₂N₄ 602.3409; found [M+1] 603.3487. C₄₂H₄₂N₄ (602.82): calcd. C 83.68, H 7.02, N 9.29; found C 83.17, H 7.20, N 9.50.

[5-(4-Ethynylphenyl)-10,20-diphenylporphyrinato]zinc(II) (94): The free base porphyrin 5-(4-ethynylphenyl)-10,20-diphenylporphyrin **93** (100 mg, 0.18 mmol) was dissolved in 100 mL dichloromethane. At room temperature 5 mL MeOH (dry) and 0.6 g of zinc(II) acetate were added and the mixture stirred for 30 min. After washing three times with water the organic phase was dried over Na₂SO₄, concentrated *in vacuo* and chromatography on alumina (dichloromethane/*n*-hexane, 1:5, v/v) to give 98 mg of **94** (1.5 mmol, 82 %) after recrystallization from CH₂Cl₂/MeOH. M.p. > 300 °C. *R*_f = 0.43 (dichloromethane/*n*-hexane, 1:5, v/v, alumina). ¹H NMR (250 MHz, CDCl₃, TMS): δ = 3.30 (s, 1H, HC≡C), 7.65-7.75 (*m*, 6H, Ph-*H*), 7.85 (*d*, 2H, ³*J* = 7.4, Ph-*H*), 8.20-8.30 (*m*, 6H, Ph-*H*), 8.80-9.0

(each d , 4H, $^3J = 5.0$, β -pyrrole- H), 9.20 (d , 2H, $^3J = 5.0$, β -pyrrole- H), 9.40 (d , 2H, $^3J = 5.0$, β -pyrrole- H), 10.20 ppm (s , 1H, meso- H). UV/vis (CH_2Cl_2): λ_{max} (log ϵ) = 413 (5.31), 539 (3.93), 562 nm (4.10). MS (EI, 80 eV, 170 °C), m/z (%): 626 (100) [M^+], 313 (7) [M^{2+}]. HRMS calcd. for $\text{C}_{40}\text{H}_{24}\text{N}_4\text{Zn}$ 626.1449; found 626.1452.

1,4-Di[4-phenyl-10,20-diphenylporphyrinato-5-yl]zinc(II)butane-1,3-diyne (95): Porphyrin **94** (50 mg, 0.08 mmol) was dissolved in 30 ml dichloromethane and CuCl (457 mg, 44.57 mmol) and 0.6 ml TMEDA were added slowly. The reaction mixture was stirred for 3 h at room temperature and washed 5 times with water (100 ml each). The organic phase was dried over Na_2SO_4 and the solvent removed *in vacuo*. Chromatographic work-up on alumina eluting with dichloromethane gave the dimer (41 mg pink crystals, 0.033 mmol, 82 % after recrystallization from $\text{CH}_2\text{Cl}_2/\text{MeOH}$). M.p. > 300 °C. $R_f = 0.76$ (dichloromethane/THF: 10:1, v/v, silica gel). ^1H NMR (250 MHz, CDCl_3 , TMS): $\delta = 7.60\text{--}7.65$ (d , 4H, $J = 7.6$, Ph- H), 7.70–7.80 (m , 12H, Ph- H), 7.90–7.95 (d , 4H, $J = 7.6$, Ph- H), 8.20–8.25 (d , 8H, $J = 7.4$, Ph- H), 8.80 (m , 8H, β -pyrrole- H), 9.05 (d , 4H, d , $^3J = 5.0$, β -pyrrole- H), 9.40 (d , 4H, d , $^3J = 5.0$, β -pyrrole- H), 10.25 ppm (s , 2H, meso- H). UV/vis (CH_2Cl_2): λ_{max} (log ϵ) = 416 (5.23), 460 (3.89), 544 nm (4.00). MS (EI, 80 eV, 170 °C), m/z (%): 1250 (43) [M^+], 600 (52) [$\text{M}^{2+} - \text{C}_2\text{H}_2$], 524 (64) [$\text{M}^{2+} - \text{C}_8\text{H}_6$]. FAB(+) calcd. for $\text{C}_{80}\text{H}_{47}\text{N}_8\text{Zn}_2$ 1251.57; found 1251.

5-(tert-Butyl)-5,24-dihydro-15-hexyl-10,20-diphenylporphyrin (96): *tert*-Butyllithium (1 ml of a 2.5 M solution in hexane, 2.5 mmol) was added under an argon atmosphere to a 50 ml Schlenk flask charged with a solution of 5-hexyl-10,20-diphenylporphyrin **88** (50 mg, 0.09 mmol) in 30 ml of dry THF at –80 °C. The reaction mixture was stirred for 5 h (TLC control). Subsequently, a mixture of 2 ml of water in 3 ml of THF was added for hydrolysis. After stirring of the mixture for 20 min, a solution of 10 equivalents of DDQ in THF (0.03 M) was added and the reaction mixture was stirred for another 60 min at room temperature. Subsequently, the mixture was filtered through silica gel and the organic solvent was removed under vacuum. Final purification was achieved by column chromatography and elution with ethyl acetate/*n*-hexane (1:6, v/v) yielding the title compound (11 mg, 0.02 mmol, 20%) as purple crystals. M.p. = 227 °C. $R_f = 0.43$ (ethyl acetate/*n*-hexane, 1:1, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): $\delta = 0.91$ (t , 3H, $J = 6$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.09 (m , 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.28 (s , 9H, $\text{C}(\text{CH}_3)_3$), 1.44 (m , 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.88 (m , 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.47 (m , 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 4.13 (t , 2H, $J = 6$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 5.32 (s , 1H, 5- H), 6.04–6.47 (m , 4H, β -pyrrole- H), 6.51–6.85 (m , 4H, β -pyrrole- H), 7.14–7.59 (m , 6H, Ph- H), 7.87–7.96 (m , 4H, Ph- H), 10.39, 10.57 and 10.94 ppm (s each, 3H, NH). UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 376 (5.03), 435 (4.82), 578 (4.84), 656 (4.17) nm.^[31]

5-(tert-Butyl)-5,24-dihydro-10,20-bis(3-methoxyphenyl)-15-phenylporphyrin (97): *tert*-Butyllithium (2 ml of a 1.8 M solution in hexane, 0.06 mmol) was slowly added (ca. 1 h) under an argon atmosphere to a 50 ml Schlenk flask charged with a solution of 5-phenyl-10,20-bis(3-methoxyphenyl)porphyrin **59** (35 mg, 0.06 mmol) in 30 ml of dry THF. The mixture was heated to 50 °C and stirred for 30 min. Subsequently, a mixture of 2 ml of water in 3 ml of THF was added for hydrolysis. After stirring of the mixture for 20 min, a solution of 10 equivalents of DDQ in THF (0.02 M) was added and the reaction mixture was stirred for another 60 min at room temperature. Subsequently, the mixture was filtered through silica gel and the organic solvent was removed under vacuum or washed with *n*-hexane. Final purification was achieved by column chromatography and elution with ethyl acetate/*n*-hexane (1:8, v/v) and gave the title compound (11 mg, 0.016 mmol, 28 %) as purple crystals. M.p. = 226 °C. $R_f = 0.47$ (ethyl acetate/*n*-hexane, 1:1, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): $\delta = 1.28$ (s , 9H, $\text{C}(\text{CH}_3)_3$), 3.91 (s , 6H, OCH_3), 5.33 (s , 1H, 5- H), 6.01–6.43 (m , 4H, β -pyrrole- H), 6.53–6.66 (m , 4H, β -pyrrole- H), 6.87 (m , 2H, Ar- H), 7.06–7.15 (m , 4H, Ar- H), 7.37 (m , 3H, Ar- H), 7.51 (m , 4H, Ar- H), 9.88, 10.02 and 10.65 ppm (s each, 3H, NH). UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 350 (4.85), 453 (4.69), 577 (5.11), 656 (4.15) nm.^[31]

5-Hexyl-10,20-bis(3-methoxyphenyl)-15-[(3-pyridyl)methyl]porphyrin (98): *n*-Hexyllithium (1 ml of a 2.5 M solution in hexane) was slowly added under an argon atmosphere to a 100 ml Schlenk flask charged with a solution of 5,15-bis(3-methoxyphenyl)porphyrin **58** (100 mg, 0.19 mmol) in 40 ml of dry THF at –80 °C under an argon atmosphere. After 15 min the solution was treated with a solution of 300 mg (0.86 mmol) 3-(iodomethyl)pyridine hydrogen iodide dissolved in 4 ml DMF and stirred for 24 h and heated to 75 °C (TLC control). Subsequently, a mixture of 2 ml of water in 3 ml of THF was added for hydrolysis. After stirring of the mixture for 20 min, a solution of 10 equivalents of DDQ in THF (0.04 M) was added and the reaction mixture was stirred for another 60 min at room

temperature. Subsequently, the mixture was filtered through silica gel and the organic solvent was removed under vacuum. Final purification was achieved by column chromatography on silica gel and elution with ethyl acetate/*n*-hexane (1:8, v/v) and yielded the title compound (8 mg, 0.01 mmol, 6 %) as purple crystals and the trisubstituted porphyrin (28 %). Data for **98**: M.p. >300 °C. $R_f = 0.35$ (ethyl acetate/*n*-hexane, 1:1, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): $\delta = -2.62$ (s , 2H, NH), 0.91 (t , $J = 7.2$, 3H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.23 (m , 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.54 (m , 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.83 (m , 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.55 (m , 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 4.03 (s , 6H, OCH_3), 4.99 (t , 2H, $J = 8.1$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 6.38 (s , 2H, pyridyl- CH_2), 7.01 (m , 2H, pyridyl- H), 7.35 (m , 3H, 2 Ar- H , 1 pyridyl- H), 7.66 (m , 2H, Ar- H), 7.79 (m , 4H, Ar- H), 8.38 (s , 1H, pyridyl- H), 8.95 (m , 4H, β -pyrrole- $H_{2,8,12,18}$), 9.33 (d , 2H, $J = 5.0$ Hz, β -pyrrole- $H_{13,17}$), 9.48 ppm (d , 2H, $J = 5.0$ Hz, β -pyrrole- $H_{3,7}$). UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 420 (5.07), 485 (3.74), 516 (3.62), 551 (2.92), 593 (3.04), 655 nm (4.00). HRMS calcd. for $\text{C}_{46}\text{H}_{43}\text{N}_5\text{O}_2$ 697.3416; found [$\text{M} + 1$] 698.3516. $\text{C}_{46}\text{H}_{43}\text{N}_5\text{O}_2$ (697.88): calcd. C 79.17, H 6.21, N 10.04; found C 79.44, H 6.51, N 9.92.

3-(5,15-Dihexyl-10-phenylporphyrin-20-yl)propionic acid (99): Phenyllithium (0.6 ml of a 1.8 M solution in hexane, 0.018 mmol) was slowly added under an argon atmosphere to a 50 ml Schlenk flask charged with a solution of 5,15-dihexylporphyrin **18** (80 mg, 0.167 mmol) in 30 ml of dry THF at R.T. under an argon atmosphere. After 15 min the solution was treated with a solution of 200 mg (1 mmol) 3-iodopropionic acid dissolved in 5 ml THF and stirring for 24 h (TLC control). Subsequently, a mixture of 2 ml of water in 3 ml of THF was added for hydrolysis. After stirring of the mixture for 20 min, a solution of 10 equivalents of DDQ in THF (0.03 M) was added and the reaction mixture was stirred for another 60 min at room temperature. Subsequently, the mixture was filtered through silica gel and the organic solvent was removed under vacuum. Final purification was achieved by column chromatography on silica gel and elution with ethyl acetate/*n*-hexane (1:10, v/v) yielded the title compound (19 mg, 0.03 mmol, 18 %) as purple crystals and the respective trisubstituted porphyrin (27 %). Data for **99**: M.p. >300 °C. $R_f = 0.47$ (ethyl acetate/*n*-hexane, 1:7, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): $\delta = -2.65$ (s , 2H, NH), 0.91 (t , 6H, $J = 7.2$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.20 (m , 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.55 (m , 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.86 (m , 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.57 (m , 6H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{COOH}$), 3.72 (t , 2H, $J = 6.4$, $\text{CH}_2\text{CH}_2\text{COOH}$), 4.98 (t , 4H, $J = 8.1$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 7.77 (m , 3H, Ph- H), 8.21 (m , 2H, Ph- H), 8.88 (m , 4H, β -pyrrole- $H_{3,7,13,17}$), 9.44 (m , 4H, β -pyrrole- $H_{2,8,12,18}$), 10.56 ppm (s , 1H, COOH). UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 418 (5.08), 440 (3.94), 517 (3.79), 552 (3.91), 593 (3.88), 655 nm (4.09). HRMS calcd. for $\text{C}_{41}\text{H}_{46}\text{N}_4\text{O}_2$ 626.3621; found 626.4512. $\text{C}_{41}\text{H}_{46}\text{N}_4\text{O}_2$ (626.84): calcd. C 78.56, H 7.40, N 8.94; found C 78.76, H 7.80, N 9.21.

5,15-Dihexyl-10-[4-(dipropylamino)phenyl]-20-propylporphyrin (101) and 5,15-dihexyl-10-[4-(dipropylamino)phenyl]-porphyrin (102): *n*-Butyllithium (4 ml of a 2.5 M solution in hexane, 10 mmol) was added under an argon atmosphere to a 100 ml Schlenk flask charged with a solution of *p*-bromoaniline (1 g, 5 mmol) in 10 ml of dry diethyl ether at 0 °C. After addition of *n*-BuLi the cold bath was removed and stirring was continued for another 2 h at room temperature. To the vigorously stirred mixture was added rapidly a solution of 5,15-dihexylporphyrin **18** (150 mg, 0.313 mmol) in 40 ml of dry THF. The mixture changed from deep purple to brown within 30 min. After 1 h the solution was treated with 1 ml propyl iodide (10 mmol) and stirred for 12 h and heated to 70 °C (TLC control). Subsequently, a mixture of 2 ml of water in 3 ml of THF was added for hydrolysis. After stirring of the mixture for 20 min, a solution of 10 equivalents of DDQ in THF (0.03 M) was added and the reaction mixture was stirred for another 60 min at room temperature. Subsequently, the mixture was filtered through silica gel and the organic solvent was removed under vacuum. Final purification was achieved by column chromatography on silica gel and elution with ethyl acetate/*n*-hexane (1 : 10, v/v). The first fraction consisted of **101** (43 mg, 0.06 mmol, 20 %) and the second fraction of **102** (34 mg, 0.05 mmol, 16 %). Data for **101**: purple crystals. M.p. >300 °C. $R_f = 0.63$ (ethyl acetate/*n*-hexane, 1:6, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): $\delta = -2.57$ (s , 2H, NH), 0.92 (t , 6H, $J = 7.2$ Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.11 (m , 6H, $\text{N}(\text{CH}_2\text{CH}_2\text{CH}_3)_2$), 1.25 (m , 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.37–1.51 (m , 7H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_3$), 1.84–1.93 (m , 8H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{N}(\text{CH}_2\text{CH}_2\text{CH}_3)_2$), 2.56 (m , 6H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_3$), 3.55 (m , 4H, $\text{N}(\text{CH}_2\text{CH}_2\text{CH}_3)_2$), 4.99 (m , 6H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{COOH}$), 7.04 (d , 2H, $J = 7.5$ Hz, Ar- H), 8.05 (d , 2H, $J = 7.5$ Hz, Ar- H), 9.02–9.11 (m , 4H, β -pyrrole- H), 9.45 ppm (m , 4H, β -pyrrole- H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 11.22$, 13.76, 14.59, 20.13, 20.24, 22.34, 29.22, 29.85, 31.27, 31.52, 32.63, 34.56, 35.03, 37.15, 38.16, 38.21, 50.71, 52.74, 109.24, 109.44, 117.84, 118.35, 118.57, 126.48, 127.55, 128.86,

146.52 ppm. UV/Vis (CH₂Cl₂): $\lambda_{\max}$ (log ϵ) = 436 (5.10), 486 (4.32), 511 (4.29), 581 (3.64), 656 nm (4.28). HRMS calcd. for C₄₇H₆₁N₅ 695.4926; found [M + 1] 696.5008. C₄₇H₆₁N₅ (696.03): calcd. C 81.09, H 8.84, N 10.07; found C 81.29, H 8.59, N 9.98. Data for **102**: purple crystals. M.p. >300 °C. R_f = 0.55 (ethyl acetate/*n*-hexane, 1:6, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.75 (s, 2H, NH), 0.92 (t, 6H, J = 7.2, CH₂CH₂CH₂CH₂CH₂CH₃), 1.14 (m, 6H, N(CH₂CH₂CH₃)₂), 1.28 (m, 4H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.37-1.51 (m, 4H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.85-1.94 (m, 8H, CH₂CH₂CH₂CH₂CH₂CH₃, N(CH₂CH₂CH₃)₂), 2.58 (m, 4H, CH₂CH₂CH₂CH₂CH₂CH₃), 3.56 (m, 4H, N(CH₂CH₂CH₃)₂), 5.02 (m, 4H, CH₂CH₂CH₂CH₂CH₂CH₃), 7.04 (d, 2H, J = 7.5 Hz, Ar-H), 8.04 (d, 2H, J = 7.5 Hz, Ar-H), 8.99 (d, 2H, J = 5.0 Hz, β -pyrrole-H), 9.34 (d, 2H, J = 5.0 Hz, β -pyrrole-H), 8.44 (d, 2H, J = 5.0 Hz, β -pyrrole-H), 9.56 (d, 2H, J = 5.0 Hz, β -pyrrole-H), 10.03 ppm (s, 1H, meso-H). ¹³C NMR (100 MHz, CDCl₃): δ = 11.22, 13.74, 20.23, 22.31, 29.29, 29.82, 31.51, 34.55, 38.15, 52.75, 102.72, 109.26, 118.57, 120.45, 126.46, 127.58, 129.33, 130.91, 131.91, 135.36, 144.73, 147.23 ppm. UV/Vis (CH₂Cl₂): $\lambda_{\max}$ (log ϵ) = 434 (5.11), 585 (3.78), 656 nm (4.11). HRMS calcd. for C₄₄H₅₅N₅ 653.4457; found [M + 1] 653.4491. C₄₄H₅₅N₅ (653.95): calcd. C 80.81, H 8.48, N 10.71; found C 81.09, H 8.71, N 10.43.

5,10-Dihexyl-15-[4-(dihexyl-hydroxymethyl)phenyl]-20-propylporphyrin (103) and 5,10-dihexyl-15-[4-(dihexyl-hydroxymethyl)phenyl]-porphyrin (104): *n*-Hexyllithium (2 ml of a 2.5 M solution in hexane, 5 mmol) was added under an argon atmosphere to a 100 ml Schlenk flask charged with a solution of 5-hexyl-15-(4-methylcarboxyphenyl)porphyrin **17** (40 mg, 0.07 mmol) in 40 ml of dry THF under an argon atmosphere. The mixture changed from deep purple to brown within 30 min. After 1 h the solution was treated with 0.5 ml propyl iodide (0.52 mmol) and stirring for 12 h and heating to 70 °C (TLC control). Subsequently, a mixture of 2 ml of water in 3 ml of THF was added for hydrolysis. After stirring of the mixture for 20 min, a solution of 10 equivalents of DDQ in THF (0.02 M) was added and the reaction mixture was stirred for another 60 min at room temperature. The mixture was filtered through silica gel and the organic solvent was removed under vacuum. Final purification was achieved by column chromatography and elution with acetone/*n*-hexane (1:5, v/v). The first fraction consisted of **103** (11 mg, 0.013 mmol, 18 %) and the second fraction of **104** (8 mg, 0.01 mmol, 14 %). Data for **103**: Purple crystals. M.p. >300 °C. R_f = 0.31 (acetone/*n*-hexane, 1:5, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.60 (s, 2H, NH), 0.96-1.15 (t, 12H, J = 7.2, CH₂CH₂CH₂CH₂CH₂CH₃, C(OH)CH₂CH₂CH₂CH₂CH₂CH₃), 1.23 (m, 8H, CH₂CH₂CH₂CH₂CH₂CH₃), C(OH)CH₂CH₂CH₂CH₂CH₂CH₃), 1.41-1.65 (m, 16H, CH₂CH₂CH₂CH₂CH₂CH₃, C(OH)CH₂CH₂CH₂CH₂CH₂CH₃, C(OH)CH₂CH₂CH₂CH₂CH₂CH₃, C(OH)CH₂CH₂CH₂CH₂CH₂CH₃), 1.81-1.91 (m, 7H, CH₂CH₂CH₂CH₂CH₂CH₃, CH₂CH₂CH₃), 2.12 (m, 4H, C(OH)CH₂CH₂CH₂CH₂CH₂CH₃), 2.55 (m, 6H, CH₂CH₂CH₂CH₂CH₂CH₃, CH₂CH₂CH₃), 4.98 (m, 6H, CH₂CH₂CH₂CH₂CH₂CH₃, CH₂CH₂CH₃), 7.78 (d, 2H, J = 7.5 Hz, Ar-H), 8.17 (m, 6H, Ar-H), 8.87 (d, 2H, J = 5.0 Hz, β -pyrrole-H_{13,17}), 9.43 (m, 2H, β -pyrrole-H_{2,18}), 9.54 ppm (m, 4H, β -pyrrole-H_{3,7,8,12}). ¹³C NMR (100 MHz, CDCl₃): δ = 13.67, 22.20, 22.38, 23.29, 28.99, 29.31, 31.52, 35.01, 35.38, 36.84, 38.28, 38.47, 42.82, 117.61, 118.28, 118.68, 118.90, 123.10, 126.83, 127.35, 127.88, 133.74, 140.02, 145.33 ppm. UV/Vis (CH₂Cl₂): $\lambda_{\max}$ (log ϵ) = 418 (5.18), 519 (3.50), 553 (3.78), 601 (3.88), 655 nm (3.75). HRMS calcd. for C₅₄H₇₄N₄O 794.5862; found 794.5725. Data for **104**: Purple crystals. M.p. >300 °C. R_f = 0.31 (acetone/*n*-hexane, 1:5, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.88 (s, 2H, NH), 0.93-1.15 (t, 12H, J = 7.2 Hz, CH₂CH₂CH₂CH₂CH₂CH₃, C(OH)CH₂CH₂CH₂CH₂CH₂CH₃), 1.23 (m, 8H, CH₂CH₂CH₂CH₂CH₂CH₃, C(OH)CH₂CH₂CH₂CH₂CH₂CH₃), 1.41-1.73 (m, 16H, CH₂CH₂CH₂CH₂CH₂CH₃, C(OH)CH₂CH₂CH₂CH₂CH₂CH₃, C(OH)CH₂CH₂CH₂CH₂CH₂CH₃, C(OH)CH₂CH₂CH₂CH₂CH₂CH₃), 1.82-1.90 (m, 4H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.09 (m, 4H, C(OH)CH₂CH₂CH₂CH₂CH₂CH₃), 2.57 (m, 4H, CH₂CH₂CH₂CH₂CH₂CH₃), 5.06 (m, 4H, CH₂CH₂CH₂CH₂CH₂CH₃), 7.79 (d, 2H, J = 7.5 Hz, Ar-H), 8.21 (d, 2H, J = 7.5 Hz, Ar-H), 8.87 (d, 2H, J = 5.0 Hz, β -pyrrole-H_{13,17}), 9.34 (d, 2H, β -pyrrole-H_{2,18}), 9.54 (m, 4H, β -pyrrole-H_{3,7,8,12}), 10.03 ppm (s, 1H, meso-H). ¹³C NMR (100 MHz, CDCl₃): δ = 13.66, 22.18, 29.29, 29.88, 29.90, 31.31, 34.87, 35.66, 38.37, 38.64, 42.80, 103.03, 117.77, 119.06, 120.08, 123.29, 127.81, 130.38, 133.91, 138.72, 139.38, 145.41 ppm. UV/Vis (CH₂Cl₂): $\lambda_{\max}$ (log ϵ) = 435 (5.10), 486 (3.11), 514 (2.57), 581 (3.42), 656 nm (3.72). HRMS calcd. for C₅₁H₆₈N₄O 752.5393; found 752.4981.

5-Hexyl-10-phenyl-15-(4-methoxyphenyl)-20-propylporphyrin (105): PhLi (0.8 ml of a 1.8 M solution in hexane, 0.02 mmol) was slowly added (ca. 1 h) under an argon atmosphere to a 100 ml Schlenk flask charged with a solution of 5-hexyl-15-(4-methoxyphenyl)porphyrin **105** (100 mg, 0.19 mmol) in 40 ml of dry THF. The mixture was heated to 50 °C and stirred

for 30 min. After 1 h the solution was treated with 0.7 propyl iodide (7.1 mmol) and stirring for 20 h and heating to 70 °C (TLC control). Subsequently, a mixture of 2 ml of water in 3 ml of THF was added for hydrolysis. After stirring of the mixture for 20 min, a solution of 10 equivalents of DDQ in THF (0.06 M) was added and the reaction mixture was stirred for another 60 min at room temperature. Subsequently, the mixture was filtered through silica gel and the organic solvent was removed under vacuum. Final purification was achieved by column chromatography (ethyl acetate/*n*-hexane = 1:10, v/v) yielded the title compound (31 mg, 0.05 mmol, 25 %) as purple crystals, besides the respective trisubstituted porphyrin (14 %) and starting material (8 %). M.p. >300 °C. R_f = 0.76 (ethyl acetate/*n*-hexane, 1:5, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.61 (s, 2H, NH), 0.99 (t, 6H, J = 8.1 Hz, CH₂CH₂CH₂CH₂CH₂CH₃, CH₂CH₂CH₃), 1.34 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.41 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.86 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.57 (m, 4H, CH₂CH₂CH₂CH₂CH₂CH₃, CH₂CH₂CH₃), 4.09 (s, 3H, OCH₃), 5.02 (t, 4H, J = 8.1 Hz, CH₂CH₂CH₂CH₂CH₂CH₃, CH₂CH₂CH₃), 7.25 (d, 2H, J = 8 Hz, Ph-H), 7.79 (m, 3H, Ph-H), 8.15 (d, 2H, J = 8 Hz, Ph-H), 8.27 (m, 2H, Ph-H), 8.81 (m, 2H, β -pyrrole-H_{13,17}), 8.98 (m, 2H, β -pyrrole-H_{2,18}), 9.43 (m, 2H, β -pyrrole-H_{3,12}), 9.56 (m, 2H, β -pyrrole-H_{7,8}) ppm. ¹³C NMR (100 MHz, CDCl₃): δ = 13.79, 14.58, 22.36, 29.33, 29.88, 31.33, 31.53, 35.25, 37.10, 38.45, 55.12, 111.71, 118.16, 119.24, 119.49, 126.22, 126.75, 127.13, 128.35, 134.07, 135.14, 141.99, 158.86 ppm. UV/Vis (CH₂Cl₂): $\lambda_{\max}$ (log ϵ) = 444 (4.97), 487 (4.30), 601 (4.03), 657 nm (4.28). HRMS: calcd. for C₄₂H₄₂N₄O 618.3358; found [M + 1] 619.3425. C₄₂H₄₂N₄O (618.82): calcd. C 81.52, H 6.84, N 9.05; found C 81.63, H 7.05, N 8.97.

5-Butyl-10-hexyl-15-pentyl-20-(4-methoxyphenyl)porphyrin (107): *n*-BuLi (0.4 ml of 2.5 M solution in hexane, 1 mmol) was slowly added under an argon atmosphere to a 100 ml Schlenk flask charged with a solution of 5-hexyl-15-(4-methoxyphenyl)porphyrin **51** (100 mg, 0.19 mmol) in 40 ml of dry THF at -80 °C under an argon atmosphere. After 15 min the solution was treated with 0.8 ml *n*-pentyl iodide (6.1 mmol) and stirred for 12 h (TLC control). Subsequently, a mixture of 2 ml of water in 3 ml of THF was added for hydrolysis. After stirring of the mixture for 20 min, a solution of 10 equivalents of DDQ in THF (0.03 M) was added and the reaction mixture was stirred for another 60 min at room temperature. Subsequently, the mixture was filtered through silica gel and the organic solvent was removed under vacuum. Final purification was achieved by column chromatography and elution with ethyl acetate/*n*-hexane (1:15, v/v) yielded the title compound (15 mg, 0.023 mmol, 13 %) as purple crystals, trisubstituted porphyrin (9 %) and starting material (7 %). M.p. >300 °C. R_f = 0.57 (ethyl acetate/*n*-hexane, 1:5, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.63 (s, 2H, NH), 0.87 (m, 6H, CH₂CH₂CH₂CH₂CH₂CH₃, CH₂CH₂CH₂CH₂CH₂CH₃), 1.37 (m, 5H, CH₂CH₂CH₂CH₂CH₂CH₃, CH₂CH₂CH₂CH₂CH₃), 1.56 (m, 4H, CH₂CH₂CH₂CH₂CH₂CH₃, CH₂CH₂CH₂CH₂CH₃), 1.86 (m, 6H, CH₂CH₂CH₂CH₂CH₂CH₃, CH₂CH₂CH₂CH₂CH₃), 2.56 (m, 6H, CH₂CH₂CH₂CH₂CH₂CH₃, CH₂CH₂CH₂CH₂CH₃, CH₂CH₂CH₂CH₂CH₃, CH₂CH₂CH₂CH₂CH₃), 4.13 (s, 3H, OCH₃), 5.02 (m, 6H, CH₂CH₂CH₂CH₂CH₂CH₃, CH₂CH₂CH₂CH₂CH₃, CH₂CH₂CH₂CH₂CH₃), 7.33 (d, 2H, J = 8 Hz, Ar-H), 8.11 (d, 2H, J = 8 Hz, Ar-H), 8.85 (d, 2H, J = 5 Hz, β -pyrrole-H), 9.40 (d, 2H, J = 5 Hz, β -pyrrole-H), 9.52 (d, 2H, J = 5 Hz, β -pyrrole-H), 9.55 ppm (d, 2H, J = 5 Hz, β -pyrrole-H). UV/Vis (CH₂Cl₂): $\lambda_{\max}$ (log ϵ) = 442 (5.09), 486 (3.51), 548 (3.27), 581 (3.51), 656 nm (4.27). HRMS: calcd. for C₄₂H₅₀N₄O 626.3984; found [M + 1] 627.4061. C₄₂H₅₀N₄O (626.89): calcd. C 80.47, H 8.04, N 8.94; found C 80.67, H 6.95, N 8.63.

5-[4-(Dipropylamino)phenyl]-10-hexyl-20-(4-methoxyphenyl)porphyrin (109): *n*-Butyllithium (2 ml of a 2.5 M solution in hexane, 5 mmol) was added under an argon atmosphere to a 100 ml Schlenk flask charged with a solution of *p*-bromoaniline (0.5 g, 2.5 mmol) in 10 ml of dry diethyl ether at 0 °C. After addition of *n*-BuLi the cold bath was removed and stirring was continued for another 2 h at room temperature. To the vigorously stirred mixture was added rapidly a solution of 5-hexyl-15-(4-methoxyphenyl)porphyrin **51** (100 mg, 0.19 mmol) in 40 ml of dry THF under an argon atmosphere at -80 °C. The mixture changed from deep purple to brown within 30 min. After 1 h the solution was treated with 1 ml propyl iodide (10 mmol) and stirring for 12 h and heating to 70 °C (TLC control). Subsequently, a mixture of 2 ml of water in 3 ml of THF was added for hydrolysis. After stirring of the mixture for 20 min, a solution of 10 equivalents of DDQ in THF (0.03 M) was added and the reaction mixture was stirred for another 60 min at room temperature. The mixture was filtered through silica gel and the organic solvent was removed under vacuum. Final purification was achieved by column chromatography and elution with ethyl acetate/*n*-hexane (1:10, v/v) yielding the title compound (41 mg, 0.06 mmol, 31 %) as purple crystals; some starting material (13 %) was recovered, too. M.p. >300 °C. R_f = 0.54 (ethyl acetate/*n*-hexane, 1:8, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.81 (s, 2H, NH), 0.92 (t,

3H, $J = 7.2$ Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.13 (*m*, 6H, $\text{N}(\text{CH}_2\text{CH}_2\text{CH}_3)_2$), 1.27 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.57 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.83–1.94 (*m*, 6H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), $\text{N}(\text{CH}_2\text{CH}_2\text{CH}_3)_2$), 2.61 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 3.54 (*m*, 4H, $\text{N}(\text{CH}_2\text{CH}_2\text{CH}_3)_2$), 4.12 (*s*, 3H, OCH_3), 5.04 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 7.05 (*d*, 2H, $J = 7.5$ Hz, *Ar-H*), 7.34 (*d*, 2H, $J = 7.5$ Hz, *Ar-H*), 8.05 (*d*, 2H, $J = 7.5$ Hz, *Ar-H*), 8.15 (*d*, 2H, $J = 7.5$ Hz, *Ar-H*), 8.88–9.01 (*d*, 2H, $J = 5.0$ Hz, β -pyrrole-*H*), 9.02–9.13 (*d*, 2H, $J = 5.0$ Hz, β -pyrrole-*H*), 9.28–9.37 (*d*, 2H, $J = 5.0$ Hz, β -pyrrole-*H*), 9.51–9.61 (*d*, 2H, $J = 5.0$ Hz, β -pyrrole-*H*), 10.09 ppm (*s*, 1H, *meso-H*). UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 443 (5.09), 486 (3.81), 585 (3.67), 656 nm (4.22). HRMS calcd. for $\text{C}_{45}\text{H}_{49}\text{N}_5\text{O}$ 675.3937; found $[\text{M} + 1]^+$ 676.4039, $\text{C}_{45}\text{H}_{49}\text{N}_5\text{O}$ (675.92): calcd. C 79.96, H 7.31, N 10.36; found C 79.65, H 7.05, N 10.01.

5-Hexyl-10-phenyl-15-(4-methoxyphenyl)porphyrin (110):

Phenyllithium (0.8 ml of a 1.8 M solution in hexane, 0.02 mmol) was slowly added (ca. 1 h) under an argon atmosphere to a 100 ml Schlenk flask charged with a solution of 5-hexyl-15-(4-methoxyphenyl)porphyrin **51** (100 mg, 0.19 mmol) in 40 ml of dry THF. The mixture was heated to 50 °C and stirred for 30 min. Subsequently, a mixture of 2 ml of water in 3 ml of THF was added for hydrolysis. After stirring of the mixture for 20 min, a solution of 10 equivalents of DDQ in THF (0.06 M) was added and the reaction mixture was stirred for another 60 min at room temperature. The mixture was filtered through silica gel and the organic solvent was removed under vacuum. Column chromatography (ethyl acetate/*n*-hexane = 1:10, v/v) gave the title compound (**51** mg, 0.088 mmol, 45 %) as purple crystals besides starting material (8 %). M.p. >300 °C. $R_f = 0.69$ (ethyl acetate/*n*-hexane, 1:5, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): $\delta = -2.87$ (*s*, 2H, *NH*), 0.95 (*t*, 3H, $J = 8.1$ Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.32 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.53 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.85 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.56 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 4.11 (*s*, 3H, OCH_3), 5.05 (*t*, 2H, $J = 8.1$ Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 7.31 (*d*, 2H, $J = 8$ Hz, *Ph-H*), 7.82 (*m*, 3H, *Ph-H*), 8.16 (*d*, 2H, $J = 8$ Hz, *Ph-H*), 8.25 (*m*, 2H, *Ph-H*), 8.85–8.91 (*d*, 2H, $J = 5$ Hz, β -pyrrole-*H*), 8.96–9.03 (*d*, 2H, $J = 5$ Hz, β -pyrrole-*H*), 9.31–9.39 (*d*, 2H, $J = 5$ Hz, β -pyrrole-*H*), 9.51–9.61 (*d*, 2H, $J = 5$ Hz, β -pyrrole-*H*), 10.14 ppm (*s*, 1H, *meso-H*). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 13.74$ (10^3-C), 22.31 (10^2-C), 29.85 (10^4-C), 31.50 (10^3-C), 34.73 (10^2-C), 38.37 (10^1-C), 55.13 (COCH_3), 103.82, 111.93, 118.26, 119.45, 119.61, 126.04, 127.03, 127.20, 127.93, 130.61, 130.75, 131.09, 131.37, 133.63, 134.01, 135.23, 142.43, 142.64, 158.93 ppm. UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 441 (4.97), 548 (2.76), 596 (3.34), 644 nm (3.95). HRMS calcd. for $\text{C}_{39}\text{H}_{36}\text{N}_4\text{O}$ 576.2889; found $[\text{M} + 1]^+$ 577.2946. $\text{C}_{39}\text{H}_{36}\text{N}_4\text{O}$ (576.74): calcd. C 81.22, H 6.29, N 9.71; found C 81.47, H 6.32, N 9.58.

5-Butyl-10-hexyl-15-(4-hydroxyphenyl)-20-(4-methoxyphenyl)porphyrin (115):

n-Butyllithium (2 ml of a 2.5 M solution in hexane, 10 mmol) was added under an argon atmosphere to a 50 ml Schlenk flask charged with a solution of *p*-bromophenol (0.43 g, 2.5 mmol) in 10 ml of dry diethyl ether at 0 °C. After addition of *n*-BuLi the cold bath was removed and stirring was continued for 18 h at room temperature. To the vigorously stirred mixture was added rapidly a solution of 5-butyl-10-hexyl-20-(4-methoxyphenyl)porphyrin **114** (40 mg, 0.07 mmol) in 30 ml of dry THF under an argon atmosphere. Subsequently, a mixture of 2 ml of water in 3 ml of THF was added for hydrolysis. After stirring of the mixture for 20 min, a solution of 10 equivalents of DDQ in THF (0.03 M) was added and the reaction mixture was stirred for another 60 min at room temperature. Subsequently, the mixture was filtered through silica gel and the organic solvent was removed under vacuum. Final purification was achieved by column chromatography and elution with ethyl acetate/*n*-hexane (1:8, v/v) yielded the title compound (28 mg, 0.04 mmol, 60 %) as purple crystals; 8 % starting material was recovered. M.p. >300 °C. $R_f = 0.52$ (ethyl acetate/*n*-hexane, 1:2, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): $\delta = -2.66$ (*s*, 2H, *NH*), 0.95 (*t*, 3H, $J = 7.5$ Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.15 (*t*, 3H, $J = 7.5$ Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.29 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.53 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.89 (*m*, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.57 (*m*, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 4.11 (*s*, 3H, OCH_3), 5.01 (*t*, 4H, $J = 7.8$ Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 7.15 (*d*, 2H, $J = 7.5$ Hz, *Ar-H*), 7.30 (*d*, 2H, $J = 7.5$ Hz, *Ar-H*), 8.04 (*d*, 2H, $J = 7.5$ Hz, *Ar-H*), 8.11 (*d*, 2H, $J = 7.5$ Hz, *Ar-H*), 8.78 (*s*, 2H, β -pyrrole-*H*), 8.97 (*d*, 2H, $J = 5$ Hz, β -pyrrole-*H*), 9.45 (*d*, 2H, $J = 5$ Hz, β -pyrrole-*H*), 9.59 ppm (*s*, 2H, β -pyrrole-*H*). UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 446 (5.08), 486 (3.36), 581 (3.15), 615 (3.39), 656 nm (4.24). HRMS calcd. for $\text{C}_{43}\text{H}_{44}\text{N}_4\text{O}_2$ 648.3464; found 648.2985. $\text{C}_{43}\text{H}_{44}\text{N}_4\text{O}_2$ (648.85): calcd. C 79.60, H 6.84, N 8.63; found C 79.99, H 7.05, N 8.58.

5-(*n*-Butyl)-15-[4-(*N,N*-dimethylamino)phenyl]-10-(1-ethylpropyl)-20-hexylporphyrin (121):

The synthesis followed the standard procedures

using *n*-BuLi (0.03 mL, 0.09 mmol of a 2.5 M solution in *n*-hexane), 5-[4-(*N,N*-dimethylamino)phenyl]-10-(1-ethylpropyl)-20-hexylporphyrin **67** (42 mg, 0.07 mmol) and oxidation with air overnight. Column chromatography on silica gel (CH_2Cl_2 with 1 % NEt_3) gave only the product after recrystallization from $\text{CH}_2\text{Cl}_2/\text{MeOH}$ as purple crystals (8.3 mg, 0.01 mmol, 18 %). M.p. 212 °C. $R_f = 0.40$ (CH_2Cl_2 /*n*-hexane, 2:1, v/v). ^1H NMR (500 MHz, CDCl_3 , TMS): $\delta = 9.67$ (*m*, 1H, β -pyrrole-*H*), 9.57 (*AB*, 2H, $^3J = 4.8$ Hz, β -pyrrole-*H*), 9.53 (*AB*, 4H, $^3J = 4.8$ Hz, β -pyrrole-*H*), 9.40 (*AB*, 1H, $^3J = 4.7$ Hz, β -pyrrole-*H*), 8.89 (*m*, 2H, β -pyrrole-*H*), 8.01 (*AB*, 2H, $^3J = 8.6$ Hz, *Ar-H*), 7.11 (*AB*, 2H, $^3J = 8.6$ Hz, *Ar-H*), 5.02 (*m*, 3H, $\text{CH}(\text{CH}_2)_2$, $\text{CH}_2\text{C}_3\text{H}_7$), 4.93 (*t*, 2H, $^3J = 8.0$ Hz, $\text{CH}_2\text{C}_3\text{H}_7$), 3.23 (*s*, 6H, $\text{N}(\text{CH}_3)_2$), 2.86 (*m*, 4H, $\text{CH}(\text{CH}_2)_2$), 2.58 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{C}_2\text{H}_5$), 2.52 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{C}_4\text{H}_9$), 1.90 (*m*, 2H, $\text{C}_2\text{H}_4\text{CH}_2\text{CH}_3$), 1.82 (*m*, 2H, $\text{C}_2\text{H}_4\text{CH}_2\text{C}_3\text{H}_7$), 1.52 (*m*, 2H, $\text{C}_3\text{H}_6\text{CH}_2\text{C}_2\text{H}_5$), 1.41 (*m*, 2H, $\text{C}_4\text{H}_8\text{CH}_2\text{CH}_3$), 1.19 (*t*, 3H, $^3J = 7.4$ Hz, $\text{C}_3\text{H}_6\text{CH}_3$), 0.95 (*t*, 6H, $^3J = 7.4$ Hz, CH_2CH_3), 0.94 (*t*, 3H, $^3J = 7.2$ Hz, $\text{C}_3\text{H}_{10}\text{CH}_3$), -2.64 ppm (*br s*, 2H, *NH*). ^{13}C NMR (126 MHz, CDCl_3): $\delta = 150.22$, ~ 145 , 135.39, ~ 131.9 , 130.66, ~ 129.1 , ~ 128.8 , ~ 128.4 , ~ 127.8 , 122.10, 119.08, 110.62, 50.26, 41.20, 40.63, 38.77, 35.80, 34.54, 32.02, 30.31, 23.83, 22.83, 13.96 ppm. UV/vis (CH_2Cl_2): λ_{max} (log ϵ) = 420 (4.81), 522 (3.98), 559 (3.88), 600 (3.60), 658 nm (3.72). MS TOF, MS, ES^+ , 70 eV: $m/z = 640$ (10%, $[\text{M}]^+$), 320 (100%, $[\text{M}]^{2+}$). HRMS calcd. for $\text{C}_{43}\text{H}_{54}\text{N}_5$ 640.7379; found 640.4355.

5-Butyl-10-hexyl-15-(3-methoxyphenyl)-20-(4-ethynylphenyl)porphyrin (122):

The synthesis followed the standard procedures using 5-butyl-10-hexyl-15-(3-methoxyphenyl)porphyrin **68** (60 mg, 0.107 mmol), 4-bromoethynylphenyl (292.47 mg, 1.615 mmol), *n*-BuLi (1.3 mL, 3.231 mmol), H_2O (0.5 mL) and DDQ (288.2 mg, 1.26 mmol). Column chromatography on silica gel (*n*-hexane/ CH_2Cl_2 = 4:1, v/v) followed by a second column using *n*-hexane/ CH_2Cl_2 (1:1) gave the title compound as purple crystals (7 mg 0.01 mmol, 10 %). M.p. >300 °C. $R_f = 0.66$ (CH_2Cl_2 /*n*-hexane, 1:1, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): $\delta = -2.69$ (*s*, 2H, *NH*), 0.96 (*t*, 3H, $J = 7.01$ Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.17 (*t*, 3H, $J = 7.01$ Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.43 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.53 (*m*, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.86 (*m*, 4H, CH_2), 2.57 (*m*, 4H, CH_2), 3.34 (*s*, 1H, *CH*), 4.00 (*s*, 3H, OCH_3), 5.02 (*m*, 4H, CH_2), 7.35 (*m*, 1H, *Ar-H*), 7.65 (*t*, 1H, $J = 7.6$ Hz, *Ar-H*), 7.78 (*m*, 2H, *Ar-H*), 7.90 (*d*, 2H, $J = 8.18$ Hz, *Ar-H*), 8.17 (*d*, 2H, $J = 8.18$ Hz, *Ar-H*), 8.73 (*d*, 1H, $J = 4.68$ Hz, β -pyrrole-*H*), 8.81 (*d*, 1H, $J = 4.67$ Hz, β -pyrrole-*H*), 8.85 (*d*, 1H, $J = 4.67$ Hz, β -pyrrole-*H*), 8.93 (*d*, 1H, $J = 4.68$ Hz, β -pyrrole-*H*), 9.48 (*m*, 2H, β -pyrrole-*H*), 9.61 ppm (*s*, 2H, β -pyrrole-*H*). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 13.7$, 22.6, 23.2, 28.9, 30.5, 31.4, 34.9, 35.2, 38.4, 40.5, 52.9, 55.0, 67.5, 83.3, 112.9, 117.0, 118.0, 119.8, 119.9, 120.9, 127.0, 130.0, 133.9, 142.6, 143.1, 157.4 ppm. UV/vis (CH_2Cl_2): λ_{max} (log ϵ) = 418 (5.03), 518 (3.57), 553 (3.29), 592 (3.01), 644 nm (3.04). HRMS (ES^+) calcd. for $\text{C}_{45}\text{H}_{45}\text{N}_4\text{O}$ $[\text{M} + \text{H}]^+$ 657.3593; found 657.357.

Crystal structure determinations: Growth and handling of crystals followed the concept developed by Hope.^[41] Intensity data were collected at 108 K with a Rigaku Saturn-724 system complete with CCD detector utilizing Mo- $\text{K}\alpha$ radiation ($\lambda = 0.71073$ Å). The intensities were corrected for Lorentz, polarization and extinction effects. The structures were solved with Direct Methods using the SHELXTL PLUS program system^[42a] and refined against $[\text{F}^2]$ with the program XL from SHELX-97 using all data.^[E42b] Nonhydrogen atoms were refined with anisotropic thermal parameters. Hydrogen atoms were generally placed into geometrically calculated positions and refined using a riding model. The N-H hydrogen atoms were located in difference maps and refined using the standard riding model. Crystal data for **77**: $\text{C}_{35}\text{H}_{34}\text{N}_4\text{O}_3\text{Br}_2$, $M = 718.46$, triclinic, space group *P*-1, $a = 8.0071(14)$, $b = 11.7648(18)$, $c = 17.446(3)$ Å, $\alpha = 98.114(11)^\circ$, $\beta = 99.002(9)^\circ$, $\gamma = 104.531(10)^\circ$, $V = 1543.1(5)$ Å³, $Z = 2$, $T = 108\text{K}$, $\mu(\text{Mo-K}\alpha) = 2.670$ cm⁻¹, 11705 reflections measured, 5398 unique reflections measured ($R_{\text{int}} = 0.0722$), 401 parameters, 4486 reflections with $I > 2.0\sigma(I)$, refinement against $[\text{F}^2]$, $R1(I > 2.0\sigma(I)) = 0.0339$, $wR2$ (all data) = 0.0859, $S = 0.957$, $\rho_{\text{max}} = 0.514$. Crystal data for **81**: $\text{C}_{35}\text{H}_{34}\text{N}_4\text{O}_3\text{Br}_2$, $M = 718.46$, triclinic, space group *P*-1, $a = 7.9861(11)$, $b = 11.6530(15)$, $c = 17.173(2)$ Å, $\alpha = 90.5430(10)^\circ$, $\beta = 98.012(2)^\circ$, $\gamma = 107.818(2)^\circ$, $V = 1504.3(3)$ Å³, $Z = 2$, $T = 108\text{K}$, $\mu(\text{Mo-K}\alpha) = 2.739$ cm⁻¹, 32735 reflections measured, 8937 unique reflections measured ($R_{\text{int}} = 0.0217$), 401 parameters, 8070 reflections with $I > 2.0\sigma(I)$, refinement against $[\text{F}^2]$, $R1(I > 2.0\sigma(I)) = 0.0338$, $wR2$ (all data) = 0.0823, $S = 0.1046$, $\rho_{\text{max}} = 0.612$.

Supporting Information CCDC-749453 and 749454 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif. Supporting information for this article is available on the WWW under <http://www.eurjoc.org/> or from the author. It contains the synthetic procedures and analytical data for

compounds **20–22, 25–28, 30, 31, 40, 43–50, 52–55, 57, 62, 64, 66–70, 74–81, 83–86, 100, 106, 108, 111, 113, 117–119, and 123–135.**

Acknowledgments

This work was supported by grants from Science Foundation Ireland (SFI Research Professorship 04/RP1/B482) and the Health Research Board (HRB Translational Research Award 2007 TRA/2007/11) and a SFI Ureka award to Y.S. We are grateful to the Centre for Synthesis and Chemical Biology (CSCB) for HRMS measurements. We are indebted to Dr. Tom McCabe for collecting the data sets of some of the crystal structures.


- [1] K. M. Kadish, K. M. Smith, R. Guilard (Eds.) *The Porphyrin Handbook*, Academic Press, New York, **2000, 2003**.
- [2] K. M. Smith (ed.) *Porphyrins and Metalloporphyrins*, Elsevier, Amsterdam, **1975**; L. R. Milgrom *The Colours of Life*, Oxford University Press, Oxford, **1997**.
- [3] J. S. Lindsey in *The Porphyrin Handbook* (Eds. K. M. Kadish, K. M. Smith, R. Guilard), Academic Press, San Diego, **2000**, Vol. 1, pp. 45–118.
- [4] R. A. Sheldon (ed.) *Metalloporphyrins in Catalytic Oxidations*, CRC Press, Boca Raton, **1994**; E. S. Nyman, P. H. Hynninen, *J. Photochem. Photobiol. B: Biol.* **2004**, *73*, 1–28; M. O. Senge, M. Fazekas, E. G. A. Notaras, W. J. Blau, M. Zawadzka, O. B. Locos, E. M. N. Mhuirheartaigh, *Adv. Mater.* **2007**, *19*, 2737–2774.
- [5] a) K. M. Smith in *The Porphyrin Handbook* (K. M. Kadish, K. M. Smith, R. Guilard, Eds.), Academic Press, San Diego, **2000**, Vol. 1, pp. 1–44; b) L. Jaquinod in *The Porphyrin Handbook* (K. M. Kadish, K. M. Smith, R. Guilard, Eds.), Academic Press, San Diego, **2000**, Vol. 1, pp. 201–237.
- [6] a) D. M. Wallace, S. H. Leung, M. O. Senge, K. M. Smith, *J. Org. Chem.* **1993**, *58*, 7245–7257; b) P. D. Rao, S. Dhanalekshmi, B. J. Littler, J. S. Lindsey, *J. Org. Chem.* **2000**, *65*, 7323–7344.
- [7] a) S. H. H. Zaidi, F. M. Fico FM, J. S. Lindsey, *Org. Proc. Res. Devel.* **2006**, *10*, 118–134; b) D. K. Dogutan, S. H. H. Zaidi, P. Thamvongkit, J. S. Lindsey, *J. Org. Chem.* **2007**, *72*, 7701–7714; c) D. K. Dogutan, J. S. Lindsey, *J. Org. Chem.* **2008**, *73*, 6728–6742; d) D. K. Dogutan, M. Ptaszek, J. S. Lindsey, *J. Org. Chem.* **2008**, *73*, 6187–6201.
- [8] M. G. H. Vicente in *The Porphyrin Handbook* (K. M. Kadish, K. M. Smith, R. Guilard, Eds.), Academic Press: San Diego, **2000**; Vol. 1, 149–199; S. Shanmugathasan, C. Edwards, R. W. Boyle, *Tetrahedron* **2000**, *56*, 1025–1046; A. K. Burrell, D. L. Officer, P. G. Plieger, D. C. W. Reid, *Chem. Rev.* **2001**, *101*, 2751–2796; M. O. Senge, J. Richter, *J. Porphyrins Phthalocyanines* **2004**, *8*, 934–953; S. Horn, K. Dahms, M. O. Senge, *J. Porphyrins Phthalocyanines* **2008**, *12*, 1053–1077; H. Shinokubo, A. Osuka, *Chem. Commun.* **2009**, 1011–1021.
- [9] a) J. S. Lindsey, H. C. Hsu, I. C. Schreiman, *Tetrahedron Lett.* **1986**, *27*, 4969–4970; b) J. S. Manka, D. S. Lawence, *Tetrahedron Lett.* **1989**, *30*, 6989–6992.
- [10] a) M. Taniguchi, M. N. Kim, D. Ra, J. S. Lindsey, *J. Org. Chem.* **2005**, *70*, 275–285; b) O. B. Locos, D. P. Arnold, *Org. Biomol. Chem.* **2006**, *1–10*; c) D. M. Shen, C. Liu, Q. Y. Chen, *Synlett*, **2007**, 3068–3072; d) T. Takanami, M. Yotsukura, W. Inoue, F. Hino, K. Suda, *Heterocycles* **2008**, *76*, 439–453; and references cited therein.
- [11] A. Wiehe, Y. M. Shaker, J. C. Brandt, S. Mebs, M. O. Senge, *Tetrahedron* **2005**, *61*, 5535–5564.
- [12] Y. M. Shaker, M. O. Senge, *Heterocycles* **2005**, *65*, 2441–2450.
- [13] a) W. W. Kalisch, M. O. Senge, *Angew. Chem. Int. Ed. Engl.* **1998**, *37*, 1107–1109; b) M. O. Senge, X. Feng, *J. Chem. Soc., Perkin Trans. 1*, **2000**, 3615–3621; c) M. O. Senge, W. W. Kalisch, I. Bischoff, *Chem. Eur. J.* **2000**, *6*, 2721–2738; d) M. O. Senge, *Acc. Chem. Res.* **2005**, *38*, 733–743.
- [14] X. Feng, M. O. Senge, *Tetrahedron* **2000**, *56*, 587–590.
- [15] X. Feng, M. O. Senge, *J. Chem. Soc., Perkin Trans. 1*, **2001**, 1030–1038.
- [16] X. Feng, I. Bischoff, M. O. Senge, *J. Org. Chem.* **2001**, *66*, 8693–8700.
- [17] S. Neya, N. Funasaki, *Tetrahedron Lett.* **2002**, *43*, 1057–1058.
- [18] a) A. Wiehe, C. Ryppa, M. O. Senge, *Org. Lett.* **2002**, *4*, 3807–3809; b) C. Ryppa, M. O. Senge, S. S. Hatscher, E. Kleinpeter, P. Wacker, U. Schilde, A. Wiehe, *Chem. Eur. J.* **2005**, *11*, 3427–3442.
- [19] a) M. B. Bakar, M. Oelgemöller, M. O. Senge, M. O. *Tetrahedron* **2009**, *65*, 7064–7078; b) M. Zawadzka, J. Wang, W. J. Blau, M. O. Senge, *Chem. Phys. Lett.* **2009**, *477*, 330–335.
- [20] M. J. Gunter, L. N. Mander, *J. Org. Chem.* **1981**, *46*, 4791–4795.
- [21] S. Hatscher, M. O. Senge, *Tetrahedron Lett.* **2003**, *44*, 157–160.
- [22] R. P. Brinas, C. Brückner, *Tetrahedron* **2002**, *58*, 4375–4381.
- [23] a) Z. Yao, J. Bhaumik, S. Dhanalekshmi, M. Ptaszek, P. A. Rodriguez, J. S. Lindsey, *Tetrahedron* **2007**, *63*, 10657–10670; b) A. Z. Muresan, J. S. Lindsey, *Tetrahedron* **2008**, *64*, 11440–11448.
- [24] Recently we reported on a mixed synthesis of the various regioisomeric mono-bromo AB-porphyrins which may be used for such purposes: M. Fazekas, M. Pintea, M. O. Senge, M. Zawadzka, *Tetrahedron Lett.* **2008**, *49*, 2236–2239.
- [25] G. Simonneaux, P. Tagliatesta, *J. Porphyrins Phthalocyanines* **2004**, *8*, 1166–1171. E. Rose, B. Andrioletti, S. Zrig, M. Quelquejeu-Ehteve, *Chem. Soc. Rev.* **2005**, *34*, 573–583.
- [26] A. Nakano, H. Shimidzu, A. Osuka, *Tetrahedron Lett.* **1998**, *39*, 9489–9492. S. G. DiMaggio, V. S. Y. Lin, M. J. Therien, *J. Org. Chem.* **1993**, *58*, 5983–5993.
- [27] a) A. Suzuki, *A. J. Organomet. Chem.* **1999**, *576*, 147–168; b) B. Shi, R. W. Boyle, *J. Chem. Soc., Perkin Trans. 1* **2002**, 1397–1400.
- [28] S. Banfi, E. Caruso, E. Fieni, L. Buccafurni, M. B. Gariboldi, R. Rivazza, E. Monti, *J. Porphyrins Phthalocyanines* **2006**, *10*, 1319–1326.
- [29] a) A. Osuka, M. Ikeda, H. Shiratori, Y. Nishimura, I. Yamazaki, *J. Chem. Soc., Perkin Trans. 2* **1999**, 1019–1026; b) J. -P. Tremblay-Morin, H. Ali, J. E. Van Lier, *Tetrahedron Lett.* **2006**, *47*, 3043–3046; c) K. Sugiura, Y. Fujimoto, Y. Sakata, *Chem. Commun.* **2000**, 1105–1106.
- [30] a) J. S. Lindsey, S. Prathapan, T. E. Johnson, R. W. Wagner, *Tetrahedron* **1994**, *50*, 8941–8968; b) R. W. Wagner, T. E. Johnson, F. Li, J. S. Lindsey, *J. Org. Chem.* **1995**, *60*, 5266–5273; c) R. W. Wagner, T. E. Johnson, J. S. Lindsey, *J. Am. Chem. Soc.* **1996**, *118*, 11166–11180.
- [31] The compound was too unstable for ¹³C NMR, CHN, and MS analysis.
- [32] a) B. Krattinger, H. G. Callot, *Tetrahedron Lett.* **1996**, *37*, 7699–7702; b) B. Krattinger, H. G. Callot, *Tetrahedron Lett.* **1998**, *39*, 1165–1168.
- [33] N. N. Sergeeva, Y. M. Shaker, E. M. Finnigan, T. McCabe, M. O. Senge, *Tetrahedron* **2007**, *63*, 12454–12464.
- [34] M. O. Senge, I. Bischoff, *Eur. J. Org. Chem.* **2001**, 1735–1751.
- [35] P. Bhyrappa, B. Purushothaman, J. Vittal, *J. Porphyrins Phthalocyanines* **2003**, *7*, 682–692.
- [36] a) W. R. Scheidt, Y. J. Lee, *Struct. Bonding (Berlin)* **1987**, *64*, 1–70; b) M. O. Senge, C. W. Eigenbrot, T. D. Brennan, J. Shusta, W. W. Scheidt, K. M. Smith, *Inorg. Chem.* **1993**, *32*, 3134–3142.
- [37] a) C. J. Medforth, M. O. Senge, T. P. Forsyth, J. D. Hobbs, J. A. Shelnutt, K. M. Smith, *Inorg. Chem.* **1994**, *33*, 3865–3872; b) M. O. Senge, T. P. Forsyth, K. M. Smith, *Z. Kristallogr.* **1996**, *211*, 176–185.
- [38] M. O. Senge, *Chem. Commun.* **2006**, 243–256.
- [39] M. S. Newman, L. F. Lee, *J. Org. Chem.* **1972**, *37*, 4468–4469.
- [40] R. Beavington, P. L. Burn, *J. Chem. Soc., Perkin Trans. 1*, **1999**, 583–592.
- [41] H. Hope, *Prog. Inorg. Chem.* **1994**, *41*, 1–19.

- [42] a) G. M. Sheldrick, SHELXS-93, Program for the Solution of Crystal Structures, Universität Göttingen, Göttingen (Germany), **1993**; b) G. M. Sheldrick, SHELXL-97, Program for the Refinement of Crystal Structures, Universität Göttingen, Göttingen (Germany), **1997**.

Received: ((will be filled in by the editorial staff))
Published online: ((will be filled in by the editorial staff))

Entry for the Table of Contents ((Please choose one layout.))

A comprehensive study of contemporary synthetic methods using organolithium and Pd-catalyzed C–C coupling reactions for ABCD porphyrins reveals that it is now possible to prepare almost any desired meso substituted porphyrin.



((Porphyrins))

Mathias O. Senge,* Yasser M. Shaker, Monica Pinteá, Claudia Ryppa, Sabine S. Hatscher, Aoife Ryan and Yulia Sergeeva Page No. – Page No.

Synthesis of meso-Substituted ABCD-type Porphyrins via Functionalization Reactions

Keywords: Porphyrinoids / C–C coupling / Organolithium / Tetrapyrroles / Conformation analysis

SUPPORTING INFORMATION

**Synthesis of meso-Substituted
ABCD-type Porphyrins via
Functionalization Reactions**

**Mathias O. Senge,^{*,[a,b]} Yasser M.
Shaker,^[a] Monica Pinteá,^[a] Claudia
Ryppa,^[c] Sabine S. Hatscher,^[c] Aoife
Ryan^[a] and Yulia Sergeeva^[a]**

[a] School of Chemistry, SFI Tetrapyrrole
Laboratory, Trinity College Dublin, Dublin 2,
Ireland

Fax: +353-1-896-8536

E-mail: sengem@tcd.ie

[b] Medicinal Chemistry, Institute of Molecular
Medicine, Trinity Centre for Health Sciences,
Trinity College Dublin, St James's Hospital,
Dublin 8, Ireland

[c] Institut für Chemie, Universität Potsdam,
Karl-Liebknecht-Str. 24-25, D-14476 Golm,
Germany

Ref. No.:

General Procedure for the synthesis of AB -type porphyrins (mixed condensation)

A three-necked-flask equipped with magnetic stirrer, gas inlet (argon) and a reflux condenser was charged with 2 L dichloromethane. Dipyrromethane (1200 mg, 8.2 mmol), the appropriate aldehyde (4.14 mmol), and the other appropriate aldehyde (4.15 mmol) were added. Then 140 μ L (1.8 mmol) TFA were added and the reaction mixture was stirred for 16 h at 20 °C. Next DDQ (2.77 g, 12.2 mmol) in 100 mL of dry dichloromethane was added and the mixture was stirred for 1 h. Subsequently, 6 mL of NEt₃ were added and the reaction mixture was filtered through 600 mL of silica, eluting with dichloromethane. Column chromatography eluting with dichloromethane gave the fractions which were recrystallized from CH₂Cl₂/MeOH to yield purple crystals.

5-(2-Methoxyphenyl)-15-(4-methylphenyl)porphyrin

(20): The procedure followed that given in the general procedure above using 435 μ L 2-methoxybenzaldehyde (3.6 mmol) and 422 μ L tolylaldehyde (3.6 mmol). Column chromatography on silica used elution with dichloromethane/*n*-hexane (1:3, v/v). The second fraction gave 400 mg of the title compound (0.8 mmol, 22 %) after recrystallization. M.p. > 320 °C. ¹H NMR (500 MHz, CDCl₃, TMS): δ = 10.25 (s, 2H, meso-*H*), 9.35 (2H, AB, ³*J* = 4.54 Hz, β -pyrrole-*H*_{12,18}), 9.34 (AB, 2H, ³*J* = 4.54 Hz, β -pyrrole-*H*_{2,8}), 9.09 (AB, 2H, ³*J* = 4.54 Hz, β -pyrrole-*H*_{13,17}), 8.99 (AB, ³*J* = 4.54 Hz, β -pyrrole-*H*_{3,7}), 8.16 (*m*, 2H, tolyl-*H*_o), 8.05 (*dd*, 1H, ³*J* = 7.54 Hz, ⁴*J* = 1.90 Hz, MeOC₆H₄-*H*_o), 7.80 (*dd*, 1H, ³*J* = 8.00 Hz, ⁴*J* = 1.90 Hz, MeOC₆H₄-*H*_p), 7.61 (AB, 2H, H₂₉, H₃₁, ³*J* = 7.54 Hz, tolyl-*H*_m), 7.40 (*td*, 1H, ³*J* = 7.40 Hz, ⁴*J* = 1.00 Hz, MeOC₆H₄-*H*_m), 7.37 (*d*, 1H, ³*J* = 8.45 Hz, MeOC₆H₄-*H*_m), 3.61 (s, 3H, OCH₃), 2.73 (s, 3H, C₆H₄-CH₃), -3.06 ppm (*bs*, 2H, NH). ¹³C NMR (126 MHz, CDCl₃): δ = 159.48, 147.60, 147.07, 145.17, 138.56, 137.35, 135.87, 134.78, 131.55, 131.25, 131.25, 130.98, 130.76, 130.23, 129.94, 127.68, 119.66, 119.35, 114.86, 111.09, 104.96, 55.79, 21.50 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ϵ) = 403 (5.59), 502 (4.18), 536 (3.71), 575 (3.72), 628 nm (3.21). MS (FAB⁺, 3 kV), *m/z* = 507.0 (16.75 %, [M + H]⁺). HRMS: calcd. for C₃₄H₂₆N₄O 506.2107; found 506.2107.

5-(4-Methylphenyl)-15-(2,4,6-trimethoxyphenyl)porphyrin

(21): The procedure followed that of the general one given above. Dipyrromethane (1 g, 6.86 mmol), 422 μ L tolylaldehyde (3.6 mmol) and 2,4,6-trimethoxybenzaldehyde (700 mg, 3.6 mmol) were dissolved in a 3-necked flask charged with 1.7 L dichloromethane and purged with argon. After addition of 110 μ L TFA the mixture was stirred for 14 h and then concentrated to about 1/10 of the volume. Column chromatography started with dichloromethane followed by addition of increasing parts of *n*-hexane and gave three fractions. The second contained the title compound and gave purple crystals after recrystallization from CH₂Cl₂/MeOH (330 mg, 0.59 mmol, 15 %). M.p. > 320 °C. ¹H NMR (500 MHz, CDCl₃, TMS): δ = 10.22 (s, 2H, meso-*H*), 9.34 (*m*, 4H, β -pyrrole-*H*), 9.08 (*d*, 2H, ³*J* = 4.54 Hz, β -pyrrole-*H*), 9.022 (*d*, 2H, ³*J* = 4.54 Hz, β -pyrrole-*H*), 8.16 (*d*, 2H, ³*J* = 7.81 Hz, tolyl-*H*_o), 7.61 (*d*, 2H, ³*J* = 4.5 Hz, tolyl-*H*_m), 6.63 (s, 2H, Ar-*H*_m), 4.12 (s, 3H, *p*-OCH₃), 3.53 (s, 6H, *m*-OCH₃), 2.71 (s, 3H, C₆H₄-CH₃), -3.05 ppm (*bs*, 2H, NH). ¹³C NMR

(126 MHz, CDCl₃): δ = 161.99, 161.12, 148.41, 146.70, 145.14, 138.62, 137.23, 134.75, 131.57, 131.02, 130.76, 130.50, 127.63, 119.12, 111.60, 110.89, 104.62, 90.89, 55.95, 55.64, 21.50 ppm. UV/vis (CH₂Cl₂): λ_{max} (log ϵ) = 406 (5.17), 503 (4.06), 538 (3.59), 576 (3.58), 628 nm (3.05). MS (300 °C, 80 eV), *m/z* = 566.0 (100.00 %, [M]⁺), 535.0 (19 %, [M - CH₃O]⁺), 283.0 (22.4 %, [M]²⁺). HRMS calcd. for C₃₆H₃₀N₄O₃ 566.2318; found 566.2318. C₃₆H₃₀N₄O₃ (566.66): calcd. C 76.31, H 5.34, N 9.89; found C 76.52, H 5.51, N 9.63.

5-(*n*-Butyl)-15-(2,4,6-trimethoxyphenyl)porphyrin

(22): The procedure followed that given above using dipyrromethane (1.2 g, 8.2 mmol), 2,4,6-trimethoxybenzaldehyde (805 mg, 4.1 mmol), pentanal (0.44 mL, 4.1 mmol), TFA (0.11 mL, 1.5 mmol) and DDQ (4.63 g, 20 mmol). Column chromatography on silica used elution with *n*-hexane/ethyl acetate (100:1, v/v) gave 5,15-dibutylporphyrin **23** (143 mg, 0.33 mmol, 16 %) as purple crystals from the first fraction, followed by **22** (300 mg, 0.56 mmol, 14 %) as purple crystals and 5,15-bis(2,4,6-trimethoxyphenyl)porphyrin **24** (150 mg, 0.28 mmol, 11 %) as purple crystals. M.p. > 300 °C. *R_f* = 0.5 (*n*-hexane : CH₂Cl₂ = 1 : 1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.92 (s, 2H, NH), 0.96 (*t*, 3H, *J* = 7.02 Hz, CH₂CH₂CH₂CH₃), 1.87 (*m*, 2H, CH₂CH₂CH₂CH₃), 2.56 (*m*, 2H, CH₂CH₂CH₂CH₃), 3.55 (s, 6H, *o*-OCH₃), 4.15 (s, 3H, *p*-OCH₃), 5.08 (*t*, 2H, *J* = 7.6 Hz, CH₂CH₂CH₂CH₃), 6.64 (s, 2H, Ar-*H*_m), 8.99 (*d*, 2H, *J* = 4.68 Hz, β -pyrrole-*H*), 9.31 (*d*, 2H, *J* = 4.68 Hz, β -pyrrole-*H*), 9.42 (*d*, 2H, *J* = 4.68 Hz, β -pyrrole-*H*), 9.62 (*d*, 2H, *J* = 4.09 Hz, β -pyrrole-*H*), 10.18 ppm (s, 2H, meso-*H*). ¹³C NMR (126 MHz, CDCl₃): δ = 13.3, 13.8, 18.7, 23.2, 24.5, 30.2, 33.9, 34.1, 40.3, 55.2, 55.5, 63.7, 90.5, 103.7, 109.7, 111.2, 119.0, 127.2, 129.8, 130.9, 131.1, 143.9, 144.3, 146.2, 148.2, 160.7, 161.5, 173.5 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ϵ) = 407 (4.93), 503 (3.90), 536(3.42), 577(3.57), 633 nm (3.32). HRMS: calcd. for C₃₃H₃₃N₄O₃ 533.2553; found 533.2534.

5-(Pentyl)-15-(2,4,6-trimethoxyphenyl)porphyrin

(25): The procedure followed that given above using dipyrromethane (1.2 g, 8.2 mmol), 2,4,6-trimethoxybenzaldehyde (805 mg, 4.1 mmol), hexanal (0.49 mL, 4.1 mmol), TFA (0.11 mL, 1.5 mmol) and DDQ (4.63 g, 20 mmol). Column chromatography on silica used elution with *n*-hexane/ethyl acetate (100:1, v/v) gave 5,15-dipentylporphyrin **26** (50 mg, 0.11 mmol, 5.4 %) as purple crystals from the first fraction, followed by **25** (200 mg, 0.36 mmol, 9 %) as purple crystals and 5,15-bis(2,4,6-trimethoxyphenyl)porphyrin **24** (30 mg, 0.046 mmol, 2.2 %) as purple crystals. M.p. > 300 °C. *R_f* = 0.4 (*n*-hexane : CH₂Cl₂ = 1:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.89 (s, 2H, NH), 1.04 (*t*, 3H, *J* = 6.85 Hz, CH₂CH₂CH₂CH₂CH₃), 1.62 (*m*, 2H, CH₂CH₂CH₂CH₂CH₃), 1.84 (*m*, 2H, CH₂CH₂CH₂CH₂CH₃), 2.61 (*m*, 2H, CH₂CH₂CH₂CH₂CH₃), 3.58 (s, 6H, *o*-OCH₃), 4.15 (s, 3H, *p*-OCH₃), 5.04 (*t*, 2H, *J* = 7.8 Hz, CH₂CH₂CH₂CH₂CH₃), 6.68 (s, 2H, Ar-*H*_m), 9.05 (*d*, 2H, *J* = 3.92 Hz, β -pyrrole-*H*), 9.35 (*d*, 2H, *J* = 4.89 Hz, β -pyrrole-*H*), 9.41 (*d*, 2H, *J* = 3.91 Hz, β -pyrrole-*H*), 9.60 (*d*, 2H, *J* = 3.91 Hz, β -pyrrole-*H*), 10.19 ppm (s, 1H, meso-*H*). ¹³C NMR (126 MHz, CDCl₃): δ = 14.1, 22.8, 30.9, 32.7, 34.8, 38.3, 55.6, 55.9, 91.0, 104.2, 104.6, 110.2, 111.6, 119.5, 127.7, 130.3, 130.4, 131.3,

131.6, 144.3, 144.8, 146.7, 148.6, 161.1, 162.0 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ϵ) = 407 (4.99), 504 (3.73), 538 (3.58), 579 (3.64), 629 nm (3.43). HRMS (ES⁺) calcd. for C₃₄H₃₄N₄O₃ (M+H⁺) 547.2709; found 547.2719.

5-(Hexyl)-15-(2,4,6-trimethoxyphenyl)porphyrin (27):

The procedure followed that given above using dipyrromethane (1.2 g, 8.2 mmol), 2,4,6-trimethoxybenzaldehyde (804.4 mg 4.1 mmol), heptanal (0.57 mL, 4.1 mmol), TFA (0.11 mL, 1.5 mmol) and DDQ (4.63 g, 20 mmol). Column chromatography on silica used elution with CH₂Cl₂ : *n*-hexane (1:1, v/v) and gave 5,15-dihexylporphyrin **18** (120 mg, 0.25 mmol, 12%) as purple crystals from the first fraction, followed by **27** (400 mg, 0.71 mmol, 17%) as purple crystals and 5,15-bis(2,4,6-trimethoxyphenyl)porphyrin **24** (150 mg, 0.23 mmol, 9%) as purple crystals. M.p. > 300 °C. *R_f* = 0.4 (*n*-hexane : CH₂Cl₂ = 1:2, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.90 (s, 2H, NH), 0.99 (t, 3H, *J* = 7.01, Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 1.45 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.57 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.86 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.60 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 3.57 (s, 6H, *o*-OCH₃), 4.15 (s, 3H, *p*-OCH₃), 5.06 (t, 2H, *J* = 8.19 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 6.67 (s, 2H, Ar-*H_m*), 9.03 (d, 2H, *J* = 4.67 Hz, β -pyrrole-*H*), 9.34 (d, 2H, *J* = 4.68 Hz, β -pyrrole-*H*), 9.42 (d, 2H, *J* = 4.68 Hz, β -pyrrole-*H*), 9.61 (d, 2H, *J* = 4.68 Hz, β -pyrrole-*H*), 10.19 ppm (s, 1H, meso-*H*). ¹³C NMR (126 MHz, CDCl₃): δ = 13.7, 22.3, 29.2, 29.8, 31.5, 34.4, 38.2, 55.2, 55.5, 90.5, 103.7, 109.7, 112.2, 119.1, 127.1, 129.8, 130.9, 131.1, 143.9, 144.3, 146.2, 148.2, 160.7, 161.5 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ϵ) = 407 (5.07), 504 (4.04), 536 (3.74), 576 (3.82), 630 nm (3.74). HRMS (ES⁺) calcd. for C₃₅H₃₆N₄O₃ (M+H⁺) 561.2866; found 561.2847.

5-(*n*-Butyl)-15-(3,4,5-trimethoxyphenyl)porphyrin (28):

The procedure followed that given above using dipyrromethane (1.2 g, 8.2 mmol), 3,4,5-trimethoxybenzaldehyde (805 mg 4.6 mmol), pentanal (0.61 mL, 4.6 mmol), TFA (0.11 mL, 1.5 mmol) and DDQ (4.63 g, 20 mmol). Column chromatography on silica used elution with CH₂Cl₂/*n*-hexane (2:1, v/v) gave 5,15-dibutylporphyrin **23** (60 mg, 0.14 mmol, 6%) as purple crystals from the first fraction, followed by **28** (175 mg, 0.32 mmol, 8%) as purple crystals and 5,15-bis(3,4,5-trimethoxyphenyl)porphyrin **29** (30 mg, 0.046 mmol, 2.2 %) as purple crystals. M.p. > 300 °C. *R_f* = 0.5 (CH₂Cl₂). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.98 (s, 2H, NH), 1.19 (t, 3H, *J* = 7.01 Hz, CH₂CH₂CH₂CH₃), 1.90 (m, 2H, CH₂CH₂CH₂CH₃), 2.56 (m, 2H, CH₂CH₂CH₂CH₃), 4.03 (s, 6H, *m*-OCH₃), 4.22 (s, 3H, *p*-OCH₃), 5.10 (t, 2H, *J* = 8.2, Hz, CH₂CH₂CH₂CH₃), 7.52 (s, 2H, Ar-*H_o*), 9.16 (d, 2H, *J* = 4.09 Hz, β -pyrrole-*H*), 9.39 (d, 2H, *J* = 4.09 Hz, β -pyrrole-*H*), 9.48 (d, 2H, *J* = 4.67 Hz, β -pyrrole-*H*), 9.67 (d, 2H, *J* = 4.68 Hz, β -pyrrole-*H*), 10.27 ppm (s, 2H, meso-*H*). ¹³C NMR (126 MHz, CDCl₃): δ = 13.8, 23.4, 24.3, 34.3, 40.7, 56.3, 61.2, 105.0, 113.1, 118.2, 120.1, 128.4, 131.0, 132.0, 132.2, 137.3, 138.1, 145.0, 145.2, 147.7, 152.1 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ϵ) = 409 (4.93), 506 (3.90), 542(3.72), 582 (3.42), 636 ppm (3.32). HRMS (ES): calcd. for C₃₃H₃₂N₄O₃ (M+H⁺) 33.2553; found 533.2562.

5-Pentyl-15-(3,4,5-trimethoxyphenyl)porphyrin (30):

Similar to the procedure given above using dipyrromethane (1 g, 6.84 mmol), 3,4,5-trimethoxybenzaldehyde (671 mg, 3.42 mmol), hexanal (0.42 mL, 3.42 mmol), TFA (0.11 mL, 1.5 mmol) and DDQ (4.63 g, 20 mmol) were used. The reaction mixture was separated by column chromatography using CH₂Cl₂/*n*-hexane (2:1, v/v) as eluent. The first fraction was 5,15-dipentylporphyrin **26** (73 mg, 0.162 mmol, 5 %) as purple crystals, the second fraction 5-pentyl-15-(3,4,5-trimethoxyphenyl)porphyrin **30** (110 mg, 0.2 mmol, 6 %) as purple crystals, and the third fraction **29** (50 mg, 0.38 mmol, 2 %) as purple crystals. M.p. >300 °C. *R_f* = 0.5 (CH₂Cl₂). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.97 (s, 2H, NH), 1.03 (t, 3H, *J* = 7.01 Hz, CH₂CH₂CH₂CH₂CH₃), 1.60 (m, 2H, CH₂CH₂CH₂CH₂CH₃), 1.86 (m, 2H, CH₂CH₂CH₂CH₂CH₃), 2.61 (m, 2H, CH₂CH₂CH₂CH₂CH₃), 4.03 (s, 6H, *m*-OCH₃), 4.23 (s, 3H, *p*-OCH₃), 5.08 (t, 2H, *J* = 7.6 Hz, CH₂CH₂CH₂CH₂CH₃), 7.52 (s, 2H, Ar-*H_o*), 9.16 (d, 2H, *J* = 4.68 Hz, β -pyrrole-*H*), 9.39 (d, 2H, *J* = 4.68 Hz, β -pyrrole-*H*), 9.47 (d, 2H, *J* = 4.68 Hz, β -pyrrole-*H*), 9.65 (d, 2H, *J* = 4.68 Hz, β -pyrrole-*H*), 10.26 ppm (s, 2H, meso-*H*). ¹³C-NMR (150 MHz, CDCl₃, TMS): δ = 13.9, 22.6, 32.6, 34.6, 38.3, 56.2, 61.1, 104.6, 112.8, 117.8, 119.7, 127.9, 130.5, 131.5, 131.7, 136.7, 137.7, 144.4, 144.7, 147.1, 151.1 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ϵ) = 407 (5.15), 505 (4.03), 538 (3.43), 577 (3.58), 634 nm (3.33). HRMS calcd. for C₃₄H₃₄N₄O₃ 547.2709; found 547.2719.

5,15-Dipentylporphyrin (26):

Obtained from the synthesis of **30**. M.p. >300 °C. *R_f* = 0.9 (CH₂Cl₂). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.88 (s, 2H, NH), 1.00 (t, 6H, *J* = 7.02 Hz, CH₂CH₂CH₂CH₂CH₃), 1.58 (m, 4H, CH₂CH₂CH₂CH₂CH₃), 1.81 (m, 4H, CH₂CH₂CH₂CH₂CH₃), 2.58 (m, 4H, CH₂CH₂CH₂CH₂CH₃), 5.02 (t, 2H, *J* = 7.6 Hz, CH₂CH₂CH₂CH₂CH₃), 9.42 (d, 4H, *J* = 4.68 Hz, β -pyrrole-*H*), 9.59 (d, 4H, *J* = 4.68 Hz, β -pyrrole-*H*), 10.18 ppm (s, 2H, meso-*H*). ¹³C-NMR (150 MHz, CDCl₃, TMS): δ = 13.9, 14.0, 22.6, 22.8, 29.5, 32.5, 34.4, 34.5, 36.5, 38.1, 104.0, 104.5, 118.6, 119.1, 127.6, 128.9, 129.0, 131.3, 131.6, 131.7, 144.0, 144.7, 144.3 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ϵ) = 405 (4.90), 505 (3.82), 538(3.35), 587 (3.49), 636 nm (3.25). HRMS calcd. for C₃₀H₃₄N₄ 451.2862; found 451.2862.

5-Hexyl-15-(3,4,5-trimethoxyphenyl)porphyrin (31):

Similar to the procedure given above using dipyrromethane (1.2 g, 8.2 mmol), 3,4,5-trimethoxybenzaldehyde (805 mg, 4.1 mmol), heptanal (0.61 mL, 4.6 mmol), TFA (0.11 mL, 1.5 mmol) and DDQ (4.63 g 2.0 mmol) were used. The reaction mixture was separated by column chromatography using CH₂Cl₂/*n*-hexane (2:1, v/v) as eluent. The first fraction was **18** (120 mg, 0.25 mmol, 6%) as purple crystals, followed by the title compound **31** (370 mg, 0.7 mmol, 17%) as purple crystals, and **29** (200 mg, 0.38 mmol, 9%) as purple crystals. M.p. >300 °C. *R_f* = 0.5 (*n*-hexane:CH₂Cl₂ = 1:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.98 (s, 2H, NH), 0.97 (t, 3H, *J* = 7.1 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 1.27 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.44 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.88 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.63 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 4.03 (s, 6H, *m*-OCH₃), 4.21 (s, 3H, *p*-OCH₃), 5.09 (t, 2H, *J* = 7.6 Hz,

$\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 7.52 (*s*, 2H, Ar- H_o), 9.15 (*d*, 2H, $J = 4.68$ Hz, β -pyrrole- H), 9.39 (*d*, 2H, $J = 4.68$ Hz, β -pyrrole- H), 9.48 (*d*, 2H, $J = 4.09$ Hz, β -pyrrole- H), 9.66 (*d*, 2H, $J = 4.68$ Hz, β -pyrrole- H), 10.27 ppm (*s*, 1H, meso- H). ^{13}C NMR (126 MHz, CDCl_3): $\delta = 14.0, 22.6, 30.1, 31.8, 34.7, 38.6, 56.2, 61.1, 97.6, 104.7, 112.8, 117.8, 119.7, 127.9, 130.5, 131.5, 131.7, 132.1, 136.7, 137.7, 144.4, 144.7, 145.1, 147.1$ ppm. UV/vis (CH_2Cl_2): λ_{max} ($\log \epsilon$) = 407 (5.14), 503 (4.04), 538 (3.44), 577 (3.52), 639 nm (3.22) nm. HRMS calcd. for $\text{C}_{35}\text{H}_{36}\text{N}_4\text{O}_3$ 561.2866;] ($\text{M}+\text{H}^+$): calcd 561.2866, found 561.2847.

General procedure for the synthesis of 5,15-AB-type porphyrins (organolithium reaction)

5-(*tert*-Butyl)porphyrin **32** (100 mg, 0.272 mmol) was dissolved in 80 mL dry THF under argon atmosphere and the reaction mixture was cooled down to -78°C . $n\text{BuLi}$ (0.435 mL, 1.088 mmol) was added dropwise over 15 minutes *via* syringe. The cold bath was removed and stirring continued 1.5 hours at room temperature. 0.5 mL H_2O was added and stirring continued 15 min. Then DDQ (243.3 mg, 1.088 mmol) was added in 10 mL THF and the reaction mixture was stirred for an additional hour. The reaction mixture was filtered through silicagel, followed by evaporation of the solvent

5-*tert*-Butyl-15-hexylporphyrin (40): The synthesis was performed similar to the above general procedure using 5-*tert*-butylporphyrin **32** (50 mg, 0.14 mmol) and a 2.5 M solution of *n*-hexyllithium in hexane (0.19 mL, 0.48 mmol, 3.5 equiv.). The crude product was purified on silica gel (CH_2Cl_2 : *n*-hexane = 1:1, v/v). The first fraction contained a mixture of higher alkylated products. The second fraction yielded the title compound as purple crystals after recrystallization from $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (9 mg, 0.02 mmol, 14 %). M.p. $>320^\circ\text{C}$. $R_f = 0.48$ (CH_2Cl_2 /*n*-hexane = 1:1, v/v). ^1H NMR (500 MHz, CDCl_3 , TMS): $\delta = 10.00$ (*s*, 2H, meso- H), 9.82 (*AB*, 2H, $^3J = ^3J = 4.8$ Hz, β -pyrrole- $H_{3,7}$), 9.45 (*AB*, 2H, $^3J = ^3J = 4.5$ Hz, β -pyrrole- $H_{13,17}$), 9.29 (*AB*, 2H, $^3J = ^3J = 4.5$ Hz, β -pyrrole- $H_{12,18}$), 9.20 (*AB*, 2H, $^3J = ^3J = 4.8$ Hz, β -pyrrole- $H_{2,8}$), 4.88 (*t*, 2H, $^3J = 8.0$ Hz, $\text{CH}_2\text{C}_5\text{H}_{11}$), 2.58 (*s*, 9H, $\text{C}(\text{CH}_3)_3$), 2.50 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{C}_4\text{H}_9$), 1.76 (*m*, 2H, $\text{C}_2\text{H}_4\text{CH}_2\text{C}_3\text{H}_7$), 1.50 (*m*, 2H, $\text{C}_3\text{H}_6\text{CH}_2\text{C}_2\text{H}_5$), 1.38 (*m*, 2H, $\text{C}_4\text{H}_8\text{CH}_2\text{CH}_3$), 0.93 (*t*, 3H, $^3J = 7.3$ Hz, $\text{C}_5\text{H}_{10}\text{CH}_3$), -1.95 ppm (*br s*, 2H, NH). ^{13}C NMR (126 MHz, CDCl_3): $\delta = 148.19, 147.09, 143.50, 141.91, 131.41, 131.13, 130.11, 127.50, 126.39, 119.62, 104.34, 40.80, 40.31, 38.31, 34.10, 31.89, 30.16, 22.71, 14.13$ ppm. UV/vis (CH_2Cl_2): λ_{max} ($\log \epsilon$) = 407 (5.55), 504 (4.37), 537 (3.73), 578 (3.83), 633 nm (3.31). MS (250 $^\circ\text{C}$, 80 eV): $m/z = 450$ (100 %, $[\text{M}]^{*+}$), 435 (72 %, $[\text{M} - \text{CH}_3]^+$), 394 (10 %, $[\text{M} - \text{C}_4\text{H}_9 + \text{H}]^{*+}$), 379 (7 %, $[\text{M} - \text{C}_5\text{H}_{11}]^+$), 323 (17 %, $[\text{M} - \text{C}_4\text{H}_9 + \text{H} - \text{C}_5\text{H}_{11}]^+$), 225 (3 %, $[\text{M}]^{2+}$). HRMS: calcd. for $\text{C}_{30}\text{H}_{34}\text{N}_4$ 450.27835, found 450.27572.

General procedure for the synthesis of 5,10-AB-type porphyrins

A 250 mL Schlenk flask was charged with the porphyrin (1 equiv.) in 50 mL THF under Ar and cooled to -70°C . The LiR reagent (hexyllithium, 3.5 equiv. of a 2.5 M solution in hexane) was added dropwise over 15 min and the color changed from red to green-brown. The cold bath was

removed and the solution stirred for 30 min at rt. Next water (1 mL in 5 mL THF) was added and the color changed to green. The solution was stirred for 15 min and then oxidized with DDQ followed by stirring for a further 30 min accompanied by a color change to red. The mixture was filtered through silica gel eluting with dichloromethane and the solvents removed *in vacuo*.

5-(1-Ethylpropyl)-10-phenylporphyrin (43): Similar to the procedure given above (0°C) 5-(1-ethylpropyl)porphyrin **33** (63 mg, 0.17 mmol) was reacted with a 1.8 M solution of PhLi in ether (0.60 mL, 1.08 mmol, 6.5 equiv.). Column chromatography on silica gel eluting with CH_2Cl_2 /*n*-hexane (1:1, v/v) gave a first fraction of higher alkylated porphyrins followed by a fraction with the title compound to yield purple crystals after recrystallization (56 mg, 0.12 mmol, 69 %). M.p. 166°C . $R_f = 0.67$ (*n*-hexane/ CH_2Cl_2 = 2:1, v/v). ^1H NMR (300 MHz, CDCl_3 , TMS): $\delta = 10.18$ (*s*, 1H, meso- H_{20}), 10.11 (*s*, 1H, meso- H_{15}), 9.74 (*m*, 2H, β -pyrrole- $H_{3,7}$), 9.39 (*m*, 3H, β -pyrrole- $H_{2,17,18}$), 9.28 (*AB*, 1H, $^3J_{12,13} = 4.7$ Hz, β -pyrrole- H_{13}), 8.99 (*m*, 2H, β -pyrrole- $H_{8,12}$), 8.27 (*m*, 2H, Ar- H_o), 7.80 (*m*, 3H, Ar- $H_{m,p}$), 5.17 (*m*, 1H, $\text{CH}(\text{CH}_2)_2$), 3.01, 2.84 (*m*, 4H, $\text{CH}(\text{CH}_2)_2$), 0.98 (*t*, 6H, $^3J = 7.4$ Hz, CH_2CH_3), -3.17 ppm (*br s*, 2H, NH). ^{13}C NMR (75 MHz, CDCl_3): $\delta = 142.8, 134.49, \sim 132, \sim 131, \sim 129, 127.67, 126.50, 123.98, \sim 119, 103.93, 103.32, 50.49, 34.84, 14.18$ ppm. UV/vis (CH_2Cl_2): λ_{max} ($\log \epsilon$) = 406 (5.48), 503 (4.13), 535 (3.31), 577 (3.57), 631 nm (2.94). MS (EI, 70 eV): $m/z = 456$ (52 %, $[\text{M}]^{*+}$), 427 (63 %, $[\text{M} - \text{C}_2\text{H}_5]^+$), 228 (12 %, $[\text{M}]^{2+}$), 214 (17 %, $[\text{M} - \text{C}_2\text{H}_5]^{2+}$). HRMS calcd. for $\text{C}_{31}\text{H}_{28}\text{N}_4$ 456.2314; found 456.2286.

5-[4-(*N,N*-Dimethylamino)phenyl]-15-(1-ethylpropyl)porphyrin (44):

The preparation followed the general procedure given for 5,15-AB porphyrins using porphyrin **33** (168 mg, 0.43 mmol) in 50 mL THF. Column chromatography on silica gel eluting with ethyl acetate/*n*-hexane (1:3, v/v) gave as the first fraction a small amount of **46** (5 mg, 0.01 mmol, $<1\%$). The second fraction contained a mixture of 5-[4-(*N,N*-dimethylamino)phenyl]-10-(1-ethylpropyl)porphyrin and 5-[4-(*N,N*-dimethylamino)phenyl]-15-(1-ethylpropyl)porphyrin. The third fraction contained 5,10-bis[4-(*N,N*-dimethylamino)phenyl]-15-(1-ethylpropyl)porphyrin **47** (9 mg (0.01 mmol, 2 %). The mixture was chromatographed a second time on silica eluting with CH_2Cl_2 /*n*-hexane (4:1, v/v) and gave **44** (34 mg, 0.07 mmol, 16 %) as the first fraction followed by **45** (94 mg (0.19 mmol, 44 %). All compounds were recrystallized from $\text{CH}_2\text{Cl}_2/\text{MeOH}$. Data for **44**: M.p. 290°C . $R_f = 0.35$ (ethyl acetate/*n*-hexane = 1:3, v/v). ^1H NMR (300 MHz, CDCl_3 , TMS): $\delta = 10.24$ (*s*, 2H, meso- H), 9.73 (*m*, 2H, β -pyrrole- $H_{13,17}$), 9.42 (*br s*, 2H, β -pyrrole- $H_{12,18}$), 9.36 (*AB*, 2H, $^3J = ^3J = 4.5$ Hz, β -pyrrole- $H_{2,8}$), 9.16 (*AB*, 2H, $^3J = ^3J = 4.5$ Hz, β -pyrrole- $H_{3,7}$), 8.14 (*AB*, 2H, $^3J = ^3J = 8.5$ Hz, Ar- H_o), 7.16 (*AB*, 2H, $^3J = ^3J = 8.5$ Hz, Ar- H_m), 5.07 (*m*, 1H, $\text{CH}(\text{CH}_2)_2$), 3.23 (*s*, 6H, $\text{N}(\text{CH}_3)_2$), 2.91 (*m*, 4H, $\text{CH}(\text{CH}_2)_2$), 0.97 (*t*, 6H, $^3J = 7.3$ Hz, CH_2CH_3), -2.63 ppm (*br s*, 2H, NH). ^{13}C NMR (75 MHz, CDCl_3): $\delta = 148.06, 135.98, \sim 131, 131.44, \sim 130, \sim 129, 119.42, 111.33, 104.43, 49.95, 40.80, 34.63, 14.13$ ppm. UV/vis (CH_2Cl_2): λ_{max} ($\log \epsilon$) = 406 (4.69), 508 (3.94), 549 (3.76), 580 (3.57), 638 nm (3.39). MS (EI, 80 eV): m/z

= 499 (66 %, [M]⁺), 470 (100 %, [M – C₂H₅]⁺), 454 (16 %, [M – C₂H₇N]⁺), 250 (16 %, [M]²⁺), 235 (75 %, [M – C₂H₅]²⁺). HRMS calcd. for C₃₃H₃₃N₅ 499.2736; found 499.2722.

5-[4-(*N,N*-Dimethylamino)phenyl]-10-(1-ethylpropyl)porphyrin (45):

Derived from the synthesis of **44**: M.p. 268 °C. *R_f* = 0.35 (ethyl acetate/*n*-hexane = 1:3, v/v). ¹H NMR (300 MHz, CDCl₃, TMS): δ = 10.14 (s, 1H, meso-*H*₁₅), 10.09 (s, 1H, meso-*H*₂₀), 9.73 (m, 2H, β-pyrrole-*H*_{8,12}), 9.37 (m, 3H, β-pyrrole-*H*_{13,17,18}), 9.27 (m, 1H, β-pyrrole-*H*₂), 9.08 (m, 2H, β-pyrrole-*H*_{3,7}), 8.11 (AB, 2H, ³*J* = ³*J* = 8.6 Hz, Ar-*H*_o), 7.15 (AB, 2H, ³*J* = ³*J* = 8.6 Hz, Ar-*H*_m), 5.16 (m, 1H, CH(CH₂)₂), 3.24 (s, 6H, N(CH₃)₂), 2.93 (m, 4H, CH(CH₂)₂), 0.97 (t, 6H, ³*J* = ³*J* = 7.4 Hz, CH₂CH₃), –3.14 ppm (br s, 2H, NH). ¹³C NMR (75 MHz, CDCl₃): δ = 150.03, 135.58, ~131, 130.86, ~130, ~129, 123.68, 122.28, 110.59, 103.15, 50.46, 40.74, 34.99, 14.18 ppm. UV/vis (CH₂Cl₂): λ_{max} (log ε) = 409 (4.33), 507 (3.62), 545 (3.53), 581 (3.54), 637 nm (3.17). MS (EI, 80 eV): *m/z* = 499 (66 %, [M]⁺), 470 (100 %, [M – C₂H₅]⁺), 454 (16 %, [M – C₂H₇N]⁺), 250 (16 %, [M]²⁺), 235 (75 %, [M – C₂H₅]²⁺). HRMS calcd. for C₃₃H₃₃N₅ 499.2736; found 499.2713.

5,10-Di(*n*-butyl)-15-(1-ethylpropyl)porphyrin (46):

Derived from the synthesis of **44**: M.p. 160 °C. *R_f* = 0.55 (ethyl acetate/*n*-hexane = 1:3, v/v). ¹H NMR (300 MHz, CDCl₃, TMS): δ = 9.95 (s, 1H, meso-*H*₂₀), 9.65 (m, 2H, β-pyrrole-*H*_{3,7}), 9.60, (AB, 2H, *J* = 4.9 Hz, β-pyrrole-*H*), 9.51 (AB, 1H, *J* = 4.9 Hz, β-pyrrole-*H*), 9.49 (AB, 1H, *J* = 4.7 Hz, β-pyrrole-*H*), 9.27 (AB, 2H, *J* = 4.7 Hz, β-pyrrole-*H*), 5.08, 4.95 (m, 4H, CH₂C₃H₇), 5.02 (m, 1H, CH(CH₂)₂), 2.87, 2.60 (m, 4H, CH(CH₂)₂), 2.50 (m, 4H, CH₂CH₂C₂H₅), 1.85 (m, 4H, C₂H₄CH₂CH₃), 1.19, 1.13 (t, 6H, ³*J* = ³*J* = 7.3 Hz, C₃H₆CH₃), 0.96 (t, 6H, ³*J* = ³*J* = 7.3 Hz, CH₂CH₃), –2.67, –2.75 ppm (br s, 2H, NH). ¹³C NMR (75 MHz, CDCl₃): δ = ~149, 131.34, ~129, 118.57, 102.94, 50.05, 41.22, 40.60, 36.26, 34.54, 23.79, 23.61, 14.20 ppm. UV/vis (CH₂Cl₂): λ_{max} (log ε) = 412 (5.12), 512 (3.96), 545 (3.57), 589 (3.47), 644 nm (3.38). MS (EI, 80 eV): *m/z* = 492 (85 %, [M]⁺), 463 (100 %, [M – C₂H₅]⁺), 449 (40 %, [M – C₃H₇]⁺), 377 (20 %, [M – 2 C₃H₇ – C₂H₅]⁺), 246 (5 %, [M]²⁺), 232 (4 %, [M – C₂H₅]²⁺). HRMS calcd. for C₃₃H₄₀N₄ 492.3253; found 492.3245.

5,10-Bis[4-(*N,N*-dimethylamino)phenyl]-15-(1-ethylpropyl)porphyrin (47):

Derived from the synthesis of **44**: M.p. 178 °C. *R_f* = 0.27 (ethyl acetate/*n*-hexane = 1:3, v/v). ¹H NMR (300 MHz, CDCl₃, TMS): δ = 10.08 (s, 1H, meso-*H*₂₀), 9.69 (m, 1H, β-pyrrole-*H*₁₇), 9.57 (m, 1H, β-pyrrole-*H*₁₃), 9.34 (AB, 1H, ³*J* = 4.7 Hz, β-pyrrole-*H*₁₈), 9.27 (AB, 1H, ³*J* = 4.6 Hz, β-pyrrole-*H*₁₈), 9.07 (AB, 1H, ³*J* = 4.6 Hz, β-pyrrole-*H*₃), 8.99 (m, 1H, β-pyrrole-*H*₁₂), 8.93, 8.88 (AB, 2H, ³*J* = ³*J* = 4.9 Hz, β-pyrrole-*H*_{7,8}), 8.07 (m, 2H, Ar-*H*_o), 7.10 (m, 2H, Ar-*H*_m), 5.06 (m, 1H, CH(CH₂)₂), 3.23, 3.22 (s, 12H, N(CH₃)₂), 2.89 (m, 4H, CH(CH₂)₂), 0.96 (t, 6H, ³*J* = 7.3 Hz, CH₂CH₃), –2.68 ppm (br s, 2H, NH). ¹³C NMR (75 MHz, CDCl₃): δ = 149.93, 135.85, 135.46, 131.85, 131.53, 130.83, 129.69, ~129, 119.74, 111.01, 110.40, ~103, 50.13, 40.76, 34.62, 14.18 ppm. UV/vis (CH₂Cl₂): λ_{max} (log ε) = 419 (4.82), 517 (4.18), 557 (4.09), 587 (3.83), 649 nm (3.80). MS (EI, 80 eV): *m/z* = 618 (52 %, [M]⁺), 589 (100 %, [M – C₂H₅]⁺), 295 (24 %, [M – C₂H₅]²⁺). HRMS calcd. for C₄₁H₄₂N₆ 618.3471; found 618.3477.

5-(*n*-Butyl)-15-(3-methoxyphenyl)porphyrin (48): The synthesis followed the general procedure given for 5,15-AB porphyrins using 5-(3-methoxyphenyl)porphyrin **34** (80 mg, 0.192 mmol), *n*BuLi (0.46 mL, 1.152 mmol), 0.5 mL H₂O and DDQ (227 mg, 1.152 mmol). Column chromatography using *n*-hexane/CH₂Cl₂ (2:1, v/v) gave first 5,10-di(*n*-butyl)-15-(3-methoxyphenyl)porphyrin **50** (17 mg, 0.032 mmol, 17 %), followed by the title compound **48** (10 mg, 0.021 mmol, 11 %), and then 5-(*n*-butyl)-10-(3-methoxyphenyl)porphyrin **49** (11 mg, 0.023 mmol, 12 %) all as purple crystals. Data for compound **48**: M.p. > 300 °C. *R_f* = 0.62 (*n*-hexane/CH₂Cl₂ = 1:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = –3.00 (s, 2H, NH), 1.20 (t, 3H, *J* = 7.1 Hz, CH₂CH₂CH₂CH₃), 1.88 (m, 2H, CH₂CH₂CH₂CH₃), 2.58 (m, 2H, CH₂CH₂CH₂CH₃), 4.05 (s, 3H, OCH₃), 5.05 (t, 2H, *J* = 7.6 Hz, CH₂CH₂CH₂CH₃), 7.40 (d, 1H, *J* = 7.6 Hz, Ar-*H*), 7.73 (t, 1H, *J* = 7.0 Hz, Ar-*H*), 7.89 (m, 2H, Ar-*H*), 9.13 (d, 2H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.38 (d, 2H, *J* = 4.09 Hz, β-pyrrole-*H*), 9.44 (d, 2H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.62 (d, 2H, *J* = 4.67 Hz, β-pyrrole-*H*), 10.24 ppm (s, 2H, meso-*H*). ¹³C-NMR (100 MHz, CDCl₃, TMS): δ = 13.7, 13.8, 22.3, 23.2, 28.9, 31.5, 34.1, 40.1, 55.1, 104.3, 122.9, 117.5, 119.3, 120.2, 127.3, 127.4, 127.5, 130.2, 131.3, 142.2, 144.0, 144.3, 146.7 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ε) = 406 (5.12), 503 (4.15), 537 (3.97), 577 (4.05), 629 nm (3.67). HRMS calcd. for C₃₁H₂₉N₄O [M+H]⁺ 473.2341; found 473.2364.

5-(*n*-Butyl)-10-(3-methoxyphenyl)porphyrin (49):

Derived from the synthesis of **48**. M.p. > 300 °C. *R_f* = 0.5 (*n*-hexane/CH₂Cl₂ = 1:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = –3.28 (s, 2H, NH), 1.18 (t, 3H, *J* = 7.0 Hz, CH₂CH₂CH₂CH₃), 1.86 (m, 2H, CH₂CH₂CH₂CH₃), 2.59 (m, 2H, CH₂CH₂CH₂CH₃), 4.03 (s, 3H, OCH₃), 5.11 (t, 2H, *J* = 7.6 Hz, CH₂CH₂CH₂CH₃), 7.40 (d, 1H, *J* = 5.84 Hz, Ar-*H*), 7.70 (t, 1H, *J* = 7.6 Hz, Ar-*H*), 7.86 (m, 2H, Ar-*H*), 9.07 (t, 2H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.31 (d, 1H, *J* = 4.67 Hz, β-pyrrole-*H*), 9.41 (m, 3H, β-pyrrole-*H*), 9.58 (d, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.67 (d, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 10.13 (s, 1H, meso-*H*), 10.16 ppm (s, 1H, meso-*H*). ¹³C-NMR (100 MHz, CDCl₃, TMS): δ = 13.7, 13.8, 22.2, 23.7, 28.9, 29.2, 31.5, 34.8, 40.6, 55.0, 102.9, 103.3, 113.0, 118.6, 120.0, 120.2, 126.9, 127.2, 143.3, 157.4 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ε) = 406 (5.45), 502 (4.45), 534 (3.97), 577 (4.27), 632 nm (4.15). HRMS calcd. for C₃₁H₂₉N₄O [M+H]⁺ 473.2341; found 473.2357.

5,10-Di(*n*-Butyl)-15-(3-methoxyphenyl)porphyrin (50):

Derived from the synthesis of **48**. M.p. > 300 °C. *R_f* = 0.62 (*n*-hexane/CH₂Cl₂ = 1:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = –2.90 (s, 2H, NH), 1.18 (dd, 6H, *J* = 7.32 Hz, CH₂CH₂CH₂CH₃), 1.88 (m, 4H, CH₂CH₂CH₂CH₃), 2.57 (m, 4H, CH₂CH₂CH₂CH₃), 4.03 (s, 3H, OCH₃), 5.08 (m, 4H, CH₂CH₂CH₂CH₃), 7.37 (d, 1H, *J* = 8.05 Hz, Ar-*H*), 7.68 (t, 1H, *J* = 8.05, 7.32 Hz, Ar-*H*), 7.81 (m, 2H, Ar-*H*), 8.98 (t, 2H, *J* = 4.39 Hz, β-pyrrole-*H*), 9.25 (d, 1H, *J* = 4.39 Hz, β-pyrrole-*H*), 9.35 (d, 1H, *J* = 4.39 Hz, β-pyrrole-*H*), 9.54 (d, 1H, *J* = 4.39 Hz, β-pyrrole-*H*), 9.60 (d, 1H, *J* = 4.39 Hz, β-pyrrole-*H*), 9.62 (d, 1H, *J* = 4.4 Hz, β-pyrrole-*H*), 9.66 (d, 1H, *J* = 5.12 Hz, β-pyrrole-*H*), 10.04 ppm (s, 1H, meso-*H*). ¹³C-NMR (100 MHz, CDCl₃, TMS): δ = 14.0, 23.5, 34.9, 35.6, 40.7, 40.9, 55.3, 103.3, 113.2, 117.7, 119.4, 120.3, 127.4, 127.5, 143.2, 157.9 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log

ϵ) = 412 (5.08), 510 (4.06), 544 (3.46), 585 (3.76), 643 nm (3.36). HRMS calcd. for $C_{35}H_{36}N_4O$ $[M+H]^+$ 529.2967; found 529.2980.

5-Hexyl-10-(4-methoxyphenyl)porphyrin (52): Prepared similar to the general procedure given for the synthesis of 5,10-AB-type porphyrins using 5-(4-methoxyphenyl)porphyrin **35** (34 mg, 0.08 mmol) and a 2.5 M solution of *n*-hexyllithium in hexane (0.11 mL, 0.28 mmol, 3.5 equiv.). Column chromatography on silica gel (CH_2Cl_2/n -hexane = 2:1, v/v) gave a first fraction containing a mixture of higher alkylated porphyrins. The second fraction contained a mixture of 5-hexyl-10-(4-methoxyphenyl)porphyrin **52** and 5-hexyl-15-(4-methoxyphenyl)porphyrin **51** in a ratio of 2.89 : 1 (determined via analytical HPLC; CH_2Cl_2/n -hexane = 1:1, v/v). A second column chromatography on silica gel (CH_2Cl_2/n -hexane = 2:3, v/v) gave a first fraction of **51** (5 mg (< 0.01 mmol, 11 %),^[11] and a second fraction that gave purple crystals of the title compound after recrystallization from $CH_2Cl_2/MeOH$ (14 mg (0.03 mmol, 32 %). M.p. 163 °C. R_f = 0.16 (CH_2Cl_2/n -hexane = 2:3, v/v). 1H NMR (500 MHz, $CDCl_3$, TMS): δ = 10.051 (s, 1H, meso- H_{20}), 10.046 (s, 1H, meso- H_{15}), 9.56 (AB, 1H, J = 4.7 Hz, β -pyrrole- H_3), 9.51 (AB, 1H, J = 4.8 Hz, β -pyrrole- H_7), 9.31 (m, 3H, β -pyrrole- $H_{2,17,18}$), 9.25 (AB, 1H, J = 4.5 Hz, β -pyrrole- H_{13}), 9.03 (AB, 1H, J = 4.8 Hz, β -pyrrole- H_8), 9.01 (AB, 1H, J = 4.5 Hz, β -pyrrole- H_{12}), 8.13 (AB, 2H, 3J = 3J = 8.5 Hz, Ar- H_o), 7.30 (AB, 2H, 3J = 3J = 8.5 Hz, Ar- H_m), 4.99 (t, 2H, 3J = 8.1 Hz, $CH_2C_3H_{11}$), 4.09 (s, 3H, OCH_3), 2.56 (m, 2H, $CH_2CH_2C_4H_9$), 1.82 (m, 2H, $C_2H_4CH_2C_3H_7$), 1.53 (m, 2H, $C_3H_6CH_2C_2H_5$), 1.42 (m, 2H, $C_4H_8CH_2CH_3$), 0.98 (t, 3H, 3J = 7.3 Hz, $C_5H_{10}CH_3$), -3.31 ppm (br s, 2H, NH). ^{13}C -NMR (126 MHz, $CDCl_3$, TMS): δ = 159.55, ~146, 135.71, 134.95, ~132, ~131, ~129, ~128, 120.72, 119.44, 112.26, 103.64, 103.44, 55.70, 39.07, 35.64, 32.05, 30.41, 22.88, 14.31 ppm. UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 407 (5.03), 503 (3.96), 536 (3.47), 578 (3.47) 633 nm (3.19). MS (TOF, MS, ES⁺, 70 eV): m/z = 500 (100 %, $[M]^+$). HRMS calcd. for $C_{33}H_{32}N_4O$ 500.2576; found 500.2571.

5-(4-Methoxyphenyl)-15-phenylporphyrin (53) and 5-(4-Methoxyphenyl)-10-phenylporphyrin (54): Prepared similar to the general procedure given for the synthesis of 5,10-AB-type porphyrins using 5-(4-methoxyphenyl)porphyrin **35** (34 mg, 0.08 mmol) and a 1.8 M solution of PhLi in dibutyl ether (0.43 mL, 0.78 mmol, 6.5 equiv.). The crude product was purified via column chromatography on silica gel (*n*-hexane/ethylacetate = 5:1, v/v) yielding a first fraction of higher alkylated porphyrins. The second fraction contained an inseparable mixture of 5-(4-methoxyphenyl)-10-phenylporphyrin **54** and 5-(4-methoxyphenyl)-15-phenylporphyrin **53** in a ratio of 7:1 (NMR). Analytical data for the mixture: R_f = 0.56 (*n*-hexane/ethylacetate = 5:1, v/v). UV/vis (CH_2Cl_2): λ_{max} (rel. intens.): 408 (1.00), 503 (0.06), 537 (0.02), 576 (0.03) 631 nm (0.01). MS (EI, 70 eV): m/z = 492 (97 %, $[M]^+$), 477 (6 %, $[M - CH_3]^+$), 246 (36 %, $[M]^{2+}$). HRMS: calcd. for $C_{33}H_{24}N_4O$ 492.1950; found 492.1936.

General procedure for the synthesis of 5,15-AB₂/A₂B-type porphyrins

A 25 mL Schlenk flask was charged with dried porphyrin (0.05 mmol) dissolved in 10 mL THF. The solution was cooled to -80 °C and treated with 0.2 mmol of a 2.5 M solution of hexyllithium (80 μ L). The color changed from red to green-brown and after 10 min changed to purple. The cold bath was removed and the solution stirred for 15 min at rt. Next water (1 mL) was added and stirring was continued for 15 min. This was accompanied by a color change to yellow-brown. Then DDQ (0.1 mmol, 23 mg) in 0.4 mL THF was added and the mixture stirred for 15 min until the color had changed to dark red. The mixture was filtered through silica gel and the solvent removed *in vacuo*.

5-(9-Anthracenyl)-10,15-dihexylporphyrin (55): The procedure followed that given above using porphyrin **36** (20 mg, 0.04 mmol), 0.2 mmol of a 2.5 M solution of hexyllithium (80 μ L) and 18 mg of DDQ (0.08 mmol). Column chromatography eluting with *n*-hexane/ CH_2Cl_2 = 1:3, v/v) gave a purple solid (14 mg, 0.02 mmol, 54 %). M.p. >330 °C. 1H NMR (500 MHz, $CDCl_3$, TMS): δ = 9.97 (s, 1H, meso- H_{20}), 9.62 (bs, 2H, β -pyrrole- $H_{12,13}$), 9.59 (d, 1H, J = 4.7 Hz, β -pyrrole- H_8), 9.32 (dd, 2H, J = J = 4.7 Hz, β -pyrrole- $H_{2,7}$), 9.04 (d, 1H, J = 4.7 Hz, β -pyrrole- H_{18}), 8.93 (s, 1H, Ar- H), 8.32 (dd, 2H, J = J = 4.7 Hz, β -pyrrole- $H_{3,17}$), 8.29 (d, 2H, J = 9.0 Hz, Ar- H), 7.46 (dd, 2H, 3J = 9.0 Hz, 4J = 6.0 Hz, Ar- H), 7.09 (d, 2H, J = 9.0 Hz, Ar- H), 6.99 (dd, 2H, 3J = 9.0 Hz, 4J = 6.5 Hz, Ar- H), 5.05, 5.00 (t, 4H, 3J = 8.1 Hz, $CH_2CH_2CH_2CH_2CH_2CH_3$), 2.61, 2.57 (m, 4H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.83 (m, 4H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.55, 1.50 (m, 4H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.44, 1.40 (m, 4H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 0.97, 0.93 (t, 6H, J = 7.4 Hz, $CH_2CH_2CH_2CH_2CH_2CH_3$), -2.63 ppm (bs, 2H, NH). ^{13}C -NMR (126 MHz, $CDCl_3$, TMS): δ = 135.68, 135.19, 130.88, 128.62, 128.19, 127.88, 125.65, 125.01, 120.66, 120.21, 113.09, 103.55, 39.02, 38.78, 35.98, 35.40, 31.86, 30.33, 22.71, 14.15 ppm. UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 415 (5.39), 512 (4.16), 544 (3.65), 587 (3.63), 643 nm (3.45). MS (EI, 290 °C, 80 eV): m/z = 656 (34 %, $[M + 2]^+$), 655 (60 %, $[M + 1]^+$), 654 (100 %, $[M]^+$), 583 (60 %, $[M - C_5H_{11}]^+$), 512 (13 %, $[M - C_{10}H_{22}]^+$), 327 (8 %, $[M]^{2+}$). HRMS calcd. for $C_{46}H_{46}N_4$ 654.37225; found 654.37225.

5,15-Dihexyl-10-pentylporphyrin (57): The procedure followed that given above using porphyrin **37** (6 mg, 0.02 mmol), 40 μ L of a 2.5 M solution of hexyllithium (0.1 mmol) and 9.1 mg of DDQ (0.04 mmol). Column chromatography eluting with *n*-hexane/ CH_2Cl_2 = 1:1, v/v) gave a mixture of two trisubstituted porphyrins. Using analytical HPLC one porphyrin could be separated and identified as the title compound (2 mg, 0.004 mmol, 19 %). M.p. 142 °C. 1H NMR (500 MHz, $CDCl_3$, TMS): δ = 9.92 (s, 1H, meso- H_{20}), 9.57 (AB, 2H, 3J = 3J = 4.8 Hz, β -pyrrole- $H_{8,12}$), 9.51 (AB, 2H, 3J = 3J = 4.8 Hz, β -pyrrole- $H_{7,13}$), 9.49 (AB, 2H, 3J = 3J = 4.6 Hz, β -pyrrole- $H_{3,17}$), 9.27 (AB, 2H, 3J = 3J = 4.6 Hz, β -pyrrole- $H_{2,18}$), 5.04 (t, 2H, 3J = 8.1 Hz, 10- $CH_2CH_2CH_2CH_2CH_3$), 4.95 (t, 4H, 3J = 8.1 Hz, 5,15- $CH_2CH_2CH_2CH_2CH_2CH_3$), 2.55 (m, 2H, 10- $CH_2CH_2CH_2CH_2CH_2CH_3$), 2.51 (m, 4H, 5,15- $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.84 (m, 2H, 10- $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.78 (m, 4H, 5,15- $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.52 (m, 2H, 10-

CH₂CH₂CH₂CH₂CH₃), 1.50 (*m*, 4H, 5,15-CH₂CH₂CH₂CH₂CH₂CH₃), 1.40 (*m*, 4H, 5,15-CH₂CH₂CH₂CH₂CH₂CH₃), 0.96 (*t*, 3H, ³*J* = 7.4 Hz, 10-CH₂CH₂CH₂CH₂CH₃), 0.94, 0.91 (*t*, 6H, ³*J* = 7.3 Hz, 5,15-CH₂CH₂CH₂CH₂CH₂CH₃), -2.88 ppm (*bs*, 2H, NH). ¹³C-NMR (126 MHz, CDCl₃, TMS): δ = 131.30, 128.77, 128.44, 119.92, 118.63, 102.94, 39.08, 38.84, 36.21, 35.02, 32.73, 31.92, 30.37, 22.78, 14.17 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ε) = 411 (5.43), 511 (4.11), 544 (3.94), 588 (3.87), 646 nm (3.73). MS (EI, 290 °C, 80 eV): *m/z* = 549 (59 %), [M + 1]⁺, 548 (100 %), [M]⁺, 491 (20 %), [M - C₄H₉]⁺, 477 (51 %), [M - C₅H₁₁]⁺, 349 (10 %), [M - C₁₄H₃₁]⁺, 71 (14 %), [C₅H₁₁]⁺, 57 (27 %), [C₄H₉]⁺. HRMS calcd. for C₃₇H₄₈N₄ 548.38790; found 548.38366.

General procedure for the synthesis of ABC-type porphyrins

Porphyrin (0.1 mmol) was dissolved in THF and cooled to -78 °C. Hexyllithium was added dropwise over the course of 15 min. The cold bath was removed and stirring was continued for another 15 min. This was followed by addition of 0.5 mL water in 5 mL THF, stirring for 15 min and addition of 6 mL of a solution of DDQ in THF. The mixture was stirred for 20 min and then filtered through neutral alumina washing with dichloromethane.

5-(4-Methylphenyl)-10-phenyl-15-(2,4,6-trimethoxyphenyl)porphyrin (62):

The porphyrin **21** (55 mg, 0.1 mmol) was dissolved in THF and cooled to 0 °C. Phenyllithium was added dropwise over the course of 15 min and subsequent workup followed the general procedure given above. Column chromatography (*n*-hexane/CH₂Cl₂ = 1:1, v/v) gave a small first fraction containing educt followed by the product as a purple solid after evaporation of the solvent (96 mg, 0.15 mmol, 75 %). M.p. >300 °C. ¹H NMR (500 MHz, CDCl₃, TMS): δ = 10.12 (*s*, 1H, meso-*H*), 8.94 (*m*, 8H, β-pyrrole-*H*), 8.12 (*m*, 4H, Ar-*H*_o), 7.73 (*m*, 3H, Ph-*H*_{mp}), 7.54 (*m*, 2H, tolyl-*H*_o), 6.56 (*s*, 2H, Ar-*H*_m), 4.03 (*s*, 3H, *p*-OCH₃), 3.53 (*s*, 6H, *o*-OCH₃), 2.73 (*s*, 3H, Ph-CH₃), -3.07 ppm (*bs*, 2H, NH). UV/vis (CH₂Cl₂): λ_{max} (log ε) = 411 (5.22), 508 (4.16), 542 (3.65), 584 (3.63), 637 nm (3.23). MS (310 °C, 8 kV): *m/z* = 643.0 (100.00 %), [M+H]⁺, 321.0 (8.99 %), [M]²⁺. HRMS calcd. for C₄₂H₃₄N₄O₃ 642.26309; found 642.26314.

5-(4-Methoxyphenyl)-10-(4-methylphenyl)-20-(2,4,6-trimethoxyphenyl)porphyrin (64):

Synthesis and workup followed the procedure given above to yield purple crystals (120 mg (0.18 mmol, 89 %). M.p. > 320 °C. ¹H NMR (500 MHz, CDCl₃, TMS): δ = 10.23 (*s*, 1H, meso-*H*), 9.33 (*m*, 4H, β-pyrrole-*H*), 9.08 (*d*, 2H, ³*J* = 4.54 Hz, β-pyrrole-*H*), 9.02 (*d*, 2H, ³*J* = 4.54 Hz, β-pyrrole-*H*), 8.16 (*d*, 2H, ³*J* = 7.81 Hz, tolyl-*H*_o), 7.61 (*d*, 2H, ³*J* = 7.45 Hz, tolyl-*H*_m), 6.63 (*s*, 2H, OMe-Ar-*H*_m), 4.12 (*s*, 3H, *p*-OH₃), 3.53 (*s*, 6H, *o*-OH₃), 2.71 (*s*, 3H, C₆H₄-CH₃), -3.05 ppm (*bs*, 2H, NH). ¹³C NMR (126 MHz, CDCl₃): δ = 162.03, 161.21, 159., 148.84-146.42, 139.13, 137.20, 135.42, 134.60, 131.56-130.08, 127.46, 119.62, 111.95, 104.02, 91.09, 56.00, 55.59, 55.47, 21.47 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ε) = 411 (5.20), 508 (4.00), 542 (3.60), 580 (3.57), 637 nm (3.03). MS (320 °C, 80 eV): *m/z* = 672.0 (100.00 %), [M]⁺, 641.0 (12 %), [M - C₁H₃O]⁺. HRMS: calcd. for C₄₃H₃₆N₄O₄ 672.27366; found 672.27370..

General procedure for the metallation with Manganese chloride

Free base porphyrin (100mg, 0.14 mmol) was dissolved together with 150 mg MnCl₂·4 H₂O in 25 mL acetic acid (96 %) and 7 mL acetic anhydride. The solution was heated for 4 h at 110 °C. After cooling to rt dichloromethane was added and the organic phase washed several times with water. The organic solvent was evaporated and the residue dried under high vacuum.

Chloro[5-(4-methylphenyl)-10-phenyl-15-(2,4,6-trimethoxyphenyl)porphyrinato]manganese(III) (66):

The free base porphyrin **62** (100 mg, 0.16 mmol) was treated as described in the procedure above. Column chromatography eluting with dichloromethane gave two minor fractions followed by the metal complex. Evaporation of the solvent gave a gree-red residue (114 mg, 0.155 mmol, 98 %). M.p. >320 °C. UV/Vis (CH₂Cl₂): λ_{max} (log ε) = 373 (4.38), 405 (4.32), 477 (4.60), 581 (3.72), 614 (3.66), 643 nm (3.45). MS (FAB⁺, 3 kV): *m/z* = calcd. for C₄₂H₃₂ClMnN₄O₃ 730.15434; found 730.0 (14.0 %), [M]⁺, 695.0 (100.0 %), [M - Cl]⁺.

5-[4-(*N,N*-Dimethylamino)phenyl]-15-(1-ethylpropyl)-10-hexylporphyrin (67):

Porphyrin **44** (51 mg, 0.10 mmol) was reacted with a 2.5 M solution of hexyllithium in hexane (0.10 mL, 0.25 mmol) as described in the general procedure for the synthesis of ABC-type porphyrins. Oxidation was performed by stirring for 12 h under air. Column chromatography on silica gel (CH₂Cl₂/*n*-hexane = 2:1, v/v) gave a single fraction which yielded purple crystals after recrystallization from CH₂Cl₂/MeOH/H₂O (31 mg, 0.05 mmol, 52 %). M.p. 138 °C. *R*_f = 0.44 (CH₂Cl₂/*n*-hexane, 2:1, v/v). ¹H NMR (300 MHz, CDCl₃, TMS): δ = 10.01 (*s*, 1H, meso-*H*), 9.70 (*m*, 2H, β-pyrrole-*H*_{13,17}), 9.63 (*m*, 1H, β-pyrrole-*H*₁₂), 9.52 (*AB*, 1H, ³*J* = 4.6 Hz, β-pyrrole-*H*₈), 9.30 (*AB*, 1H, ³*J* = 4.6 Hz, β-pyrrole-*H*₁₈), 9.23 (*AB*, 1H, ³*J* = 4.6 Hz, β-pyrrole-*H*₂), 9.05 (*AB*, 1H, ³*J* = 4.6 Hz, β-pyrrole-*H*₇), 9.04 (*AB*, 1H, ³*J* = 4.6 Hz, β-pyrrole-*H*₃), 8.09 (*AB*, 2H, ³*J* = ³*J* = 8.7 Hz, Ar-*H*_o), 7.13 (*AB*, 2H, ³*J* = ³*J* = 8.7 Hz, Ar-*H*_m), 5.08 (*m*, 3H, CH(CH₂)₂, CH₂C₅H₁₁), 3.23 (*s*, 6H, N(CH₃)₂), 2.90 (*m*, 4H, CH(CH₂)₂), 2.61 (*m*, 2H, CH₂CH₂C₄H₉), 1.87 (*m*, C₂H₄CH₂C₃H₇), 1.56 (*m*, 2H, C₃H₆CH₂C₂H₅), 1.45 (*m*, 2H, C₄H₈CH₂CH₃), 0.98 (*t*, 6H, ³*J* = 7.4 Hz, CH₂CH₃), 0.97 (*t*, 3H, ³*J* = 7.3 Hz, C₅H₁₀CH₃), -2.72 ppm (*br s*, 2H, NH). ¹³C NMR (63 MHz, CDCl₃): δ = 149.96, ~145, 135.72, 131.30, 130.66, 129.79, 128.27, 122.03, 119.31, 110.97, 103.31, 50.18, 40.76, 39.09, 36.50, 34.65, 31.94, 30.40, 22.78 14.18 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ε) = 415 (4.19), 515 (3.96), 554 (3.80), 591 (3.54), 649 nm (3.54). MS (EI, 70 eV): *m/z* 583 (4 %), [M]⁺, 554 (8 %), [M - C₂H₅]⁺, 512 (8 %), [M - C₅H₁₁]⁺, 483 (4 %), [M - C₅H₁₁ - C₂H₅]⁺, 292 (33 %), [M]²⁺, 242 (59 %), [M - C₅H₁₁ - C₂H₅]²⁺. HRMS calcd. for C₃₉H₄₅N₅ 583.3675; found 583.3665.

5-Butyl-10-hexyl-15-(3-methoxyphenyl)porphyrin (68):

Following the general procedure given for the synthesis of 5,15-AB-type porphyrins (organolithium reaction), 5-butyl-15-(3-methoxyphenyl)porphyrin **48** (100 mg, 0.211 mmol), hexyllithium (0.551 mL, 1.26 mmol), 0.5 mL water and DDQ, (288.2 mg, 1.26 mmol) were used. The crude reaction mixture was filtered through silicagel, eluting with *n*-hexane/CH₂Cl₂ = 2:1 (v/v) and afforded the title compound

(70 mg, 0.125 mmol, 60 %) as purple crystals. M.p. > 300 °C. R_f = 0.7 (CH₂Cl₂/*n*-hexane, 1:2, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.90 (s, 2H, NH), 0.99 (t, 3H, J = 7.02 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 1.20 (t, 3H, J = 7.01 Hz, CH₂CH₂CH₂CH₃), 1.45 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.57 (m, 4H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.87 (m, 4H, CH₂), 2.59 (m, 4H, CH₂), 4.03 (s, 3H, OCH₃), 5.05 (m, 4H, CH₂), 7.38 (m, 1H, Ar-H), 7.70 (t, 1H, J = 7.6 Hz, Ar-H), 7.84 (m, 2H, Ar-H), 9.01 (t, 2H, J = 4.67 Hz, β -pyrrole-H), 9.25 (d, 1H, J = 4.1 Hz, β -pyrrole-H), 9.33 (d, 1H, J = 4.68 Hz, β -pyrrole-H), 9.54 (d, 1H, J = 5.26 Hz, β -pyrrole-H), 9.57 (d, 1H, J = 4.67 Hz, β -pyrrole-H), 9.60 (d, 1H, J = 4.68 Hz, β -pyrrole-H), 9.63 (d, 1H, J = 5.26 Hz, β -pyrrole-H), 10.03 ppm (s, 1H, meso-H). ¹³C NMR (100 MHz, CDCl₃): δ = 13.7, 13.8, 22.3, 23.2, 29.9, 31.5, 34.6, 35.6, 38.6, 40.4, 55.0, 103.0, 112.9, 117.4, 119.1, 120.0, 120.1, 127.1, 127.2, 127.6, 128.0, 130.5, 142.8, 157.6 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ϵ) = 412 (4.82), 511 (3.92), 545 (3.74), 588 (3.64), 641 nm (3.34). HRMS (ES) calcd. for C₃₅H₃₆N₄O [M+H]⁺ 557.3280; found 557.3280.

5-sec-Butyl-10-hexyl-20-(3,4,5-trimethoxyphenyl)porphyrin (69):

Following the general procedure given for the synthesis of ABC-type porphyrins, 5-hexyl-15-(3,4,5-trimethoxyphenyl)porphyrin **31** (160 mg, 0.285 mmol), *sec*-BuLi (1.22 mL, 1.712 mmol), 0.5 mL H₂O and DDQ (243.3 mg, 1.088 mmol) were used. The crude reaction mixture was purified by column chromatography using *n*-hexane/ethyl acetate = 10:1 (v/v) affording the title compound (84 mg, 0.136 mmol, 47 %) as purple crystals: Mp > 300 °C; R_f = 0.4 (*n*-hexane/CH₂Cl₂ = 1:2, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.94 (s, 2H, NH), 0.99 (t, 3H, J = 7.02 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 1.09 (t, 3H, J = 7.01 Hz, CHCH₂CH₃), 1.40 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.48 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.88 (m, 2H, CH₂CH₂CH₂CH₂CH₃), 2.47 (d, 3H, J = 7.01 Hz, CHCH₃), 2.56 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.87 (m, 1H, CHCH₂CH₃), 3.00 (m, 2H, CHCH₂CH₃), 4.00 (s, 6H, OCH₃), 4.21 (s, 3H, *p*-OCH₃), 5.06 (t, 2H, J = 8.1 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 5.48 (m, 1H, CHCH₂CH₃), 7.47 (s, 2H, Ar-H), 9.00 (d, 2H, J = 4.1 Hz, β -pyrrole-H), 9.25 (d, 1H, J = 4.67 Hz, β -pyrrole-H), 9.35 (d, 1H, J = 4.68 Hz, β -pyrrole-H), 9.60 (d, 1H, J = 4.68 Hz, β -pyrrole-H), 9.62 (d, 1H, J = 4.68 Hz, β -pyrrole-H), 9.67 (d, 1H, J = 4.1 Hz, β -pyrrole-H), 9.79 (d, 1H, J = 4.09 Hz, β -pyrrole-H), 10.03 ppm (s, 1H, meso-H). ¹³C NMR (100 MHz, CDCl₃): δ = 13.7, 13.8, 22.3, 27.0, 29.2, 29.9, 31.5, 35.2, 35.5, 38.5, 42.6, 55.9, 60.8, 103.1, 112.2, 117.3, 119.2, 125.0, 137.1, 137.4, 150.9 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ϵ) = 414 (4.81), 511 (3.79), 544 (3.48), 592 (3.56), 644 nm (3.39). HRMS (ES+) calcd. for C₃₉H₄₄N₄O₃ [M+H]⁺ 617.3492; found 617.3480.

5,10-Bis(2-methoxyphenyl)-15-(4-methylphenyl)porphyrin (70):

The synthesis followed the procedure given for **63** by adding 100 mg **20** (0.2 mmol), dissolved in 40 ml THF to the LiR solution. The product was obtained as a purple residue as a mixture of two atropisomers (100 mg (0.16 mmol, 80 %)). The two atropisomers could be separated by HPLC (eluting with dichloromethane/*n*-hexane, 3:2, v/v) and were stored at -20 °C. Even during NMR measurements at rt the original

equilibrium between the two isomers was restored. M.p. 304 °C. ¹H NMR (500 MHz, CDCl₃, TMS): δ = 10.14 (s, 1H, meso-H), 9.28 (m, 2H, β -pyrrole-H), 9.00 (AB, 1H, 3J = 4.65 Hz, β -pyrrole-H₁₇), 8.90 (m, 1H, β -pyrrole-H₁₃), 8.88 (AB, 1H, 3J = 4.65 Hz, β -pyrrole-H₁₂), 8.79 (m, 3H, β -pyrrole-H_{3,7,8}), 8.11 (m, 2H, tolyl-H_o), 8.00 (m, 2H, Ar-H_m), 7.76 (m, 2H, Ar-H_p), 7.76 (m, 2H, tolyl-H_m), 7.34 (m, 4H, Ar-H_m), 3.57 and 3.54 (each s, 2 $\times$ 3H, OCH₃), 2.70 (s, 3H, C₆H₄CH₃), -2.91 ppm (bs, 2H, NH). UV/Vis (CH₂Cl₂): λ_{max} (log ϵ) = 413 (5.41), 510 (4.22), 545 (3.80), 584 (3.70), 635 nm (3.36). MS (300 °C, 80 eV): m/z = 612.0 (100.00 %, [M]⁺), 506.0 (41 %, [M - C₇H₇O]⁺), 306.0 (7 %, [M]²⁺). HRMS: calcd. for C₄₁H₃₂N₄O₂ 612.25253; found 612.25256.

5-[4-(*N,N*-Dimethylamino)phenyl]-10-(1-ethylpropyl)-20-hexylporphyrin (74): Porphyrin **45** (48 mg, 0.10 mmol) was reacted with hexyllithium as described in the general procedure for the synthesis of ABC-type porphyrins. Column chromatography on silica gel gave the product in the first fraction which yielded purple crystals after recrystallization from CH₂Cl₂/MeOH/H₂O (25 mg, 0.04 mmol, 46 %). M.p. 92 °C. R_f = 0.36 (CH₂Cl₂/*n*-hexane, 2:1, v/v). ¹H NMR (300 MHz, CDCl₃, TMS): δ = 10.04 (s, 1H, meso-H), 9.66 (m, 1H, β -pyrrole-H₁₂), 9.54 (m, 1H, β -pyrrole-H₈), 9.52 (AB, 1H, 3J = 4.7 Hz, β -pyrrole-H₁₈), 9.40 (AB, 1H, 3J = 4.9 Hz, β -pyrrole-H₂), 9.33 (AB, 2H, 3J = 4.9 Hz, β -pyrrole-H_{13,17}), 8.97 (AB, 2H, 3J = 4.9 Hz, β -pyrrole-H_{3,7}), 8.06 (AB, 2H, 3J = 3J = 8.6 Hz, Ar-H_o), 7.10 (AB, 2H, 3J = 3J = 8.7 Hz, Ar-H_m), 5.03 (m, 1H, CH(CH₂)₂), 4.95 (t, 2H, 3J = 7.8 Hz, CH₂C₅H₁₁), 3.24 (s, 6H, N(CH₃)₂), 2.87 (m, 4H, CH(CH₂)₂), 2.51 (m, 2H, CH₂CH₂C₄H₉), 1.78 (m, 2H, C₂H₄CH₂C₃H₇), 1.49 (m, 2H, C₃H₆CH₂C₂H₅), 1.38 (m, 2H, C₄H₈CH₂CH₃), 0.95 (t, 6H, 3J = 7.4 Hz, CH₂CH₃), 0.92 (t, 3H, 3J = 7.3 Hz, C₅H₁₀CH₃), -2.65 ppm (br s, 2H, NH). ¹³C NMR (63 MHz, CDCl₃): δ = 149.95, 135.38, 132.37, 131.44, ~129, ~127, ~120, ~119, 110.46, ~103, 50.03, 40.85, 38.47, 34.74, 34.54, 31.92, 30.19, 22.71, 14.12 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ϵ) = 413 (4.73), 513 (3.92), 549 (3.59), 588 (3.46), 645 nm (3.41). MS (EI, 70 eV): m/z 583 (38 %, [M]⁺), 554 (69 %, [M - C₂H₅]⁺), 512 (27 %, [M - C₅H₁₁]⁺), 483 (13 %, [M - C₅H₁₁ - C₂H₅]⁺), 292 (52 %, [M]²⁺). HRMS calcd. for C₃₉H₄₅N₅ 583.3675; found 583.3652.

5-[4-(*N,N*-Dimethylamino)phenyl]-10-(1-ethylpropyl)-20-(1-methylpropyl)porphyrin (75): Porphyrin **45** (48 mg, 0.10 mmol) was reacted with a 1.3 M solution of *sec*-butyllithium in cyclohexane/*n*-hexane (0.30 mL, 0.39 mmol) as described in the general procedure for the synthesis of ABC-type porphyrins. Column chromatography on silica gel gave a single fraction which yielded purple crystals after recrystallization from CH₂Cl₂/MeOH (19 mg (0.03 mmol, 31 %)). M.p. 140 °C. R_f = 0.53 (*n*-hexane/CH₂Cl₂, 3:1, v/v). ¹H NMR (300 MHz, CDCl₃, TMS): δ = 10.04 (s, 1H, meso-H), 9.64 (m, 2H, β -pyrrole-H_{8,12}), 9.51 (m, 2H, β -pyrrole-H_{2,18}), 9.32 (m, 2H, β -pyrrole-H_{13,17}), 8.92 (m, 2H, β -pyrrole-H_{3,7}), 8.02 (AB, 2H, 3J = 3J = 8.7 Hz, Ar-H_o), 7.08 (AB, 2H, 3J = 3J = 8.7 Hz, Ar-H_m), 5.31 (m, 1H, CH(CH₃)CH₂), 4.98 (m, 1H, CH(CH₂)₂), 3.24 (s, 6H, N(CH₃)₂), 2.85 (m, 6H, CH(CH₂)₂, CH(CH₃)CH₂), 2.38 (d, 3H, 3J = 7.4 Hz, CH(CH₂)₂(CH₃)), 1.01 (t, 3H, 3J = 7.3 Hz, CH(CH₃)CH₂CH₃), 0.93 (t, 6H, 3J = 7.4 Hz, CH(CH₂)₂(CH₃)), -2.49 ppm (br s, 2H, NH). UV/Vis (CH₂Cl₂): λ_{max} (log ϵ) = 413 (4.49), 513 (3.69), 548 (3.39), 587 (3.26), 646 nm (3.23). MS (EI, 70 eV): m/z 555

(33 %, $[M]^{*+}$), 526 (25 %, $[M - C_2H_5]^+$). HRMS calcd. for $C_{37}H_{42}N_5$ 556.3440; found 556.3444.

General procedure for bromination of 5,15-AB-type porphyrins

The porphyrin (1 equiv.) was dissolved in 100 mL of chloroform and cooled to 0 °C. Pyridine (1.0 mL) was added to act as an acid scavenger. NBS (1.1 equiv.) was added directly to the flask, and the reaction was followed by TLC. When the reaction reached completion it was quenched with 10 mL of acetone. The solvents were evaporated under reduced pressure, and the product was dissolved in CH_2Cl_2 followed by filtration through silica gel.

5-Bromo-10-hexyl-20-(2,4,6-trimethoxyphenyl)porphyrin (78):

The compound was derived during the preparation of **79**. M.p. > 300 °C. R_f = 0.5 (*n*-hexane/ CH_2Cl_2 , 1:1, v/v). 1H NMR (400 MHz, $CDCl_3$, TMS): δ = -2.98 (s, 2H, NH), 0.97 (t, 3H, J = 7.01 Hz, $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.41 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.51 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.78 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 2.49 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 3.58 (s, 6H, OCH₃), 4.15 (s, 3H, *p*-OCH₃), 4.81 (t, 2H, J = 7.6 Hz, $CH_2CH_2CH_2CH_2CH_2CH_3$), 6.65 (s, 2H, Ar-*H*), 8.94 (d, 2H, J = 4.68 Hz, β -pyrrole-*H*), 9.20 (d, 1H, J = 4.68 Hz, β -pyrrole-*H*), 9.22 (d, 1H, J = 4.68 Hz, β -pyrrole-*H*), 9.38 (d, 1H, J = 4.67 Hz, β -pyrrole-*H*), 9.41 (d, 1H, J = 4.68 Hz, β -pyrrole-*H*), 9.74 (d, 1H, J = 4.67 Hz, β -pyrrole-*H*), 9.97 ppm (s, 1H, meso-*H*). ^{13}C NMR (100 MHz, $CDCl_3$): δ = 14.1, 22.7, 30.2, 31.8, 35.1, 38.7, 55.6, 55.9, 90.9, 104.5, 120.7, 128.4, 131.5, 132.3, 132.7, 161.1, 162.1 ppm. UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 414 (5.25), 513 (4.20), 547 (3.50), 585 (3.65), 648 nm (3.28). HRMS (ES^+) calcd. for $C_{35}H_{36}N_4O_3Br$ $[M+H]^+$ 639.1971; found 639.1983.

5,15-Dibromo-10-hexyl-20-(2,4,6-trimethoxyphenyl)porphyrin (79):

Following the general procedure for bromination 5-hexyl-15-(2,4,6-trimethoxyphenyl)porphyrin **27** (145.2 mg, 0.217 mmol), NBS (46.4 mg, 0.33 mmol) and acetone (10 mL) were used. The product was purified using column chromatography eluting with *n*-hexane/ CH_2Cl_2 (1:2, v/v). The first fraction was 5,15-dibromo-10-hexyl-20-(2,4,6-trimethoxyphenyl)porphyrin **79** (186 mg, 0.25 mmol, 40 %) as purple crystals, the second fraction gave 5-bromo-10-hexyl-20-(2,4,6-trimethoxyphenyl)porphyrin **78** (194 mg, 0.3 mmol, 38 %) as purple crystals. Data for **79**: M.p. > 300 °C. R_f = 0.6 (*n*-hexane/ CH_2Cl_2 , 1:1, v/v). 1H NMR (400 MHz, $CDCl_3$, TMS): δ = -2.78 (s, 2H, NH), 0.95 (t, 3H, J = 7.01 Hz, $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.42 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.51 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.79 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 2.47 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 3.56 (s, 6H, OCH₃), 4.14 (s, 3H, *p*-OCH₃), 4.8 (t, 2H, J = 8.19 Hz, $CH_2CH_2CH_2CH_2CH_2CH_3$), 6.61 (s, 2H, Ar-*H*), 8.80 (d, 2H, J = 4.1 Hz, β -pyrrole-*H*), 9.39 (d, 2H, J = 4.68 Hz, β -pyrrole-*H*), 9.55 (d, 2H, J = 4.68 Hz, β -pyrrole-*H*), 9.64 ppm (d, 2H, J = 4.71 Hz, β -pyrrole-*H*). ^{13}C NMR (63 MHz, $CDCl_3$): δ = 13.9, 22.5, 29.9, 31.6, 34.9, 38.5, 55.5, 55.8, 90.7, 102.4, 11.6, 112.6, 121.5, 130.0, 131.0, 134.3, 160.9, 162 ppm. UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 421 (5.21), 522

(4.03), 558 (3.85), 603 (3.75), 662 nm (3.81). HRMS (ES^+) calcd. for $C_{35}H_{35}N_4O_3Br_2$ $[M+H]^+$ 719.0854; found 719.0828.

5-Bromo-10-hexyl-20-(3,4,5-trimethoxyphenyl)porphyrin (80):

The compound was derived during the preparation of **81**. M.p. > 300 °C. R_f = 0.3 (*n*-hexane/ CH_2Cl_2 , 1:2, v/v). 1H NMR (400 MHz, $CDCl_3$, TMS): δ = -2.91 (s, 2H, NH), 0.97 (t, 3H, J = 7.02 Hz, $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.43 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.54 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.86 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 2.56 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 4.01 (s, 6H, OCH₃), 4.21 (s, 3H, *p*-OCH₃), 5.01 (t, 2H, J = 8.18 Hz, $CH_2CH_2CH_2CH_2CH_2CH_3$), 7.46 (s, 2H, Ar-*H*), 9.03 (t, 2H, J = 4.68 Hz, β -pyrrole-*H*), 9.28 (d, 1H, J = 4.68 Hz, β -pyrrole-*H*), 9.37 (d, 1H, J = 4.68 Hz, β -pyrrole-*H*), 9.58 (d, 2H, J = 5.26 Hz, β -pyrrole-*H*), 9.74 (d, 1H, J = 4.67 Hz, β -pyrrole-*H*), 9.85 (d, 1H, J = 4.67 Hz, β -pyrrole-*H*), 10.12 ppm (s, 1H, meso-*H*). ^{13}C NMR (63 MHz, $CDCl_3$): δ = 14.1, 22.7, 30.2, 31.9, 35.2, 38.9, 56.4, 61.2, 103.3, 105.1, 112.9, 119.1, 121.1, 136.8, 138.0, 151.5 ppm. UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 416 (5.10), 513 (3.98), 547 (3.50), 592 (3.65), 646 nm (3.40). HRMS (ES^+) calcd. for $C_{35}H_{36}N_4O_3Br$ $[M+H]^+$ 639.1971; found 639.1990.

5,15-Dibromo-10-hexyl-20-(3,4,5-trimethoxyphenyl)porphyrin (81):

Following the general procedure for bromination, 5-hexyl-15-(3,4,5-trimethoxyphenyl)porphyrin **31** (200 mg, 0.35 mmol), NBS (76.2 mg, 0.42 mmol), acetone (10 mL) were used. Column chromatography eluting with *n*-hexane/ CH_2Cl_2 = 2:1 (v/v) gave 5,15-dibromo-10-hexyl-20-(3,4,5-trimethoxyphenyl)porphyrin **81** (100 mg, 0.13 mmol, 38 %) as purple crystals from the first fraction. The second fraction gave purple crystals of 5-bromo-10-hexyl-20-(3,4,5-trimethoxyphenyl)porphyrin **80** (120 mg, 0.18 mmol, 53 %) as purple crystals. Data for **81**: M.p. > 300 °C. R_f = 0.37 (*n*-hexane/ CH_2Cl_2 , 1:8, v/v). 1H NMR (400 MHz, $CDCl_3$, TMS): δ = -2.71 (s, 2H, NH), 0.97 (t, 3H, J = 7.02 Hz, $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.42 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.53 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 1.82 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 2.52 (m, 2H, $CH_2CH_2CH_2CH_2CH_2CH_3$), 4.00 (s, 6H, OCH₃), 4.20 (s, 3H, *p*-OCH₃), 4.92 (t, 2H, J = 8.19 Hz, $CH_2CH_2CH_2CH_2CH_2CH_3$), 7.41 (s, 2H, Ar-*H*), 8.92 (d, 2H, J = 3.51 Hz, β -pyrrole-*H*), 9.48 (d, 2H, J = 4.09 Hz, β -pyrrole-*H*), 9.61 (d, 2H, J = 4.68 Hz, β -pyrrole-*H*), 9.72 ppm (d, 2H, J = 4.68 Hz, β -pyrrole-*H*). ^{13}C NMR (63 MHz, $CDCl_3$): δ = 13.7, 22.3, 29.8, 31.4, 35.1, 38.56, 55.9, 60.8, 102.9, 112.3, 119.8, 121.9, 136.3, 137.4, 151.0 ppm; UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 422 (4.93), 523 (4.15), 558 (3.85), 603 (3.55), 661 nm (3.70). HRMS (ES^+) calcd. for $C_{35}H_{35}N_4O_3Br_2$ $[M+H]^+$ 719.1055; found 719.1050.

General procedure for Suzuki coupling

A Schlenk flask was charged with K_3PO_4 (20 equiv.) and anhydrous THF (60 mL) under an argon atmosphere. Then the porphyrin (1 equiv.), arylboronic acid or arylboronic ester (10 equiv.) and $Pd(PPh_3)_4$ (0.1 equiv.) were added. The reaction was heated at reflux temperature for 12-24 hours

(TLC control) and protected from light. After completion, the solvent was removed under reduced pressure and the residue was dissolved in CH₂Cl₂. The mixture was washed with saturated NaHCO₃, H₂O, and brine and then dried over Na₂SO₄. The organic solvent was evaporated and the crude product was purified by flash chromatography.

5-Hexyl-10-(3-nitrophenyl)-15-(2,4,6-trimethoxyphenyl)porphyrin (83):

Following the above general procedure, 5-bromo-10-hexyl-20-(2,4,6-trimethoxyphenyl)porphyrin **78** (99 mg, 0.1544 mmol), 3-nitrophenyl boronic acid (257 mg, 1.54 mmol), Pd(PPh₃)₄ (17.79 mg, 0.0154 mmol) and K₃PO₄ (657.14 mg, 2.89 mmol) in THF (50 mL) were used. Column chromatography on silica gel using *n*-hexane/ethyl acetate (5:1, v/v) gave the title compound as purple crystals (46 mg, 0.067 mmol, 46 %). M.p. > 300 °C. *R*_f = 0.25 (*n*-hexane/ethyl acetate, 5:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.84 (*s*, 2H, NH), 0.98 (*t*, 3H, *J* = 7.02 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 1.43 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.55 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.81 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.55 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 3.58 (*s*, 6H, OCH₃), 4.15 (*s*, 3H, *p*-OCH₃), 4.95 (*t*, 2H, *J* = 7.6 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 6.66 (*s*, 2H, Ar-*H*), 7.85 (*t*, 1H, *J* = 7.6 Hz, Ar-*H*), 8.46 (*d*, 1H, *J* = 7.6 Hz, Ar-*H*), 8.68 (*d*, 2H, *J* = 4.68 Hz, β-pyrrole-*H*), 8.76 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 8.94 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.02 (*d*, 1H, *J* = 4.1 Hz, Ar-*H*), 9.13 (*s*, 1H, Ar-*H*), 9.31 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.35 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.49 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.55 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 10.11 ppm (*s*, 1H, meso-*H*). ¹³C NMR (100 MHz, CDCl₃): δ = 13.7, 22.3, 29.3, 29.8, 31.4, 34.7, 38.3, 55.2, 55.5, 90.5, 104.0, 115.0, 120.1, 122.2, 136.8, 127.3, 139.1, 144.3, 146.2, 160.6, 161.6 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ε) = 414 (5.17), 510 (4.13), 542 (3.53), 581 (3.83), 639 nm (3.43). HRMS (ES⁺) calcd. for C₄₁H₄₀N₅O₅ [M+H]⁺ 682.3029; found 682.3044.

5-Hexyl-10-(2-phenylethenyl)-15-(2,4,6-trimethoxyphenyl)porphyrin (84):

Following the above general procedure, 5-bromo-10-hexyl-20-(2,4,6-trimethoxyphenyl)porphyrin **27** (95 mg, 0.148 mmol), *trans*-betastylene boronic acid (218.89 mg, 1.48 mmol), Pd(PPh₃)₄ (17.1 mg, 0.0148 mmol) and K₃PO₄ (630 mg, 2.97 mmol) in THF (50 mL) were used. Column chromatography on silica gel (*n*-hexane/ethyl acetate = 10:1, v/v) gave the desired product as purple crystals (43 mg, 0.064 mmol, 43 %). M.p. > 300 °C. *R*_f = 0.52 (*n*-hexane/ethyl acetate, 5:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.66 (*s*, 2H, NH), 0.97 (*t*, 3H, *J* = 7.02 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 1.43 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.55 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.84 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.57 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 3.56 (*s*, 6H, OCH₃), 4.14 (*s*, 3H, *p*-OCH₃), 4.99 (*t*, 2H, *J* = 7.6 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 6.64 (*s*, 2H, Ar-*H*), 7.38 (*d*, 1H, *J* = 15.78 Hz, CH=CH), 7.50 (*t*, 1H, *J* = 7.6 Hz, Ar-*H*), 7.62 (*t*, 2H, *J* = 7.6 Hz, Ar-*H*), 7.98 (*d*, 2H, *J* = 7.6 Hz, Ar-*H*), 8.92 (*m*, 2H, β-pyrrole-*H*), 9.22 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.31 (*d*, 1H, *J* = 4.67 Hz, β-pyrrole-*H*), 9.51 (*d*, 2H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.35 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.59 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.70 (*d*,

1H, *J* = 15.79 Hz, CH=CH), 10.0 ppm (*s*, 1H, meso-*H*). ¹³C NMR (100 MHz, CDCl₃): δ = 13.7, 22.3, 29.8, 31.5, 34.8, 38.2, 55.2, 55.5, 90.5, 103.3, 110.5, 111.7, 115.5, 119.7, 126.4, 127.6, 128.6, 129.4, 137.6, 142.0, 60.7, 161.5 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ε) = 422 (4.99), 516 (3.99), 556 (3.82), 599 (3.52), 653 nm (3.42). HRMS (ES⁺) calcd. for C₄₃H₄₂N₄O₃ [M+H]⁺ 663.3335; found 663.3340.

5-Hexyl-10-(3-nitrophenyl)-15-(3,4,5-trimethoxyphenyl)porphyrin (85):

Following the above general procedure, 5-bromo-10-hexyl-20-(3,4,5-trimethoxyphenyl)porphyrin **80** (60 mg, 0.093 mmol), 3-nitrophenyl boronic acid (156.58 mg, 0.938 mmol), Pd(PPh₃)₄ (10.83 mg, 0.0093 mmol) and K₃PO₄ (398.31 mg, 1.87 mmol) in THF (50 mL) were used. Column chromatography on silica gel (*n*-hexane/CH₂Cl₂ = 2:1, v/v) gave the title compound as purple crystals (53 mg, 0.077 mmol, 82 %). M.p. > 300 °C. *R*_f = 0.3 (*n*-hexane/ethyl acetate, 10:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.91 (*s*, 2H, NH), 0.96 (*t*, 3H, *J* = 7.0 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 1.43 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.55 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.86 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.59 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 4.01 (*s*, 6H, OCH₃), 4.22 (*s*, 3H, *p*-OCH₃), 5.06 (*t*, 2H, *J* = 7.61 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 7.49 (*s*, 2H, Ar-*H*), 7.97 (*t*, 1H, *J* = 8.18 Hz, Ar-*H*), 8.56 (*d*, 1H, *J* = 7.6 Hz, Ar-*H*), 8.72 (*d*, 2H, *J* = 5.27 Hz, Ar-*H*), 8.83 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.02 (*m*, 1H, β-pyrrole-*H*), 9.11 (*m*, 2H, Ar-*H*), 9.36 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.44 (*d*, 1H, *J* = 4.04 Hz, β-pyrrole-*H*), 9.56 (*d*, 1H, *J* = 5.26 Hz, β-pyrrole-*H*), 9.65 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 10.21 ppm (*s*, 1H, meso-*H*). ¹³C NMR (100 MHz, CDCl₃): δ = 22.2, 29.8, 31.4, 34.7, 38.4, 55.9, 60.86, 104.7, 112.4, 115.7, 118.5, 120.5, 122.4, 127.0, 128.3, 139.1, 144.1, 146.3, 151.1 ppm. UV/Vis (CH₂Cl₂): λ_{max} (log ε) = 415 (4.97), 511 (4.00), 547 (3.83), 588 (3.91), 640 nm (3.67). HRMS (ES⁺) calcd. for C₄₁H₄₀N₅O₅ [M+H]⁺ 682.3029; found 682.3021.

5-Hexyl-10-(4-methoxycarbonylphenyl)-15-(3,4,5-trimethoxyphenyl)porphyrin (86):

Following the above general procedure, 5-bromo-10-hexyl-20-(3,4,5-trimethoxyphenyl)porphyrin **80** (60 mg, 0.0938 mmol), 4-methoxycarbonylphenyl boronic acid (168.8 mg, 0.938 mmol), Pd(PPh₃)₄ (10.83 mg, 0.00938 mmol) and K₃PO₄ (398.21 mg, 1.876 mmol) in THF (50 mL) were used. Column chromatography on silica gel (*n*-hexane/CH₂Cl₂ = 2:1, v/v) gave the title compound as purple crystals (57 mg, 0.083 mmol, 87 %). M.p. > 300 °C. *R*_f = 0.37 (*n*-hexane/CH₂Cl₂, 1:2, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.90 (*s*, 2H, NH), 0.97 (*t*, 3H, *J* = 7.02 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 1.43 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.54 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.86 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.58 (*m*, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 4.00 (*s*, 6H, OCH₃), 4.16 (*s*, 3H, OCH₃), 4.21 (*s*, 3H, -OCH₃), 5.05 (*t*, 2H, *J* = 8.19 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 7.49 (*s*, 2H, Ar-*H*), 8.32 (*d*, 2H, *J* = 8.18 Hz, Ar-*H*), 8.48 (*d*, 2H, *J* = 8.19 Hz, Ar-*H*), 8.80 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 8.90 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.00 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.11 (*d*, 1H, *J* = 4.68 Hz, β-pyrrole-*H*), 9.35 (*d*, 1H, *J* = 4.68 Hz,

β -pyrrole-*H*), 9.43 (*d*, 1H, $J = 4.09$ Hz, β -pyrrole-*H*), 9.54 (*d*, 1H, $J = 4.67$ Hz, β -pyrrole-*H*), 9.64 (*d*, 1H, $J = 4.68$ Hz, β -pyrrole-*H*), 10.19 ppm (*s*, 1H, meso-*H*). ^{13}C NMR (150 MHz, CDCl_3): $\delta = 13.7, 22.2, 22.3, 28.9, 29.2, 29.8, 31.4, 34.8, 52.0, 55.9, 60.8, 104.3, 112.3, 117.9, 118.3, 120.2, 126.7, 127.2, 129.1, 129.7, 130.6, 130.9, 133.9, 136.6, 137.3, 147.2, 151.0, 166.9$ ppm. UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 414 (5.18), 510 (4.14), 544 (3.84), 591 (3.92), 639 nm (3.74). HRMS (ES^+) calcd. for $\text{C}_{43}\text{H}_{43}\text{N}_4\text{O}_5$ $[\text{M}+\text{H}]^+$ 695.3233; found 695.3214.

5,15-Dihexyl-10-phenyl-20-propylporphyrin (100): The reaction followed the procedure given for **99** using 0.8 ml *n*-propyl iodide (8.1 mmol) and gave the title compound (20 mg, 0.03 mmol, 21 %) as purple crystals and trisubstituted porphyrin (24 %). Data for **100**: M.p. >300 °C. $R_f = 0.47$ (ethyl acetate/*n*-hexane, 1:7, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): $\delta = -2.63$ (*s*, 2H, *NH*), 0.92 (*t*, 6H, $J = 7.2$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.25 (*m*, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.37-1.51 (*m*, 7H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_3$), 1.86 (*m*, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.58 (*m*, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 3.75 (*t*, 2H, $J = 6.4$, $\text{CH}_2\text{CH}_2\text{CH}_3$), 4.99 (*m*, 6H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_3$), 7.77 (*m*, 3H, *Ph-H*), 8.18 (*m*, 2H, *Ph-H*), 8.82 (*d*, 2H, $J = 5$ Hz, β -pyrrole- $H_{8,12}$), 9.39 (*d*, 2H, $J = 5$ Hz, β -pyrrole- $H_{2,18}$), 9.52 (*d*, 2H, $J = 5$ Hz, β -pyrrole- $H_{3,17}$), 9.56 ppm (*d*, 2H, $J = 5$ Hz, β -pyrrole- $H_{7,13}$). UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 440 (5.03), 486 (3.92), 577 (3.77), 594 (3.77), 659 nm (3.37). HRMS calcd. for $\text{C}_{41}\text{H}_{48}\text{N}_4$ 596.3878; found $[\text{M} + 1]$ 596.3953. $\text{C}_{41}\text{H}_{48}\text{N}_4$ (596.86): calcd. C 82.51, H 8.11, N 9.39; found C 82.33, H 8.25, N 9.11.

5-Hexyl-10-pentyl-15-(4-methoxyphenyl)-20-phenylporphyrin (106): The reaction was performed following the method described for **105** using phenyllithium (0.8 ml of a 1.8 M solution in hexane, 0.02 mmol), 5-hexyl-15-(4-methoxyphenyl)porphyrin **51** (100 mg, 0.19 mmol) and 0.8 ml pentyl iodide (6.1 mmol) and gave the title compound (18 mg, 0.027 mmol, 14 %) as purple crystals, trisubstituted porphyrin (9 %) and starting material (6 %). M.p. >300 °C. $R_f = 0.75$ (ethyl acetate/*n*-hexane, 1:5, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): $\delta = -2.64$ (*s*, 2H, *NH*), 0.98 (*t*, 6H, $J = 8.1$ Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.43 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.58 (*m*, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.87 (*m*, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.59 (*m*, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 4.09 (*s*, 3H, OCH_3), 5.03 (*t*, 4H, $J = 8.1$ Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 7.31 (*d*, 2H, $J = 8$ Hz, *Ph-H*), 7.77 (*m*, 3H, *Ph-H*), 8.13 (*d*, 2H, $J = 8$ Hz, *Ph-H*), 8.23 (*m*, 2H, *Ph-H*), 8.78 (*m*, 2H, β -pyrrole- $H_{13,17}$), 8.92 (*m*, 2H, β -pyrrole- $H_{2,18}$), 9.46 (*m*, 2H, β -pyrrole- $H_{3,12}$), 9.58 ppm (*m*, 2H, β -pyrrole- $H_{7,8}$). UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 418 (5.09), 485 (4.00), 517 (3.91), 553 (3.64), 593 (3.66), 655 nm (4.23). HRMS: calcd. for $\text{C}_{44}\text{H}_{46}\text{N}_4\text{O}$ 646.3671; found $[\text{M} + 1]$ 647.3771. $\text{C}_{44}\text{H}_{46}\text{N}_4\text{O}$ (646.88): calcd. C 81.70, H 7.17, N 8.66; found C 81.43, H 7.35, N 8.88.

5-Butyl-10-hexyl-15-propyl-20-(4-methoxyphenyl)porphyrin (108): The reaction was

performed following the method described for **107** using *n*-BuLi (0.4 ml of 2.5 M solution in hexane, 1 mmol), 5-hexyl-15-(4-methoxyphenyl)porphyrin **51** (100 mg, 0.19 mmol), and 0.7 ml *n*-propyl iodide (7.1 mmol). Final purification yielded the title compound (22 mg, 0.036 mmol, 18 %) as purple crystals, trisubstituted porphyrin (10 %) and starting material (8 %). M.p. >300 °C. $R_f = 0.62$ (ethyl acetate/*n*-hexane, 1:5, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): $\delta = -2.63$ (*s*, 2H, *NH*), 0.98 (*m*, 6H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_3$), 1.33-1.37 (*m*, 5H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.56 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.87 (*m*, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.59 (*m*, 6H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_3$), 4.13 (*s*, 3H, OCH_3), 5.01 (*m*, 6H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 7.28 (*d*, 2H, $J = 8$ Hz, *Ar-H*), 8.09 (*d*, 2H, $J = 8$ Hz, *Ar-H*), 8.86 (*d*, 2H, $J = 5$ Hz, β -pyrrole-*H*), 9.42 (*d*, 2H, $J = 5$ Hz, β -pyrrole-*H*), 9.52-9.56 ppm (*m*, 4H, β -pyrrole-*H*). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 13.79, 14.52, 22.36, 23.22, 28.49, 29.29, 29.91, 31.12, 31.52, 34.70, 35.37, 36.84, 38.45, 40.34, 50.97, 55.14, 111.61, 117.43, 118.25, 118.57, 118.80, 126.04, 128.38, 130.49, 134.55, 134.94, 158.79$ ppm. UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 418 (5.00), 487 (4.60), 517 (3.33), 555 (2.85), 598 (2.77), 655 nm (4.17). HRMS: calcd. for $\text{C}_{40}\text{H}_{46}\text{N}_4\text{O}$ 598.3671; found $[\text{M} + 1]$ 599.3739. $\text{C}_{40}\text{H}_{46}\text{N}_4\text{O}$ (598.83): calcd. C 80.22, H 7.75, N 9.36; found C 80.43, H 7.56, N 9.05.

5-(*sec*-Butyl)-10-hexyl-20-(4-methoxyphenyl)porphyrin

(111): 5-Hexyl-15-(4-methoxyphenyl)porphyrin **51** (100 mg, 0.19 mmol) was reacted with *sec*-butyllithium (1 ml of a 2.5 M solution in hexane, 2.5 mmol) similar to the procedure given for **110**. The title compound was obtained as purple crystals (15 mg, 0.026 mmol, 14 %) accompanied by 10 % recovered starting material. M.p. >300 °C. $R_f = 0.62$ (ethyl acetate/*n*-hexane, 1:5, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): $\delta = -2.92$ (*s*, 2H, *NH*), 0.89 (*t*, 3H, $J = 7.6$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.10 (*t*, 3H, $J = 7.6$ Hz, $\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$), 1.42 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.57 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.89 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.46 (*d*, 3H, $J = 7.4$ Hz, $\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$), 2.60 (*m*, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 3.02 (*m*, 2H, $\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$), 4.13 (*s*, 3H, OCH_3), 5.03 (*t*, 2H, $J = 7.6$ Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 5.46 (*m*, 1H, $\text{CH}(\text{CH}_3)(\text{CH}_2\text{CH}_3)$), 7.33 (*d*, 2H, $J = 8$ Hz, *Ar-H*), 8.12 (*d*, 2H, $J = 8$ Hz, *Ar-H*), 8.93 (*d*, 2H, $J = 5$ Hz, β -pyrrole- $H_{2,18}$), 9.23-9.33 (*d*, 2H, $J = 5$ Hz, β -pyrrole- $H_{13,17}$), 9.58 (*d*, 2H, $J = 5$ Hz, β -pyrrole- $H_{8,12}$), 9.66-9.78 (*d*, 2H, $J = 5$ Hz, β -pyrrole- $H_{3,7}$), 10.01 ppm (*s*, 1H, meso-*H*). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 13.75, 13.83, 22.35, 27.02, 29.28, 31.51, 35.18, 35.49, 38.43, 42.60, 55.15, 102.96, 111.74, 117.56, 118.88, 124.84, 135.06, 158.87$ ppm. UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 440 (4.96), 486 (3.61), 585 (3.63), 656 nm (3.94). HRMS calcd. for $\text{C}_{37}\text{H}_{40}\text{N}_4\text{O}$ 556.3202; found 556.3239. $\text{C}_{37}\text{H}_{40}\text{N}_4\text{O}$ (556.75): calcd. C 79.82, H 7.24, N 10.06; found C 80.08, H 7.37, N 9.88.

5-Butyl-10-(3,5-dimethoxyphenyl)-20-hexylporphyrin

(113): 5-Hexyl-15-(3,5-dimethoxyphenyl)porphyrin **112** (100 mg, 0.18 mmol) was reacted with *n*-butyllithium (0.4 ml of 2.5 M solution in hexane, 1 mmol) similar to the reaction described for **110**. The title compound was obtained

as purple crystals (16 mg, 0.027 mmol, 15 %) and 12 % of starting material was recovered. M.p. >300 °C. R_f = 0.60 (ethyl acetate/*n*-hexane, 1:5, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): δ = -2.91 (s, 2H, NH), 0.93 (t, 3H, J = 7.5 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.16 (t, 3H, J = 7.5 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.26 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.47 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.88 (m, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.54 (m, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 4.00 (s, 6H, OCH_3), 5.04 (t, 4H, J = 7.8 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 6.94 (m, 1H, Ar-*H*), 7.42 (m, 2H, Ar-*H*), 9.03 (2d, 2H, J = 5 Hz, β -pyrrole- $H_{13,17}$), 9.23-9.34 (2d, 2H, J = 5 Hz, β -pyrrole- $H_{13,17}$), 9.52-9.65 (m, 4H, β -pyrrole- $H_{2,3,7,18}$), 10.14 ppm (s, 1H, meso-*H*). ^{13}C NMR (100 MHz, CDCl_3): δ = 13.74, 13.80, 22.33, 25.17, 29.27, 29.89, 30.51, 31.50, 38.40, 40.70, 55.19, 99.51, 103.07, 117.31, 120.08, 127.71, 130.45, 143.45, 158.47 ppm. UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 436 (5.02), 486 (3.11), 544 (2.86), 581 (3.50), 656 nm (3.72). HRMS calcd. for $\text{C}_{38}\text{H}_{42}\text{N}_4\text{O}_2$ 586.3307; found [M + 1] 586.5437. $\text{C}_{38}\text{H}_{42}\text{N}_4\text{O}_2$ (586.78): calcd. C 77.78, H 7.21, N 9.55; found C 77.41, H 7.42, N 9.21.

5-(3,5-Dimethoxyphenyl)-10-(4-hydroxyphenyl)-15-hexyl-20-phenyl-porphyrin (117): The reaction was performed similar to **115** using 3,5-dimethoxy-*p*-phenol (0.43 g, 2.5 mmol) and 5-(3,5-dimethoxyphenyl)-10-phenyl-15-hexylporphyrin **116** (40 mg, 0.07 mmol). Final purification was achieved by column chromatography and elution with ethyl acetate/*n*-hexane (1:5, v/v) yielded the title compound (17 mg, 0.024 mmol, 37 %) as purple crystals and 10 % recovered starting material. M.p. >300 °C. R_f = 0.28 (ethyl acetate/*n*-hexane, 1:1, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): δ = -2.74 (s, 2H, NH), 0.92 (t, 3H, J = 8.1 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.41 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.53 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.84 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.59 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 3.97 (s, 6H, OCH_3), 5.04 (t, 2H, J = 8.1 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 6.77 (m, 4H, Ph-*H*), 7.22-7.39 (m, 4H, Ph-*H*), 7.79 (m, 2H, Ph-*H*), 8.07 (d, 1H, J = 8 Hz, Ph-*H*), 8.21 (d, 1H, J = 6.4 Hz, Ph-*H*), 8.79-8.83 (2d, 2H, J = 5 Hz, β -pyrrole- $H_{3,7}$), 8.90-8.97 (m, 4H, β -pyrrole- $H_{2,8,12,18}$), 9.50 ppm (d, 2H, J = 5 Hz, β -pyrrole- $H_{13,17}$). UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 449 (4.04), 514 (3.66), 552 (3.60), 585 (3.60), 656 nm (4.34). HRMS calcd. for $\text{C}_{46}\text{H}_{42}\text{N}_4\text{O}_3$ 698.3256; found [M + 1] 699.3335. $\text{C}_{46}\text{H}_{42}\text{N}_4\text{O}_3$ (698.86): calcd. C 79.06, H 6.06, N 8.02; found C 79.17, H 5.99, N 8.13.

5-(4-Methoxyphenyl)-10-(4-methylphenyl)-15-phenyl-20-(2,4,6-trimethoxyphenyl)porphyrin (118): The synthesis followed the procedure given for **61** using 4-methoxyphenyllithium and 100 mg of **62** (0.16 mmol). Column chromatography eluting with dichloromethane:*n*-hexane (1:2, v/v) gave a small amount of starting material followed by the title compound as a purple solid (47 mg, 64 μmol , 40 %). M.p. >320 °C. ^1H NMR (500 MHz, CDCl_3 , TMS): δ = 8.945 (m, 8H, β -pyrrole-*H*), 8.24 (m, 2H, *p*-MeOPh-*H*), 8.12 (m, 4H, Ar-*H*), 7.73 (m, 3H, Ph- $H_{m,p}$), 7.54 (m, 2H, tolyl- H_m), 7.20 (d, 2H, MeOPh- H_m), 6.56 (s, 2H, triMeOPh- H_m), 4.03 (s, 3H, triMeOC $_6$ H $_2$ -*p*CH $_3$), 3.98 (s, 3H, C $_6$ H $_4$ -OCH $_3$), 3.53 (s, 6H, triMeOC $_6$ H $_2$ -*o*CH $_3$), 2.71 (s, 3H, C $_6$ H $_4$ -CH $_3$), -3.04 ppm (*br s*, 2H, NH). UV/vis (CH_2Cl_2):

λ_{max} (log ϵ) = 410 (5.25), 508 (4.20), 543 (3.67), 584 (3.65), 640 nm (3.33). MS (FAB $^+$, 3 kV): m/z 750.0 (100.00 %, [M + H] $^+$). HRMS calcd. for $\text{C}_{49}\text{H}_{40}\text{N}_4\text{O}_4$ 748.30496; found 748.30410. $\text{C}_{49}\text{H}_{40}\text{N}_4\text{O}_4$ (748.88): calcd. C 78.59, H 5.38, N 7.48; found C 78.65, H 5.71, N 7.73.

5-Hexyl-10-(4-methylphenyl)-15-phenyl-20-(2,4,6-trimethoxyphenyl)porphyrin (119): The synthesis followed the procedure given for **62** using hexyllithium and 100 mg of **62** (0.16 mmol). Column chromatography eluting with dichloromethane:*n*-hexane (1:2, v/v) gave the title compound as the sole product as a purple solid (80 mg, 0.11 mmol, 70 %). M.p. >300 °C. ^1H NMR (500 MHz, CDCl_3 , TMS): δ = 9.46 (d, 2H, 3J = 4.54 Hz, β -pyrrole-*H*), 8.94 (m, 2H, β -pyrrole-*H*), 8.82 (d, 2H, 3J = 4.54 Hz, β -pyrrole-*H*), 8.78 (d, 2H, 3J = 4.54 Hz, β -pyrrole-*H*), 8.21 (d, 2H, 3J = 7.63 Hz, tolyl- H_o), 8.12 (d, 2H, 3J = 7.63 Hz), 7.73 (m, 3H, Ph- $H_{m,p}$), 7.56 (d, 2H, 3J = 7.45 Hz, tolyl- H_m), 6.62 (s, 2H, Ar- H_m), 5.00 (t, 2H, 3J = 7.9 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 4.11 (s, 3H, *p*-OCH $_3$), 3.53 (s, 6H, *o*-OCH $_3$), 2.73 (s, 3H, C $_6$ H $_4$ -CH $_3$), 2.60 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.85 (q, 2H, 3J = 7.36 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.54 (q, 2H, 3J = 6.99 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.44 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 0.97 (t, 3H, 3J = 7.30 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), -3.06 ppm (*b s*, 2H, NH). ^{13}C NMR (100 MHz, CDCl_3): δ = 162.00, 161.12, 148.41-145.14, 142.37, 139.65, 137.13, 134.51, 134.42, 131.02-130.50, 127.41, 127.25, 126.60, 119.12-118.57, 102.88, 90.99, 56.06, 55.63, 38.81, 35.51, 32.00, 30.34, 22.80, 21.51, 14.15 ppm. UV/vis (CH_2Cl_2): λ_{max} (log ϵ) = 413 (5.24), 508 (4.15), 542 (3.65), 584 (3.63), 638 nm (3.23). MS (FAB $^+$, 3 kV): m/z 728.0 (100.00 %, [M + H] $^+$). HRMS calcd. for $\text{C}_{48}\text{H}_{46}\text{N}_4\text{O}_3$ 726.356; found 726.3561.

Chloro[5-hexyl-10-(4-methylphenyl)-15-phenyl-20-(2,4,6-trimethoxyphenyl)porphyrinato]manganese(III) (120): The free base porphyrin **119** (100 mg, 0.13 mmol) was treated as described in the procedure for **65**. Column chromatography eluting with dichloromethane gave several minor fractions and the metal complex as the main fraction. Evaporation of the solvent gave a green-red residue (98 mg, 0.12 mmol, 90 %). M.p. >300 °C. UV/Vis (CH_2Cl_2): λ_{max} (log ϵ) = 375 (4.69), 402 (4.60), 479 (4.95), 528 (3.7), 583 (3.95), 619 nm (4.01). MS (FAB $^+$, 3 kV): m/z = calcd. for $\text{C}_{48}\text{H}_{44}\text{ClMnN}_4\text{O}_3$ 814.24824; found 814.0 (11.0 %, [M] $^{+}$), 799.0 (100.0 %, [M - Cl] $^{+}$).

5-Bromo-10-hexyl-15-(3-nitrophenyl)-20-(3,4,5-trimethoxyphenyl)porphyrin (123): The synthesis followed the standard procedure given for **77** using 5-hexyl-10-(3-nitrophenyl)-15-(3,4,5-trimethoxyphenyl)porphyrin **85** (90 mg, 0.132 mmol), NBS (25.8 mg, 0.14 mmol) and acetone (10 mL). The mixture was filtered through a short silica gel column and gave the title compound as purple crystals (77 mg, 0.10 mmol, 71 %). M.p. >300 °C. R_f = 0.3 (CH_2Cl_2 /*n*-hexane, 2:1, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): δ = -3.60 (s, 2H, NH), 0.96 (t, 2H, J = 7.01 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.49 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.53 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.84 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.56 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 4.0 (s, 6H, OCH $_3$), 4.2 (s, 3H, *p*-OCH $_3$), 5.00 (t, 2H, J = 8.18, Hz,

CH₂CH₂CH₂CH₂CH₂CH₃), 7.43 (s, 2H, Ar-H), 7.98 (t, 2H, *J* = 8.18 Hz, Ar-H), 8.52 (d, 1H, *J* = 7.6 Hz, Ar-H), 8.65 (d, 1H, *J* = 5.27 Hz, β-pyrrole-H), 8.71 (d, 1H, *J* = 8.19 Hz, Ar-H), 8.76 (d, 1H, *J* = 4.68 Hz, β-pyrrole-H), 8.91 (d, 1H, *J* = 4.67 Hz, β-pyrrole-H), 8.99 (d, 1H, *J* = 4.68 Hz, β-pyrrole-H), 9.06 (s, 1H, Ar-H), 9.50 (d, 1H, *J* = 5.26 Hz, β-pyrrole-H), 9.58 (d, 1H, *J* = 4.67 Hz, β-pyrrole-H), 9.69 (d, 1H, *J* = 4.68 Hz, β-pyrrole-H), 9.81 ppm (d, 1H, *J* = 4.67 Hz, β-pyrrole-H). ¹³C NMR (100 MHz, CDCl₃): δ = 14.1, 22.7, 30.2, 31.9, 35.2, 38.9, 56.4, 61.2, 103.3, 105.1, 112.9, 119.1, 121.1, 136.8, 138.0, 151.1 ppm. UV/vis (CH₂Cl₂): λ_{max} (log ε) = 422 (5.05), 519 (4.05), 556 (3.88), 599 (3.35), 653 nm (3.48). HRMS (ES⁺) calcd. for C₄₁H₃₇N₄O₅Br [M+H]⁺ 761.2213; found 761.2216.

5-Bromo-10-hexyl-15-(sec-butyl)-20-(3,4,5-trimethoxyphenyl)porphyrin (124): The synthesis followed the standard procedure given for **77** using 5-(sec-butyl)-10-hexyl-20-(3,4,5-trimethoxyphenyl)porphyrin **69**, (20 mg, 0.0324 mmol) NBS (7.5 mg, 0.042 mmol) and acetone (5 mL). Column chromatography (*n*-hexane/CH₂Cl₂ = 1:2, v/v) gave the title compound as purple crystals (16 mg, 0.024 mmol, 77 %). M.p. >300 °C. *R_f* = 0.28 (CH₂Cl₂/*n*-hexane, 2:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.60 (s, 2H, NH), 0.98 (t, 3H, *J* = 7.6 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 1.08 (t, 3H, *J* = 7.01 Hz, CHCH₂CH₃), 1.45 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.56 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.86 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.42 (d, 3H, *J* = 7.01 Hz, CHCH₃), 2.56 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.81 (m, 1H, CHCH₂CH₃), 2.95 (m, 1H, CHCH₂CH₃), 4.03 (s, 6H, OCH₃), 4.21 (s, 3H, *p*-OCH₃), 4.97 (t, 2H, *J* = 8.18 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 5.34 (m, 1H, CHCH₂CH₃), 7.42 (s, 2H, Ar-H), 8.90 (m, 2H, β-pyrrole-H), 9.52 (m, 2H, β-pyrrole-H), 9.55 (m, 2H, β-pyrrole-H), 9.68 ppm (d, 2H, *J* = 4.67 Hz, β-pyrrole-H). ¹³C NMR (100 MHz, CDCl₃): δ = 13.7, 13.7, 22.3, 26.8, 29.2, 29.8, 31.4, 35.2, 38.6, 42.3, 55.9, 60.8, 100.9, 112.1, 118.4, 120.4, 125.4, 137.2, 137.3 ppm. UV/vis (CH₂Cl₂): λ_{max} (log ε) = 423 (4.91), 522 (4.12), 558 (3.99), 603 (3.72), 660 nm (3.51). HRMS calcd. for C₃₉H₄₃N₄Br [M+H]⁺ 646.2671; found 646.2655.

5-Bromo-10-(*n*-butyl)-15-(4-methoxycarbonylphenyl)-20-(3-bromo-2,4,6-trimethoxyphenyl)porphyrin (125): The synthesis followed the standard procedure given for **77** using 5-(*n*-butyl)-10-(4-methoxycarbonylphenyl)-15-(2,4,6-trimethoxyphenyl)porphyrin **82** (145.2 mg, 0.217 mmol), NBS (46.4 mg, 0.33 mmol) and acetone (10 mL). Column chromatography (*n*-hexane/CH₂Cl₂ = 1:2, v/v) gave the title compound as purple crystals (96 mg, 0.11 mmol, 54 %). M.p. >300 °C. *R_f* = 0.5 (CH₂Cl₂/*n*-hexane, 2:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.63 (s, 2H, NH), 1.14 (t, 3H, *J* = 7.02 Hz, CH₂CH₂CH₂CH₃), 1.81 (m, 2H, CH₂CH₂CH₂CH₃), 2.50 (m, 2H, CH₂CH₂CH₂CH₃), 3.02 (s, 3H, OCH₃), 3.59 (s, 3H, OCH₃), 4.15 (s, 3H, OCH₃), 4.23 (s, 3H, OCH₃), 4.95 (t, 2H, *J* = 7.16 Hz, CH₂CH₂CH₂CH₃), 6.79 (s, 1H, Ar-H), 8.28 (t, 2H, *J* = 7.6 Hz, Ar-H), 8.46 (d, 2H, *J* = 8.18 Hz, Ar-H), 8.71 (d, 1H, *J* = 4.69 Hz, β-pyrrole-H), 8.79 (d, 2H, *J* = 5.27 Hz, β-pyrrole-H), 8.88 (d, 1H, *J* = 4.67 Hz, β-pyrrole-H), 9.43 (d, 1H, *J* = 4.68 Hz, Ar-H), 9.50 (d, 1H, *J* = 4.68 Hz, β-pyrrole-H), 9.65 (d, 1H, *J* = 5.26 Hz, β-pyrrole-H), 9.74 ppm (d, 1H, *J* = 4.67 Hz, β-pyrrole-H). ¹³C NMR (150 MHz, CDCl₃): δ = 13.7, 23.1, 34.8, 40.4, 52.0, 55.8, 56.3, 60.5, 92.0, 98.0, 102.2, 110.1,

118.0, 121.6, 126.7, 127.4, 129.1, 129.7, 133.8, 133.9, 146.5, 157.7, 158.1, 159.6, 166.9 ppm. UV/vis (CH₂Cl₂): λ_{max} (log ε) = 422 (5.36), 520 (4.21), 555 (3.91), 598 (3.61), 653 nm (3.51). HRMS calcd. for C₄₁H₃₇N₄O₅Br₂ [M+H]⁺ 825.110; found 825.1090.

5-Hexyl-10-(3-nitrophenyl)-15-(3,4,5-trimethoxyphenyl)-20-(4-methoxycarbonylphenyl)porphyrin (126): The synthesis followed the standard procedure given for **82** using 5-bromo-10-hexyl-15-(3-nitrophenyl)-20-(3,4,5-trimethoxyphenyl)porphyrin **123** (30.4 mg, 0.04 mmol), 4-methoxycarbonylphenyl boronic acid (89.98 mg, 0.5 mmol), Pd(PPh₃)₄ (5.77 mg, 0.005 mmol) and K₃PO₄ (212.2 mg, 1 mmol) in THF (50 mL). Column chromatography on silica gel (*n*-hexane/ethyl acetate = 10:1, v/v) afforded the desired product (21.9 mg, 0.026 mmol, 67 %) as a purple solid. M.p. >300 °C. *R_f* = 0.5 (*n*-hexane/ethyl acetate, 6:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.71 (s, 2H, NH), 0.96 (t, 3H, *J* = 7.02 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 1.43 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.54 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.85 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.58 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 3.99 (s, 6H, OCH₃), 4.16 (s, 3H, OCH₃), 4.20 (s, 3H, OCH₃), 5.04 (t, 2H, *J* = 7.6 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 7.46 (s, 2H, Ar-H), 7.99 (t, 1H, *J* = 7.6 Hz, Ar-H), 8.33 (d, 2H, *J* = 7.6 Hz, Ar-H), 8.49 (d, 2H, *J* = 7.6 Hz, Ar-H), 8.57 (d, 1H, *J* = 7.02 Hz, Ar-H), 8.73 (m, 2H, Ar-H, β-pyrrole-H), 8.79 (d, 1H, *J* = 4.09 Hz, β-pyrrole-H), 8.83 (d, 1H, *J* = 4.68 Hz, β-pyrrole-H), 8.90 (d, 1H, *J* = 4.68 Hz, β-pyrrole-H), 8.97 (t, 2H, *J* = 4.68 Hz, β-pyrrole-H), 9.11 (m, 1H, Ar-H), 9.55 ppm (m, 2H, β-pyrrole-H). ¹³C NMR (150 MHz, CDCl₃): δ = 13.9, 22.5, 29.2, 29.5, 30.1, 31.7, 35.4, 38.8, 52.3, 56.2, 61.1, 112.7, 127.4, 127.7, 128.1, 129.5, 134.3, 137.0, 139.4, 144.0, 146.8, 146.9, 151.3, 167.1 ppm. UV/vis (CH₂Cl₂): λ_{max} (log ε) = 422 (5.00), 516 (3.91), 553 (3.61), 594 (3.51), 648 nm (3.38). HRMS calcd. for C₄₉H₄₅N₅O₇ [M+H]⁺ 816.3397; found 816.3363.

5-(*n*-Butyl)-10-hexyl-15-(3-nitrophenyl)-20-(3,4,5-trimethoxyphenyl)porphyrin (127): The synthesis followed the standard procedure given for **82** using **123** (30.4 mg, 0.04 mmol), 4-dimethylaminophenyl boronic acid (82.5 mg, 0.5 mmol), Pd(PPh₃)₄ (5.77 mg, 0.005 mmol), and K₃PO₄ (212.2 mg, 1 mmol) in THF (50 mL). Column chromatography on silica gel (*n*-hexane/ethyl acetate = 6:1, v/v) gave the desired product as purple crystals (3 mg, 0.004 mmol, 10 %) and 5-hexyl-10-(3-nitrophenyl)-15-(3,4,5-trimethoxyphenyl)porphyrin **85** (11 mg, 0.16 mmol, 40 %) as a purple solid. Data for **127**: M.p. >300 °C. *R_f* = 0.4 (*n*-hexane/ethyl acetate, 6:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.67 (s, 2H, NH), 0.97 (t, 3H, *J* = 6.43 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 1.19 (t, 3H, *J* = 7.01 Hz, CH₂CH₂CH₂CH₃), 1.44 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.55 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.87 (m, 4H, CH₂), 2.58 (m, 4H, CH₂), 3.99 (s, 6H, OCH₃), 4.21 (s, 3H, *p*-OCH₃), 5.05 (m, 4H, CH₂), 7.46 (s, 2H, Ar-H), 7.96 (t, 1H, *J* = 8.18 Hz, Ar-H), 8.31 (d, 1H, *J* = 7.6 Hz, Ar-H), 8.64 (d, 1H, *J* = 4.67 Hz, β-pyrrole-H), 8.70 (d, 1H, *J* = 7.6 Hz, Ar-H), 8.76 (d, 1H, *J* = 4.1 Hz, β-pyrrole-H), 8.91 (d, 1H, *J* = 4.09 Hz, β-pyrrole-H), 9.00 (d, 1H, *J* = 4.09 Hz, β-pyrrole-H), 9.07 (s, 1H, Ar-H), 9.52 (m, 2H, β-pyrrole-H), 9.63 ppm (s, 2H, β-pyrrole-H). ¹³C NMR (150 MHz, CDCl₃): δ = 13.9, 14.0, 22.5, 23.5, 25.4, 29.5, 30.1, 31.7, 35.3, 38.8,

40.8, 56.2, 61.1, 67.8, 112.6, 114.8, 118.7, 120.6, 120.9, 122.6, 127.4, 128.1, 137.4, 139.5, 144.0, 146.8, 151.2 ppm. UV/vis (CH₂Cl₂): λ_{max} (log ϵ) = 421 (5.16), 519 (4.04), 555 (3.86), 594 (3.77), 653 nm (3.64). HRMS calcd. for C₄₅H₄₇N₅O₅ [M+H]⁺ 738.3655; found 738.3662.

5-Hexyl-10-(3-nitrophenyl)-15-(3,4,5-trimethoxyphenyl)-20-(N,N-dimethylaminophenyl) porphyrin (128): The synthesis followed the standard procedure given for **82** using **123** (30.4 mg, 0.04 mmol), 4-dimethylaminophenyl boronic acid (82.5 mg, 0.5 mmol), Pd(PPh₃)₄ (5.77 mg, 0.005 mmol), and K₃PO₄ (212.2 mg, 1 mmol) in THF (50 mL). Column chromatography on silica gel (*n*-hexane/ethyl acetate = 20:1, v/v) gave the desired product as purple crystals (14.5 mg, 0.018 mmol, 22 %). M.p. >300 °C. *R*_f = 0.75 (CH₂Cl₂). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.73 (s, 2H, NH), 0.95 (t, 3H, *J* = 7.02 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 1.42 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.52 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.84 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.20 (s, 6H, N(CH₃)₂), 2.6 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 3.98 (s, 6H, OCH₃), 4.15 (s, 3H, OCH₃), 4.15 (s, 3H, OCH₃), 5.06 (t, 2H, *J* = 7.6 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 7.45 (s, 2H, Ar-*H*), 7.99 (t, 1H, *J* = 7.6 Hz, Ar-*H*), 8.32 (d, 2H, *J* = 7.6 Hz, Ar-*H*), 8.48 (d, 2H, *J* = 7.6 Hz, Ar-*H*), 8.57 (d, 1H, *J* = 7.6 Hz, Ar-*H*), 8.71 (m, 2H, Ar-*H*), 8.78 (d, 1H, *J* = 5.27 Hz, β -pyrrole-*H*), 8.82 (d, 1H, *J* = 5.26 Hz, β -pyrrole-*H*), 8.90 (d, 1H, *J* = 5.26 Hz, β -pyrrole-*H*), 8.96 (t, 2H, *J* = 5.26 Hz, β -pyrrole-*H*), 9.09 (s, 1H, β -pyrrole-*H*), 9.56 ppm (dd, 2H, *J* = 4.68, 5.27 Hz, β -pyrrole-*H*). ¹³C NMR (150 MHz, CDCl₃): δ = 14.1, 22.7, 35.6, 36.0, 37.0, 37.4, 38.1, 38.5, 38.7, 39.3, 40.5, 42.0, 45.9, 52.4, 53.0, 56.3, 61.2, 107.2, 112.8, 121.7, 122.9, 127.6, 127.9, 128.2, 134.4, 139.6 ppm. UV/vis (CH₂Cl₂): λ_{max} (log ϵ) = 422 (5.10), 519 (4.07), 556 (3.90), 598 (3.80), 653 nm (3.85). HRMS calcd. for C₄₉H₄₈N₆O₅ [M+H]⁺ 801.3764; found 801.3745.

5-Hexyl-10-(3-nitrophenyl)-15-(3,4,5-trimethoxyphenyl)-20-(4-hydroxyphenyl)porphyrin (129): The synthesis followed the standard procedure given for **82** using **123** (30.4 mg, 0.04 mmol), 4-hydroxyphenyl boronic acid (68.95 mg, 0.5 mmol), Pd(PPh₃)₄ (5.77 mg, 0.005 mmol) and K₃PO₄ (212.2 mg, 1 mmol) in THF (50 mL). Column chromatography on silica gel (*n*-hexane/ethyl acetate = 6:1, v/v) gave the desired product as purple crystals (21 mg, 0.027 mmol, 70 %). M.p. >300 °C. *R*_f = 0.3 (*n*-hexane/ethyl acetate, 6:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.71 (s, 2H, NH), 0.95 (t, 3H, *J* = 7.01 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 1.41 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.54 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.85 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.58 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 3.99 (s, 6H, OCH₃), 4.20 (s, 3H, *p*-OCH₃), 5.04 (t, 2H, *J* = 7.6 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 5.21 (br s, 1H, OH), 7.24 (d, 2H, *J* = 8.77 Hz, Ar-*H*), 7.46 (s, 2H, Ar-*H*), 7.98 (t, 1H, *J* = 7.6 Hz, Ar-*H*), 8.09 (d, 2H, *J* = 8.18 Hz, Ar-*H*), 8.57 (d, 1H, *J* = 7.6 Hz, Ar-*H*), 8.69 (d, 1H, *J* = 4.67 Hz, β -pyrrole-*H*), 8.72 (m, 1H, Ar-*H*), 8.81 (d, 1H, *J* = 4.67 Hz, β -pyrrole-*H*), 8.88 (d, 1H, *J* = 4.67 Hz, β -pyrrole-*H*), 8.94 (d, 1H, *J* = 4.68 Hz, β -pyrrole-*H*), 8.96 (d, 1H, *J* = 4.68 Hz, β -pyrrole-*H*), 8.99 (d, 1H, *J* = 5.26 Hz, β -pyrrole-*H*), 9.10 (m, 2H, Ar-*H*), 9.52 (d, 1H, *J* = 4.68 Hz, β -pyrrole-*H*), 9.55 ppm (d, 2H, *J* = 5.26 Hz, β -pyrrole-*H*).

1H, *J* = 4.68 Hz, β -pyrrole-*H*). ¹³C NMR (150 MHz, CDCl₃): δ = 13.9, 22.5, 29.2, 29.5, 30.1, 31.7, 35.4, 38.8, 56.2, 61.1, 122.7, 113.5, 115.5, 119.3, 120.0, 121.2, 122.7, 128.1, 134.5, 135.4, 137.2, 137.7, 139.5, 144.1, 146.8, 151.3 ppm. UV/vis (CH₂Cl₂): λ_{max} (log ϵ) = 422 (4.88), 519 (3.88), 552 (3.73), 596 (3.66), 653 nm (3.49). HRMS calcd. for C₄₇H₄₃N₅O₆ [M+H]⁺ 774.3292; found 774.3260.

5-(4-Bromophenyl)-10-(3,4,5-trimethoxyphenyl)-15-(3-nitrophenyl)-20-hexylporphyrin (130): The synthesis followed the standard procedure given for **82** using **123** (30.4 mg, 0.04 mmol), 4-bromophenyl boronic acid (100.4 mg, 0.5 mmol), Pd(PPh₃)₄ (5.77 mg, 0.005 mmol), and K₃PO₄ (212.2 mg, 1 mmol) in THF (50 mL). Column chromatography on silica gel (*n*-hexane/ethyl acetate = 10:1, v/v) gave the desired product as purple crystals (18 mg, 0.022 mmol, 53 %) accompanied by **85** (40%). M.p. >300 °C. *R*_f = 0.25 (*n*-hexane/ethyl acetate, 4:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.62 (s, 2H, NH), 0.96 (t, 3H, *J* = 6.84 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 1.43 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.56 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.58 (s, 6H, CH₃), 1.86 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.60 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.68 (s, 3H, *p*-CH₃), 3.99 (s, 6H, OCH₃), 4.19 (s, 3H, *p*-OCH₃), 5.03 (t, 2H, *J* = 7.83 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 7.33 (s, 2H, Ar-*H*), 7.47 (s, 2H, Ar-*H*), 7.97 (t, 1H, *J* = 7.83 Hz, Ar-*H*), 8.55 (d, 1H, *J* = 7.83 Hz, Ar-*H*), 8.68 (d, 1H, *J* = 3.91 Hz, β -pyrrole-*H*), 8.72 (d, 1H, *J* = 4.89 Hz, β -pyrrole-*H*), 8.80 (d, 2H, *J* = 3.91 Hz, β -pyrrole-*H*), 8.82 (d, 1H, *J* = 7.82 Hz, Ar-*H*), 8.90 (d, 1H, *J* = 3.92 Hz, β -pyrrole-*H*), 8.95 (d, 1H, *J* = 4.89 Hz, β -pyrrole-*H*), 9.09 (s, 1H, Ar-*H*), 9.48 (d, 1H, *J* = 4.89 Hz, β -pyrrole-*H*), 9.55 ppm (d, 1H, *J* = 4.89 Hz, β -pyrrole-*H*). ¹³C NMR (150 MHz, CDCl₃): δ = 14.1, 21.5, 21.6, 22.7, 29.6, 29.7, 30.3, 31.9, 35.5, 38.9, 56.3, 61.3, 112.7, 112.8, 115.6, 118.7, 119.1, 121.1, 122.8, 127.5, 127.8, 128.2, 137.2, 138.3, 139.4, 139.6, 144.3 ppm. UV/vis (CH₂Cl₂): λ_{max} (log ϵ) = 420 (4.71), 517 (3.60), 552 (3.30), 594 (3.44), 647 nm (3.07). HRMS calcd. for C₅₀H₄₉N₅O₅ [M+H]⁺ 800.3812; found 800.3820.

5-Hexyl-10-(3-nitrophenyl)-15-(3,4,5-trimethoxyphenyl)-20-(2,4,6-trimethylphenyl)porphyrin (131): The synthesis followed the standard procedure given for **82** using **123** (30.4 mg, 0.04 mmol), 2,4,6-trimethylphenyl boronic acid (164 mg, 1 mmol) and Pd(PPh₃)₄ (5.77 mg, 0.005 mmol) and K₃PO₄ (212.2 mg, 1 mmol) in THF (50 mL). Column chromatography on silica gel (*n*-hexane/ethyl acetate = 10:1, v/v) gave the desired product as purple crystals (15 mg, 0.017 mmol, 45 %). M.p. >300 °C. *R*_f = 0.37 (*n*-hexane/ethyl acetate, 6:1, v/v). ¹H NMR (400 MHz, CDCl₃, TMS): δ = -2.74 (s, 2H, NH), 0.96 (t, 3H, *J* = 7.02 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 1.42 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.53 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 1.85 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 2.57 (m, 2H, CH₂CH₂CH₂CH₂CH₂CH₃), 3.99 (s, 6H, OCH₃), 4.20 (s, 3H, *p*-OCH₃), 5.05 (t, 2H, *J* = 7.01 Hz, CH₂CH₂CH₂CH₂CH₂CH₃), 7.46 (s, 2H, Ar-*H*), 7.98 (d, 2H, *J* = 8.18 Hz, Ar-*H*), 7.99 (t, 1H, *J* = 8.19 Hz, Ar-*H*), 8.10 (d, 2H, *J* = 7.6 Hz, Ar-*H*), 8.57 (d, 1H, *J* = 7.02 Hz, Ar-*H*), 8.71 (d, 1H, *J* = 4.68 Hz, β -pyrrole-*H*), 8.73 (s, 1H, Ar-*H*), 8.82 (d, 2H, *J* = 4.68 Hz, β -pyrrole-*H*), 8.95 (m, 3H, β -pyrrole-*H*), 9.10 (s, 1H, Ar-*H*), 9.55 ppm (d, 2H, *J* = 5.26 Hz, β -pyrrole-*H*).

pyrrole-*H*). ^{13}C NMR (150 MHz, CDCl_3): δ = 13.9, 22.5, 22.8, 23.6, 28.7, 29.2, 29.5, 29.8, 30.0, 30.1, 30.2, 31.7, 32.6, 35.4, 36.9, 38.6, 38.8, 56.2, 61.1, 68.0, 112.7, 115.9, 118.4, 119.5, 121.4, 122.4, 122.7, 127.4, 128.1, 128.6, 128.7, 129.7, 130.7, 131.1, 132.0, 135.6, 137.0, 137.8, 139.4 ppm. UV/vis (CH_2Cl_2): λ_{max} (log ϵ) = 421 (5.01), 519 (4.09), 552 (3.82), 592 (3.76), 648 nm (3.62). HRMS calcd. for $\text{C}_{49}\text{H}_{48}\text{N}_6\text{O}_5$ [$\text{M}+\text{H}$] $^+$ 836.2448; found 836.2481.

5-(*sec*-Butyl)-10-hexyl-15-(5-bromo-3-pyridyl)-20-(3,4,5-trimethoxyphenyl)porphyrin (132): The synthesis followed the standard procedure given for **82** using **124** (16 mg, 0.022 mmol), 3-bromopyridine-5-boronic acid (44.4 mg, 0.22 mmol), $\text{Pd}(\text{PPh}_3)_4$ (2.54 mg, 0.002 mmol) and K_3PO_4 (93.4 mg, 0.44 mmol) in THF (50 mL). Column chromatography on silica gel (*n*-hexane/ CH_2Cl_2 = 3:1, v/v) followed by a second column using *n*-hexane/ethyl acetate (3:1, v/v) gave the desired product as purple crystals (8 mg, 0.010 mmol, 47 %). M.p. >300 °C. R_f = 0.25 (*n*-hexane/ CH_2Cl_2 , 1:3, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): δ = -2.64 (s, 2H, NH), 0.98 (t, 3H, J = 7.16 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.09 (t, 3H, J = 7.15 Hz, CHCH_2CH_3), 1.44 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.57 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.88 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.46 (d, 3H, J = 7.7 Hz, CHCH_3), 2.58 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.85 (m, 1H, CHCH_2CH_3), 2.99 (m, 1H, CHCH_2CH_3), 3.99 (s, 6H, OCH_3), 4.21 (s, 3H, *p*- OCH_3), 5.05 (t, 2H, J = 8.05 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 5.41 (m, 1H, CHCH_2CH_3), 7.42 (s, 2H, Ar-*H*), 8.66 (s, 1H, Ar-*H*), 8.69 (d, 1H, J = 4.95 Hz, β -pyrrole-*H*), 8.81 (d, 1H, J = 4.95 Hz, β -pyrrole-*H*), 8.97 (d, 1H, J = 4.4 Hz, β -pyrrole-*H*), 9.10 (s, 1H, Ar-*H*), 9.33 (s, 1H, Ar-*H*), 9.55 (d, 1H, J = 4.95 Hz, β -pyrrole-*H*), 9.62 (d, 2H, J = 4.4 Hz, β -pyrrole-*H*), 9.76 ppm (d, 2H, J = 4.4 Hz, β -pyrrole-*H*). ^{13}C NMR (150 MHz, CDCl_3): δ = 13.9, 14.0, 14.1, 18.4, 19.6, 22.5, 22.6, 22.6, 27.2, 29.2, 29.5, 29.9, 30.2, 31.7, 35.7, 36.9, 38.9, 42.7, 56.2, 61.1, 111.7, 112.4, 118.7, 119.4, 120.7, 125.1, 126.0, 128.0, 128.6, 128.9, 130.7, 130.8, 137.5, 137.7, 139.3, 143 ppm. UV/vis (CH_2Cl_2): λ_{max} (log ϵ) = 420 (4.88), 519 (3.88), 552 (3.58), 598 (3.49), 647 nm (3.36). HRMS calcd. for $\text{C}_{44}\text{H}_{46}\text{N}_5\text{O}_3$ [$\text{M}+\text{H}$] $^+$ 772.2862; found 772.2895.

5-(*n*-Butyl)-10-(4-methoxycarbonylphenyl)-15-(3-bromo-2,4,6-trimethoxyphenyl)-20-(3-nitrophenyl)porphyrin (133): The synthesis followed the standard procedure given for **82** using **125** (30 mg, 0.0351 mmol), 3-nitrophenyl boronic acid (58.29 mg, 0.351 mmol), $\text{Pd}(\text{PPh}_3)_4$ (4.05 mg, 0.00351 mmol) and K_3PO_4 (149.37 mg, 0.703 mmol) in THF (50 mL). Column chromatography on silica gel (*n*-hexane/ CH_2Cl_2 = 1:1, v/v) gave as the first fraction 5-(*n*-butyl)-10-(4-methoxycarbonylphenyl)-15-(3-bromo-2,4,6-trimethoxyphenyl)-20-(3-nitrophenyl)porphyrin **133** (17 mg, 0.0196 mmol, 55 %) as purple crystals and as the second fraction 5-(*n*-butyl)-10-(4-methoxycarbonylphenyl)-15-(2,4,6-trimethoxy-3'-nitrobiphenyl-3-yl)-20-(3-nitrophenyl)porphyrin **134** (6 mg, 0.0066 mmol, 18 %) as purple crystals. Data for **133**: M.p. >300 °C. R_f = 0.5 (*n*-hexane/ CH_2Cl_2 , 1:4, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): δ = -2.63 (s, 2H, NH), 1.12 (t, 3H, J = 7.15 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.83 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.54 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 3.00 (s, 3H, OCH_3), 3.57 (s, 3H, OCH_3), 4.15 (s, 3H, OCH_3), 4.22 (s, 3H, OCH_3), 5.06 (t, 2H, J = 7.53 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 6.78 (s, 1H, Ar-*H*), 7.96

(dd, 1H, J = 8.66 Hz, Ar-*H*), 8.32 (m, 2H, Ar-*H*), 8.47 (d, 2H, J = 8.18 Hz, Ar-*H*), 8.54 (d, 1H, J = 7.15 Hz, Ar-*H*), 8.57 (d, 1H, J = 7.16 Hz, Ar-*H*), 8.68 (s, 1H, β -pyrrole-*H*), 8.71 (d, 1H, J = 8.66 Hz, β -pyrrole-*H*), 8.76 (d, 1H, J = 4.52 Hz, β -pyrrole-*H*), 8.79 (s, 1H, β -pyrrole-*H*), 8.84 (m, 1H, β -pyrrole-*H*), 9.10 (m, 1H, β -pyrrole-*H*), 9.54 (d, 1H, J = 4.52 Hz, β -pyrrole-*H*), 9.56 ppm (d, 1H, J = 4.89 Hz, β -pyrrole-*H*). ^{13}C NMR (150 MHz, CDCl_3): δ = 8.10, 13.9, 14.0, 22.5, 23.4, 23.7, 29.2, 29.5, 31.7, 35.2, 40.8, 52.2, 56.1, 56.5, 60.8, 92.4, 98.3, 98.4, 110.2, 115.7, 118.3, 118.4, 121.7, 122.7, 127.3, 127.4, 127.7, 128.0, 129.4, 134.3, 139.4, 139.5, 144.0, 146.8, 146.9, 158.0, 159.9, 167.1 ppm. UV/vis (CH_2Cl_2): λ_{max} (log ϵ) = 421 (5.16), 516 (4.01), 552 (3.63), 594 (3.84), 648 nm (3.78). HRMS calcd. for $\text{C}_{47}\text{H}_{42}\text{N}_5\text{O}_7\text{Br}$ [$\text{M}+\text{H}$] $^+$ 868.2169; found 868.2150.

5-(*n*-Butyl)-10-(4-methoxycarbonylphenyl)-15-(2,4,6-trimethoxy-3'-nitro-biphenyl-3-yl)-20-(3-nitrophenyl)porphyrin (134): This compound was derived from the synthesis of **133**. M.p. >300 °C. R_f = 0.37 (*n*-hexane/ CH_2Cl_2 , 1:4, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): δ = -2.62 (s, 2H, NH), 1.15 (t, 3H, J = 7.15 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.83 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.30 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.55 (s, 3H, OCH_3), 3.63 (s, 3H, OCH_3), 4.10 (s, 3H, OCH_3), 4.16 (s, 3H, OCH_3), 5.07 (t, 2H, J = 7.91 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 6.84 (s, 1H, Ar-*H*), 7.60 (t, 1H, J = 7.9 Hz, Ar-*H*), 7.97 (m, 1H, Ar-*H*), 8.04 (d, 1H, J = 7.9 Hz, Ar-*H*), 8.20 (d, 1H, J = 7.9 Hz, Ar-*H*), 8.32 (m, 2H, Ar-*H*), 8.47 (d, 2H, J = 8.28 Hz, Ar-*H*), 8.56 (m, 2H, Ar-*H*), 8.71 (m, 2H, Ar-*H*), 8.79 (s, 2H, β -pyrrole-*H*), 8.87 (d, 1H, J = 4.89 Hz, β -pyrrole-*H*), 8.95 (m, 2H, β -pyrrole-*H*), 9.09 (s, 1H, β -pyrrole-*H*), 9.54 (d, 1H, J = 4.52 Hz, β -pyrrole-*H*), 9.55 ppm (d, 1H, J = 4.52 Hz, β -pyrrole-*H*). UV/vis (CH_2Cl_2): λ_{max} (log ϵ) = 421 (4.93), 516 (3.95), 552 (3.65), 592 (3.73), 649 nm (3.80). HRMS calcd. for $\text{C}_{53}\text{H}_{44}\text{N}_6\text{O}_9$ [$\text{M}+\text{H}$] $^+$ 909.3248; found 909.3229. $\text{C}_{53}\text{H}_{44}\text{N}_6\text{O}_9$ (908.97): calcd. C 70.03, H 4.88, N 9.25; found C 70.33, H 5.15, N 9.52.

5-(*n*-Butyl)-10-(4-methoxycarbonylphenyl)-15-(3-bromo-2,4,6-trimethoxyphenyl)-20-(5-bromo-2-fluoro-3-pyridinyl)porphyrin (135): The synthesis followed the standard procedure given for **82** using **125** (30 mg, 0.0351 mmol), 5-bromo-2-fluoro-3-pyridine boronic acid (80.2 mg, 0.365 mmol), $\text{Pd}(\text{PPh}_3)_4$ (4.2 mg, 0.00365 mmol) and K_3PO_4 (165.87 mg, 0.729 mmol) in THF (50 mL). Column chromatography on silica gel (*n*-hexane/ CH_2Cl_2 = 1:1, v/v) gave the title compound as purple crystals (11 mg, 0.011 mmol, 35 %). M.p. >300 °C. R_f = 0.22 (*n*-hexane/ CH_2Cl_2 , 1:1, v/v). ^1H NMR (400 MHz, CDCl_3 , TMS): δ = -2.63 (s, 2H, NH), 1.15 (t, 3H, J = 7.15 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.83 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.54 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 3.00 (s, 3H, OCH_3), 3.57 (s, 3H, OCH_3), 4.15 (s, 3H, OCH_3), 4.23 (s, 3H, OCH_3), 5.06 (t, 2H, J = 7.6 Hz, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 6.79 (s, 1H, Ar-*H*), 8.30 (t, 2H, J = 8.18 Hz, Ar-*H*), 8.47 (d, 2H, J = 8.18 Hz, Ar-*H*), 8.66 (d, 1H, J = 2.34 Hz, Ar-*H*), 8.71 (m, 1H, β -pyrrole-*H*), 8.75 (d, 1H, J = 4.68 Hz, β -pyrrole-*H*), 8.79 (m, 1H, β -pyrrole-*H*), 8.85 (m, 3H, β -pyrrole-*H*), 9.53 (d, 1H, J = 4.67 Hz, β -pyrrole-*H*), 9.59 ppm (d, 1H, J = 4.68 Hz, β -pyrrole-*H*). ^{13}C NMR (150 MHz, CDCl_3): δ = 13.9, 14.0, 19.5, 22.5, 23.4, 24.7, 29.2, 29.5, 31.7, 35.1, 40.8, 52.2, 56.1, 56.6, 60.8, 92.4, 98.4, 103.7, 107.8, 109.6, 109.8, 110.5, 115.5, 118.3, 119.0, 122.0, 127.6, 129.5, 134.2, 146.8, 147.5, 147.6, 148.5,

148.6, 158.0, 158.5, 159.9, 160.3 ppm. UV/vis (CH₂Cl₂):
 λ_{max} (log ϵ) = 420 (5.3), 516 (4.26), 550 (3.96), 592 (3.66),
648 nm (3.74). HRMS calcd. for C₄₆H₃₈N₅O₅Br₂F [M+H]⁺
920.1281; found 920.1276.