

Figure #	Figure title One sentence only	Filename This should be the name the file is saved as when it is uploaded to our system. Please include the file extension. i.e.: <i>Smith_ED Fig1.jpg</i>	Figure Legend If you are citing a reference for the first time in these legends, please include all new references in the Online Methods References section, and carry on the numbering from the main References section of the paper.
Extended Data Fig. 1	Spatial distribution of Australian fish survey data used coloured according to the mean annual temperature	Audzi_ED_Fig1.pdf	Spatial distribution of Australian fish survey data used coloured according to the mean annual temperature. All sites are grouped onto 0.5 degree grid cells. Colours represent mean annual SST over the entire sampling period in that cell (ranging from 12°C for yellow to 29°C for red) and circle size is proportional to the total number of species used for the analyses in the cell (determined by species richness and number of surveys in the cell). Black stars indicate the nine long-term monitoring locations. The Australian coastline shapefile was downloaded from the Australian Natural Resources Data Library website (Commonwealth of Australia)
Extended Data Fig. 2	Correlation between species body size – SST slopes and abundance – SST slopes from literature.	Audzi_ED_Fig2.pdf	Species-specific slopes between body size and mean annual SST through space (β_1 , on y axis) are compared with the species specific abundance and mean annual SST slopes estimated in Waldock et al. (2019). Data on both slopes was available for 300 species. If abundance was a major driver of average body size, we could expect an overall negative correlation between size-temperature and abundance-temperature slopes, such as mean body size is smaller at sites with higher abundances. Alternatively, larger average body sizes might have positive effects on abundance through e.g. improved recruitment or inter-specific effects. In this case we could expect a positive correlation between size-temperature and abundance-temperature slopes. The correlation between body size slopes and abundances was close to zero ($r = 0.09$, $p = 0.14$), suggesting that abundance is unlikely to be a major driver of body sizes and vice versa.

Extended Data Fig. 3	Species-specific body length changes at nine long-term monitored and warming locations	Audzi_ED_Fig3.pdf	Location and species specific temporal responses, represented by slopes of body length change in 105 coastal fish species in nine warming and long-term monitored locations. Each dot represents a species, arranged according to the temperature at the centre of their distribution area (temperature midpoint). Blue and red colours indicate species for which 90% of the posterior probability density range for the slope of the annual body length change (on y axis) was above or below zero, respectively. For the map of the nine locations see Extended Data Fig. 1.
----------------------	--	-------------------	---

2

Item	Present?	Filename This should be the name the file is saved as when it is uploaded to our system, and should include the file extension. The extension must be .pdf	A brief, numerical description of file contents. <i>i.e.: Supplementary Figures 1-4, Supplementary Discussion, and Supplementary Tables 1-4.</i>
Supplementary Information	Yes	Audzi_SupplementaryInfo.pdf	Supplementary Tables 2-3, Supplementary Figures 1-4.
Reporting Summary	Yes	Audzi_ReportingSummary.pdf	

3

Type	Number If there are multiple files of the same type this should be the numerical indicator. i.e. "1" for Video 1, "2" for Video 2, etc.	Filename This should be the name the file is saved as when it is uploaded to our system, and should include the file extension. i.e.: <i>Smith_Supplementary Video 1.mov</i>	Legend or Descriptive Caption Describe the contents of the file
Supplementary Table	Supplementary Table 1	Audzi_SupplementaryTables1_4.xlsx	List of Bayesian estimates of body size – SST slopes through space for each of the 335 individual species used in the analyses (Table S1)

			and Bayesian estimates of annual changes in body lengths (slopes of size vs year) in nine long-term locations (Table S4)
Supplementary Data	Data files	Audzi_SupplementaryData.zip	Filtered and final datasets used in the analyses.

Fish body sizes change with temperature, but not all species shrink with warming

Asta Audzijonyte^{1,2}, Shane A. Richards³, Rick D. Stuart-Smith¹, Greta Pecl^{1,2}, Graham J. Edgar¹, Neville S. Barrett¹, Nicholas Payne⁴ and Julia L. Blanchard^{1,2}

¹Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, Australia

² Centre for Marine Socioecology, University of Tasmania, Hobart, Tasmania, Australia

³School of Natural Sciences, University of Tasmania, Hobart, Tasmania, Australia

⁴School of Natural Sciences, Trinity College Dublin, Ireland

Summary

Ectotherms generally shrink under experimental warming, but whether this pattern extends to wild populations is uncertain. We analysed ten million visual survey records, spanning the Australian continent, multiple decades and comprising all common coastal reef fish (335 species). We found that temperature indeed drives spatial and temporal changes in fish body size, but not consistently in the negative fashion expected. Around 55% of species were smaller in warmer waters (especially among small-bodied species) while 45% were bigger. The direction of a species' response to temperature through space was generally consistent with its response to temperature increase at any

given location, suggesting that spatial trends could help forecast fish responses to warming. However, temporal changes were ~10x faster than spatial trends (~4% versus ~40% body size change per 1°C change through space and time respectively). The rapid and variable responses of fish sizes to warming may herald unexpected impacts on ecosystem restructure, with potentially greater consequences than if all species were shrinking.

One sentence summary (125 characters): Spatial relationships between fish body size and temperature can help predict responses to warming through time

Main

Body size is a key biological and ecological trait^{1,2}, but many natural populations have undergone declines in average body size due to harvesting, with consequent reductions in ecosystem functional diversity and resilience³⁻⁵. More equivocal are declines in ectotherm body size caused by increased temperature, even though such declines are considered a third universal response to global warming⁶, and, unlike harvesting, act on all species in an ecosystem. While some models forecast a 15-30% decrease in body sizes due to warming by 2050^{7,8}, the generality of these responses remains debated on the theoretical basis⁹ and in the light of some empirical observations¹⁰. Within species, smaller adult body sizes at warmer temperatures are commonly found under experimental conditions, having been observed in animals as diverse as insects, worms, reptiles and fishes¹¹. Nevertheless, the mechanisms and adaptive significance of this phenomenon, often described as a temperature-size rule, remain elusive and controversial since Bergman's publication in 1847¹¹⁻¹³. Experimental temperature-size responses are particularly apparent for aquatic species, a pattern that has spawned hotly debated hypotheses on the role of oxygen supply¹⁴⁻¹⁶. For example, an average adult body weight decrease of 5% per 1°C of warming was observed across 169 aquatic species, versus just 0.5% decrease in terrestrial taxa¹⁴. Yet, the applicability of experimental studies to wild populations remains questionable, because few experiments have explored body size responses through multiple

generations where intergenerational plasticity and rapid adaptations are likely to alter the observed responses¹⁷. Moreover, the realised size in wild populations integrates growth, recruitment, mortality, competition, predation risk and food availability simultaneously, and the interplay of these factors cannot be adequately addressed in experiments¹⁰.

Climate change ecology has thus far largely focused on shifting geographic distributions, phenology and abundance of organisms, and the consequences of these¹⁸, whereas warming-driven changes in body sizes across a broad range of species (both unexploited and exploited) have not been systematically assessed. This study capitalises on large underwater visual survey datasets from the National Reef Monitoring Network in Australia (consisting of the Australian Temperate Reef Collaboration (<https://atrc.org.au/>) and Reef Life Survey (<https://reeflifesurvey.com/>))^{19,20}. The datasets consist of >30,000 surveys of rocky and coral reefs around the Australian continent (Extended Data Fig. 1), where abundance and sizes of all fish species along standardised transects is recorded by trained divers. Some locations have been continuously monitored for over 26 years. From these surveys we selected all fish species that satisfied the minimum abundance and occurrence frequency criteria, resulting in 335 common coastal species spanning a range of maximum body size and life-history characteristics. Majority of these species (254 out of 335) are unlikely to be targeted by fishing in the study region, and only 42 species are known to be commonly caught by recreational or commercial fishers (the remaining 39 species could be occasionally caught, but fishing mortality is likely to be low, see Supplementary Information Table 1). We then use Bayesian methods to fit hierarchical, mixed-effects models that account for random errors in space and time to quantify spatial and temporal relations between mean individual body length and temperature or time.

Results

In the first set of analyses, species' mean annual body length was assessed against the annual mean sea surface temperature (SST) at locations spanning their distributions. Approximately half of the species showed clear trends in body size with temperature (for which lower or upper 10% of the posterior probability density or PPD range of the size-SST slope was above or below zero respectively), with 97 species declining in size towards the warmer edge of their distribution, and 64 species increasing (Fig. 1A, Supplementary Information Table 2). Fish species included in our study grouped into two major thermal guilds²¹, corresponding to temperate and tropical distribution types. In our analyses, temperate species constituted about a third of the total species analysed and were equally likely to have negative or positive body size responses to temperature (Fig. 1A, Supplementary Information Table 2). In contrast, tropical species were twice as likely to be smaller than larger at their warm distribution edge (see Supplementary Information Table 1 showing the list of species and their individual body size responses to temperature). In addition, relative change in species' mean body length was, on average, positively related to the maximum body length (Fig. 1B), meaning that small-bodied species tended to be smaller in warmer areas and larger species tended to be larger. This contradicts experimental temperature-size observations, where strongest declines in adult body sizes occurred in largest species¹⁴ and where oxygen supply was suggested as playing a critical role. Given that small individuals are more likely to be overestimated in size by divers undertaking surveys (due to inherent and consistent biases in diver perceptions of sizes underwater²²), our result is likely conservative. Notably, body size-temperature trends in space were most often linear, or at least not strongly curved, suggesting that for most species there was not an "optimal" temperature in the central part of the distribution range at which body size was maximised (Supplementary Information Fig. 1). Density-dependent processes, such as abundance skew statistics across species distribution ranges²³, did not appear to explain the observed size trends either (see Methods and Extended Data Fig. 2).

While the spatial analyses ascertained length-temperature relations across many species' distributions, we sought further evidence that these body size changes related specifically to temperature by quantifying intra-specific changes in mean body length over time for all common fish species at nine sites monitored for 15 to 26 years. Specifically, we estimated temporal trends in body size for 105 species; of these 77 species were found at eight locations that have warmed significantly over the time of observations (average trend of 0.025°C per year). Clear temporal changes in body size (90% PPD outside zero) were seen in more than half of the studied fish species (40 out of 77) in these warming locations (Supplementary Information Table 2), and from these, most (36 out of 40) were consistently decreasing or increasing in sizes across different, geographically distant locations. No significant linear long-term warming trend was observed in one of the monitored locations in Western Australia (Jurien Bay, Supplementary Information Table 3, Extended Data Figs. 1, 3) and this location also had the smallest proportion of species with temporal trends in body size (Supplementary Information Table 3). Some of the fastest rates of warming were observed in the southernmost locations in Tasmania, where 50-66% of species showed clear changes in body size through time. The correlation between the rate of warming and the number of species changing in size was weaker closer to the tropics (Jervis Bay site), where, despite the fast warming, only a third of species were clearly increasing or decreasing in size. Importantly, for species that were included in both spatial and temporal analyses (71 species), long-term rates of change in body length at the eight warming locations were positively correlated with the relative change in body length observed across the temperature gradient in space (Fig 2). In other words, species that were smaller at the warmer edge of their geographical distribution were also more likely to become smaller in locations that are warming, and vice-versa.

The two sets of analyses strongly suggest that temperature is a major determinant of reef fish body sizes in the wild, and that the magnitude of average body size change across space and time can be surprisingly large (Fig. 2, Extended data Fig. 3). For example, for species that showed clear body

size changes with temperature through space (90% PPD outside zero, Fig. 1, constituting about half of all species), mean body length changed by ~4% for each degree of warming throughout their distribution (Supplementary Information Tables 1, 2). For a median length temperate fish (15 cm) with a typical body shape, this change in length corresponds to approximately 12% change in body mass per 1°C, and nearly a three-fold difference in mean mass between the equatorial and poleward range of an average species distribution (8°C of annual SST range, giving 12 cm and 32 g versus 17 cm and 89 g on the extreme temperature ranges across the distribution, see Supplementary Information Table 2). Temporal changes in body size were even greater, averaging roughly 1% change in length per year in the 40 clearly changing species (90% PPD outside zero). If the rate of body size change and warming observed over the last 20 years was maintained, then 1°C of warming predicted over the next 40 years would result in ~ 40% change (either increase or decrease) in fish body length. This means that body size responses to 1°C temperature change through time were about ten times faster than those seen through space (depicted with the orange line in Fig. 2). The result was similar if the space-time comparison was constrained to species for which both temporal and spatial data was available (for these species the average change through space was ~3.5% for each degree of warming).

Several alternative explanations exist for the different rates of response to temperature gradients in space and time. First, rapid warming in time also leads to species redistributions and changing species interactions may accentuate emergent body size changes through time. Second, rapid warming is also likely to coincide with changes in productivity and changes in the growing season itself, which again is likely to magnify changes in body sizes observed in the field. Third, species are adapted to the local temperature range, more so than to temperatures experienced across their full distributional range, and are consequently sensitive to relatively slight temperature changes in the local environment²⁴. Fourth, size patterns through space have emerged over many generations and likely involve adaptations that may reduce the temperature effects on growth. Counter-gradient

variation, or opposite influences of genetic and environmental factors on phenotypes, has been detected in dozens of fish species and often involve changes in physiological rates that dampen the effects of temperature²⁵. Presumably, rapid warming over the last two decades would not have allowed for the sufficient time for such adaptations to emerge. Nevertheless, even though the observed temporal rates of change are high, they are comparable to the rates of growth or maturation change reported in harvested fish species (ca 0.5-4% per year) and smaller than evolutionary change in size under strong selection experiments (2-17% per year)²⁶.

Discussion

Our study provides strong empirical support for the differential effects of warming on body sizes of many common coastal marine fishes. Until now, the majority of size trend information for marine species was only available from commercially harvested or charismatic species, where effects of harvesting and warming are difficult to separate²⁷. In our study, the effects of exploitation on body size are unlikely to affect overall conclusions because the number of commercially or recreationally fished species is small relative to the number of unexploited fish species inhabiting the shallow reefs surveyed, and because the temporal body size - temperature correlations inside and outside no-take marine protected areas were broadly similar (Supplementary Information Fig. 2). Species that showed the largest responses of body size to temperature represent useful priorities for further study to understand the underlying drivers, such as growth, mortality, recruitment, food availability and other demographic changes. Future work is also needed to disentangle the multiple ecological mechanisms and environmental stressors at play, as well as potential feedbacks. For example, how do warming-driven species redistributions interact with changes in body sizes? And what are the main reasons for different rates of responses through space and time?

Differential responses of species' body size to warming have implications for the restructuring of food webs and ecosystems, with consequences for the stability and resilience of local communities to other external stressors, such as fishing, coastal pollution, and the multi-faceted effects of climate change. Many biological functions scale allometrically with body size, hence the mean, variance, and shape of body size distributions within a community all influence aggregate ecosystem functioning²⁸. Modelling studies show that even small changes in species body sizes, such as 4% decrease over 50 years could lead to increased mortality and up to 30% decrease in biomasses and productivity^{29,30}, yet the body size changes of common species in warming locations are often much greater (Extended Data Fig. 3). Of particular consequence is our finding that body size changes across the community are not proportionate across species – total biomass and production of larger predatory species are likely to increase with rising temperatures whereas production of small bodied prey species will decline. Complex changes to food web structure can thus be expected. Understanding how such body size trends might change functional roles of dominant species in marine ecosystems should be one of the top priorities for management and protection of marine biodiversity in a rapidly changing ocean.

Methods

Reef fish data

The study is based on up to 26 years of underwater surveys from around Australian continent and includes around 10 million observations of 335 fish species from around 30000 surveys. Data were obtained from standardised quantitative censuses of fishes undertaken as part of the Australian Temperate Reef Collaboration (ATRC) monitoring program conducted from 1992 to present, and Reef Life Survey (RLS) program from 2008 to present. The RLS program was developed based on the ATRC program, and the two approaches use consistent methodology other than extent of replication. Full details of fish census methods are provided in^{19,20} and an online methods manual (<http://www.reeflifesurvey.com>) describes the different data collection methods. To maximise

consistency in this study, we only used data from Method 1 surveys. These surveys involve divers laying 50 m transect lines and recording all fish species present within duplicate 5 m wide blocks (total area = 500 m²), tallying abundance in size classes as the divers move along the transect. Fish sizes are recorded in $K = 28$ size (length) bin categories, with bin widths ranging from 2.5 cm for fish lengths below 15 cm, 5 cm for fish lengths between 15 and 40 cm, then 12.5 cm bins for fish lengths over 50 cm. Data quality and training of divers are detailed in ^{20,31}. ATRC data are only collected by professional marine scientists with field experience, while the RLS data were collected by selected, trained and experienced recreational divers, who were each provided one-on-one training and calibration with experienced scientists before contributing data to the database. The data collected by these trained RLS volunteers has been formally evaluated and found to be indistinguishable from those of professional scientists at the same time and place ³¹.

Temperature data

For the spatial analyses, we grouped all fish survey sites into half a degree grid cells, resulting in 280 cells spanning mean annual sea surface temperature (SST) from 12°C to 29°C (Extended Data Fig. 1). For each of these cells we then extracted mean yearly SST values using daily temperature records from the NOAA Optimum Interpolation 1/4 Degree Daily Sea Surface Temperature model data ³² from 1982 until the end of 2018. Alternative temperature metrics (such as growth degree days, i.e. days with temperature above 12°C, or mean annual SST in previous years) gave overall similar results. Long-term annual change in SST was also estimated for nine locations using monthly average SST values, and accounting for a sinusoidal component of seasonal temperature variation. For each location mean SST at time t was described by: $\mu(t) = \beta_0 + \beta_1 t + \beta_2 \cos(2\pi(t - \phi))$, where t is years since January 1, 1982. Variation in observed SST about this mean was assumed to be normally distributed with fixed standard deviation, σ . All parameters were estimated using Bayesian methods and the resulting long term rates of change, given by β_1 , are presented in the Supplementary Information Table 2 (analysis code available on <https://github.com/astaudzi/RLSfishSize>).

230 **Data filtering and selection**

231 Reef fish size data were assessed for outliers and possible taxonomic errors by: i) filtering out
232 observations in cells above and below 97.5% and 2.5% of the species distribution temperature (~1%
233 of records) and ii) removing records that were 1.5 times larger than maximum species size known in
234 FishBase (a larger cut-off was used because reported maximum size in FishBase is smaller than the
235 true maximum size for many unexploited species with limited life-history data) (~1.5% of records).
236 To ensure enough data and statistical power to infer body size responses and account for random
237 annual and spatial variation, the dataset used in the spatial analyses was restricted to taxa identified
238 to species level, and only species containing at least 1000 individuals occurring in at least 10
239 geographic cells and over at least 5 years (e.g. a species with 1000 individuals observed in a single
240 geographic cell or only recorded in 3 years was not used). We also excluded the smallest species that
241 had maximum body sizes below 10 cm, because their data spanned too few size bins to discriminate
242 trends. This gave 335 fish and shark species, across a range of temperature preferences and life-
243 history strategies. To reduce the effect of recruitment pulses and vagrant recruits occurring outside
244 species' normal distribution limits on estimates of body size - temperature responses, we removed
245 observations below the 10th percentile of the annual observed species body length (annual rather than
246 one overall length was used to allow for the body size trends) (~3% of observations). While out-of-
247 range observations may be important indicators of redistributions, we were interested in temperature
248 – body size responses over the core ranges occupied consistently by species. The possible effects of
249 this exclusion procedure on the conclusions from the hierarchical models (Fig.1A) were tested by
250 repeating spatial analyses using all data from 10 small and 10 large species; we found that data
251 exclusion did not affect the main conclusion (Supplementary Information Fig. 3). Temporal analyses
252 were restricted to locations where surveys spanned at least 15 years with at least 10 annual surveys
253 conducted in total. This left nine locations (Extended Data Fig. 1), each with an average of 44 sites
254 repeatedly monitored (Supplementary Information Table 2). To improve estimates of temporal trends

and associated random temporal errors, we restricted temporal analyses to species/location combinations where at least 20 individuals of a given species were observed per year at a location and at least 8 years of such observations were available (a species that was common for 5 years, but then fewer than 20 individuals observed per year in later years was not used, and likewise a location that was only sampled in e.g. 1992, 2000 and 2015 was not used).

Statistical analyses

The association between SST and the distribution of observed body sizes across a species' range was quantified using a hierarchical, mixed-effect model. We assumed that observed log-transformed fish lengths were drawn from a normal distribution with standard deviation, $\sigma_{\ln\text{length}}$, which was constant across all surveys but differed among species. Size distributions within one species generally followed a lognormal distribution (Supplementary Information Fig. 4). For survey i the mean of this distribution is given by:

$$\mu_i = \beta_0 + \beta_1 x_i + \alpha_{\text{cell}[i]} + \alpha_{\text{year}[i]} + \alpha_i,$$

where μ_i is the ln transformed body length, x_i is the normalised mean annual SST in the cell, and β_0 and β_1 are species-specific regression coefficients to be estimated. We normalised SST by subtracting the median SST observed across the species' range. This variable transformation improved convergence when fitting and reduced the potential correlation between β_0 and β_1 , thereby removing the need to also estimate an additional correlation parameter. A positive estimate for β_1 would indicate a positive association between local SST and mean fish length. The α -terms describe random effects due to spatial variation among surveyed cells ($\alpha_{\text{cell}[i]}$, due to e.g. different productivity, human pressure), temporal variation among survey years ($\alpha_{\text{year}[i]}$, due to random variation across years not associated with temperature), and random variation among the surveys themselves (α_i , e.g. observer, site, or weather differences). In this study a 'survey' is treated as a

specific day in a given cell and can comprise several 50-m transect surveys. Each of the three α -terms were assumed to be drawn from a normal distribution with zero mean and standard deviation to be estimated (denoted σ_{cell} , σ_{year} , σ_{survey}). To account for the observed fish lengths being necessarily binned we integrated this normal distribution across each of the K bin ranges to calculate the probability of fish being observed in each bin. To account for potential outliers in estimated lengths we also assumed that there was a small probability $\varepsilon = 0.01$ that a fish was observed in a randomly chosen bin. These probabilities form the basis of our likelihood function and Bayesian methods were then used to estimate the six model parameters for each species and assess their uncertainty. Uniform priors were used for all parameters: $[1.5-3.5]$ for β_0 , $[-0.5-0.5]$ for β_1 , $[0.001-1]$ for σ_{cell} and σ_{survey} , $[0.001-1.5]$ for σ_{year} , and $[0.1-0.3]$ for $\sigma_{\ln\text{length}}$. After the first set of analyses the posterior probability density plots were visually inspected for each species, and if the posterior densities were not fully included in the prior value range the priors were adjusted and analyses repeated. Yet, we found that priors had little influence on their posteriors due to large sample sizes. Markov Chain Monte Carlo (MCMC) sampling was run with 3 chains of 1000 iterations each, of which the first 500 were discarded as the burn-in and the last 500 were used to generate posterior probability density ranges. Some species did not converge at these settings and analyses were repeated using 1500 iterations and 1000 burn-in period. Further details and the code can be found at <https://github.com/astaaudzi/RLSfishSize>. Because μ_i is the log transformed body length, the estimated values of β_1 are converted to relative (or %) change in body length as $\exp(\beta_1)$, where β_1 of e.g. -0.02 corresponds to $\exp(-0.02) = 0.98$ or 2% decrease in body length. Thus β_1 values ranging from -0.2 to 0.2 can be interpreted as proportional change in body length.

Given estimates of the slope of mean body size against SST (β_1) for all species, we then assessed if these slopes were linearly associated with either the thermal affinity of a species or its maximum body-size. Species thermal affinity is defined as the SST at the midpoint of its realised thermal distribution, as described in ^{21,33} and also often referred to as the Species Temperature Index. For the

maximum body size we used the median value of five largest observed body size records, which alleviated potential observation errors in the data sets. We accounted for variation in uncertainty associated with each estimate of β_1 as follows. For each species, we calculated the median and the standard deviation of the posterior for β_1 . These standard deviations weighted the influence of each median on the linear regression, which we implemented using a hierarchical mixed-effects model. Again, Bayesian methods were used to estimate the regression coefficients and their associated uncertainty for both linear regressions (Fig. 1). Uniform priors were adopted for the y-intercept and slope ($[-0.1-0.1]$ in both cases) and a uniform prior of $[0.01,0.1]$ was also assumed for the standard deviation describing variation in the β_1 estimates about the fitted regression line (Fig. 1).

Evidence of local, linear trends in mean body size were based on observations for 105 species at nine long-term monitoring locations¹⁹. We adopted a model very similar to the one used for the spatial analysis, except we assumed a linear relation of body length to year, rather than SST, and allowed both the y-intercept and the slope to vary across locations. For survey i at location j the mean of the ln-length distribution of a species is given by:

$$\mu_{i,j} = \beta_{0,j} + \beta_{1,j}y_i + \alpha_{\text{year}[i]} + \alpha_i,$$

where y_i and $\text{year}[i]$ are the year of the survey described as a covariate and a factor, respectively. The $\beta_{1,j}$ describes long-term temporal change in log-length across locations, which is of specific interest, and the $\alpha_{\text{year}[i]}$ describe stochastic year-to-year changes in length common across locations due to unknown environmental factors. The binning of observed fish lengths was treated in the same way as in spatial analysis. Bayesian methods were again used to estimate the model parameters. Uniform priors were used for all parameters: $[1-4]$ for $\beta_{0,j}$, $[-0.1-0.1]$ for $\beta_{1,j}$, $[0.001-0.5]$ for σ_{year} and σ_{survey} , and $[0.1-0.5]$ for σ_{lnlength} . These priors had little influence on the posteriors.

Five of the long-term locations included observations both inside and outside of no-take marine protected areas (MPA) and we also tested whether temporal trends in body size differed between sites inside and outside MPAs. This was done to assess whether harvesting could be driving temporal trends in observed body sizes. The MPAs are officially no-take areas, and even though small amount of illegal harvesting is likely to occur, the fishing rates are more intense outside MPA areas. The two datasets had 84 and 74 species, respectively, with 69 species found with sufficient abundance both inside and outside MPAs. The overall average body size trends through time were similar inside and outside MPAs (Supplementary Information Fig. 2), suggesting that temporal body size trends were not driven by exploitation. This was not surprising, given that majority of the species analysed (after the filtering described above) are not routinely targeted by commercial or recreational fishing. Final temporal analyses were therefore done on a combined dataset ignoring the MPA effects (105 species across 9 locations).

To compare rates of body size change across space and time we explored correlation between the body size responses in spatial analyses (β_1) and annual body size trends ($\beta_{1,j}$) at eight locations with significant warming trends. Out of 105 species included in the temporal analyses, 28 species were found only in one non-warming location (Jurien Bay) and were not included in the space-time rate comparison. From the remaining 77 species, six species did not have sufficient spatial data, so the final dataset for the space-time rate comparison included 71 species. For these species correlations between spatial and temporal slopes were assessed using bootstrapped weighted Spearman's correlation (10000 bootstrap replicates), where each value was weighted by the inverse of its uncertainty range (code available at <https://github.com/astaudzi/RLSfishSize>).

Species abundance and therefore potential density dependence effects were not included in the main model, because the relationship describing potential abundance effects on body size are unlikely to be linear, and because competition is likely to occur at both intra- and inter-specific level.

Accounting for all these effects would require a very complex, parameter-rich model and much larger datasets that was available for most species. However, to assess whether density dependence could still be the major driver of the observed body trends (if abundance was determined by SST) we compared the species-specific spatial body size slopes (β_1) with abundance-SST skew statistics assessed in ²³. The skew statistics describe relative abundances of a species across its distribution range, where positive skew means that a species is more abundant at the warmer half of its distribution and vice versa. If abundance was a major driver of average body size, we could expect an overall negative correlation between size-temperature and abundance-temperature slopes, such as mean body size is smaller at sites with higher abundances. Alternatively, larger average body sizes might have positive effects on abundance through e.g. improved recruitment or inter-specific effects. In this case we could expect a positive correlation between size-temperature and abundance-temperature slopes. Estimates of both size-temperature slopes (this study) and abundance-temperature skew ²³ were available for 300 species. The correlation between body size slopes and abundances was close to zero ($r = 0.09$, $p = 0.14$), suggesting that abundance is unlikely to be a major driver of body sizes and vice versa (Extended Data Fig. 2).

All statistical analyses were performed using R v.3.2, Bayesian analyses were done using the *rstan* package³⁴.

Data availability. Underwater visual survey datasets are available through the Reef Life Survey site www.reeflifesurvey.com. The final datasets and all codes used in this analysis are available through the code depository at <https://github.com/astaaudzi/RLSfishSize> and as Supplementary datasets linked to this article.

References

- 1 Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. & West, G. B. Toward a metabolic theory of ecology. *Ecology* **85**, 1771-1789 (2004).

387 2 Blanchard, J. L., Heneghan, R. F., Everett, J. D., Trebilco, R. & Richardson, A. J. From
388 bacteria to whales: using functional size spectra to model marine ecosystems. *Trends in*
389 *ecology & evolution* **32**, 174-186 (2017).

390 3 Fisher, J. A., Frank, K. T. & Leggett, W. C. Breaking Bergmann's rule: truncation of
391 Northwest Atlantic marine fish body sizes. *Ecology* **91**, 2499-2505 (2010).

392 4 Shackell, N. L., Frank, K. T., Fisher, J. A., Petrie, B. & Leggett, W. C. Decline in top
393 predator body size and changing climate alter trophic structure in an oceanic ecosystem.
394 *Proceedings of the Royal Society of London B: Biological Sciences*, rspb20091020 (2009).

395 5 Rouyer, T., Sadykov, A., Ohlberger, J. & Stenseth, N. C. Does increasing mortality change
396 the response of fish populations to environmental fluctuations? *Ecology letters* **15**, 658-665
397 (2012).

398 6 Gardner, J. L., Peters, A., Kearney, M. R., Joseph, L. & Heinsohn, R. Declining body size: a
399 third universal response to warming? *Trends in ecology & evolution* **26**, 285-291 (2011).

400 7 Cheung, W. W. *et al.* Shrinking of fishes exacerbates impacts of global ocean changes on
401 marine ecosystems. *Nature Climate Change* **3**, 254 (2013).

402 8 Pauly, D. & Cheung, W. W. L. Sound physiological knowledge and principles in modeling
403 shrinking of fishes under climate change. *Glob Chang Biol* **24**, e15-e26,
404 doi:10.1111/gcb.13831 (2018).

405 9 Lefevre, S., McKenzie, D. J. & Nilsson, G. E. In modelling effects of global warming, invalid
406 assumptions lead to unrealistic projections. *Global change biology* **24**, 553-556 (2018).

407 10 Morrongiello, J. R., Sweetman, P. C. & Thresher, R. E. Fishing constrains phenotypic
408 responses of marine fish to climate variability. *Journal of Animal Ecology* (2019).

409 11 Angilletta Jr, M. J., Steury, T. D. & Sears, M. W. Temperature, growth rate, and body size in
410 ectotherms: fitting pieces of a life-history puzzle. *Integrative and comparative biology* **44**,
411 498-509 (2004).

412 12 Bergman, C. Uber die Verhaltnisse der Warmeokonomie der Thiere zu ihrer Grosse.
413 *Gottinger Studien* **3**, 595-708 (1847).

414 13 Audzijonyte, A. *et al.* Is oxygen limitation in warming waters a valid mechanism to explain
415 decreased body sizes in aquatic ectotherms? *Global Ecology and Biogeography* **28**, 64-77
416 (2019).

417 14 Forster, J., Hirst, A. G. & Atkinson, D. Warming-induced reductions in body size are greater
418 in aquatic than terrestrial species. *Proceedings of the National Academy of Sciences* **109**,
419 19310-19314, doi:10.1073/pnas.1210460109 (2012).

- 15 Lefevre, S., McKenzie, D. J. & Nilsson, G. E. Models projecting the fate of fish populations
under climate change need to be based on valid physiological mechanisms. *Glob Chang Biol*
23, 3449-3459, doi:10.1111/gcb.13652 (2017).
- 16 Pauly, D. The relationships between gill surface area and growth performance in fish: a
generalization of von Bertalanffy's theory of growth. *Meeresforschung* **28**, 251-282 (1981).
- 17 Donelson, J. M., Munday, P. L., McCormick, M. I. & Pitcher, C. R. Rapid transgenerational
acclimation of a tropical reef fish to climate change. *Nature Climate Change* **2**, 30-32,
doi:10.1038/nclimate1323 (2011).
- 18 Pecl, G. T. *et al.* Biodiversity redistribution under climate change: Impacts on ecosystems
and human well-being. *Science* **355**, eaai9214 (2017).
- 19 Edgar, G. J. & Barrett, N. S. An assessment of population responses of common inshore
fishes and invertebrates following declaration of five Australian marine protected areas.
Environmental Conservation **39**, 271-281 (2012).
- 20 Edgar, G. J. & Stuart-Smith, R. D. Systematic global assessment of reef fish communities by
the Reef Life Survey program. *Scientific Data* **1**, 140007 (2014).
- 21 Stuart-Smith, R. D., Edgar, G. J., Barrett, N. S., Kininmonth, S. J. & Bates, A. E. Thermal
biases and vulnerability to warming in the world's marine fauna. *Nature* **528**, 88 (2015).
- 22 Edgar, G. J., Barrett, N. S. & Morton, A. J. Biases associated with the use of underwater
visual census techniques to quantify the density and size-structure of fish populations.
Journal of Experimental Marine Biology and Ecology **308**, 269-290 (2004).
- 23 Waldoock, C., Stuart-Smith, R. D., Edgar, G. J., Bird, T. J. & Bates, A. E. The shape of
abundance distributions across temperature gradients in reef fishes. *Ecology letters* **22**, 685-
696 (2019).
- 24 Bennett, S., Wernberg, T., Joy, B. A., De Bettignies, T. & Campbell, A. H. Central and rear-
edge populations can be equally vulnerable to warming. *Nature Communications* **6**, 10280
(2015).
- 25 Conover, D. O., Duffy, T. A. & Hice, L. A. The covariance between genetic and
environmental influences across ecological gradients: reassessing the evolutionary
significance of countergradient and cgradient variation. *Ann N Y Acad Sci* **1168**, 100-129,
doi:10.1111/j.1749-6632.2009.04575.x (2009).
- 26 Audzijonyte, A., Kuparinen, A. & Fulton, E. A. How fast is fisheries-induced evolution?
Quantitative analysis of modelling and empirical studies. *Evolutionary applications* **6**, 585-
595 (2013).

- 27 Audzijonyte, A. *et al.* Trends and management implications of human-influenced life history changes in marine ectotherms. *Fish and Fisheries* **17**, 1005-1028 (2016).
- 28 Fritschie, K. J. & Olden, J. D. Disentangling the influences of mean body size and size structure on ecosystem functioning: An example of nutrient recycling by a non-native crayfish. *Ecology and evolution* **6**, 159-169 (2016).
- 29 Audzijonyte, A., Kuparinen, A., Gorton, R. & Fulton, E. A. Ecological consequences of body size decline in harvested fish species: positive feedback loops in trophic interactions amplify human impact. *Biology Letters* **9**, 20121103, doi:doi:10.1098/rsbl.2012.1103 (2013).
- 30 Jørgensen, C. & Fiksen, Ø. Modelling fishing-induced adaptations and consequences for natural mortality. *Canadian Journal of Fisheries and Aquatic Sciences* **67**, 1086-1097 (2010).
- 31 Edgar, G. J. & Stuart-Smith, R. D. Ecological effects of marine protected areas on rocky reef communities—a continental-scale analysis. *Marine Ecology Progress Series* **388**, 51-62 (2009).
- 32 Reynolds, R. & Banzon, V. NOAA Optimum Interpolation 1/4 Degree Daily Sea Surface Temperature (OISST) Analysis, Version 2. *NOAA National Centers for Environmental Information*. doi **10**, V5SQ8XB5 (2008).
- 33 Stuart-Smith, R. D., Edgar, G. J. & Bates, A. E. Thermal limits to the geographic distributions of shallow-water marine species. *Nature ecology & evolution* **1**, 1846 (2017).
- 34 Stan Development Team. RStan: the R interface to Stan. R package version 2.19.2. <http://mc-stan.org/>. (2019).

Reporting summary. Further information on experimental design is available in the Nature Research Reporting Summary linked to this paper.

Acknowledgements

The authors wish to acknowledge all divers and support personnel who have contributed to the data collection and repository, especially Antonia Cooper, Just Berkhout, Elizabeth Oh, Jemina Stuart-Smith, Justin Hulls and Ella Clausius. We also thank Michael Sumner for help with the SST data, Freddie Heather for help with figures, Just Berkhout for help with parallel computing and Amanda Bates and Conor Waldock for thermal distribution information. This research used the NCRIS-enabled Integrated Marine Observing System (IMOS) infrastructure for database support and storage.

This study was supported by the ARC Discovery grant DP170104240 (to JLB, GP and RDSS), UTAS visiting scholars fellowship (to AA for NP visit to IMAS), ARC Future Fellowship to GP, and the Marine Biodiversity Hub, a collaborative partnership supported through the Australian Government's National Environmental Science Programme (NESP).

Author contributions

AA designed the study and led the writing. AA and SAR designed and conducted the statistical analyses. RDS-S, GJE and NSB led field surveys and data collection. GP, NP and JLB contributed critically to the development of the study. All authors contributed to the manuscript writing and gave final approval for publication.



Author information


The authors declare no competing interests. Reprints and permissions information is available at www.nature.com/reprints. Correspondence should be addressed to asta.audzijonute@utas.edu.au

Figure legends


Fig. 1. Relative change in the mean body length of 335 coastal marine fish species per one degree change in SST observed across their geographic distributions. Change varies with species' thermal affinity²¹ (A) and maximum observed body length (B). Dots and vertical bars represent the median and 80% PPD ranges of individual species responses, respectively (wide intervals that fall outside the picture limits are not shown). Species that show clear body size responses to SST (90% PPD range above or below zero) are shown in red and blue colours. Solid black line is the linear regression weighted according to the individual species uncertainty level. Orange shading depicts the 95% credible interval for the regression.


510 **Fig. 2.** Long-term annual relative change in mean body length of 71 species at eight warming
511 locations. Change through time is positively correlated with the change in mean body length per 1°C
512 SST across their geographic distribution (see Fig. 1). The orange line shows the case where spatial
513 and temporal trends are perfectly correlated but change in body length per 1°C of warming through
514 time is 10 times faster than change per 1°C of warming through space (see text for details).
515 Correlation between spatial and temporal responses was calculated using bootstrapped weighted
516 Spearman's correlation (ρ).





Mean annual SST of each geographic cell (12 to 29)





Annual change in body length in 9 long term locations

